

GEOMETRIC VERTEX OPERATORS

ILYA ZAKHAREVICH

ABSTRACT. We investigate relationships between (infinite-dimensional) algebraic geometry of loop spaces, smooth families of partial linear operators, and classical vertex operators. Vertex operators are shown to be actions of birational transformations of infinite-dimensional algebraic “varieties” M on appropriate line bundles on M .

These vertex operators act in the vector space $\mathcal{M}(M)$ of meromorphic functions on M as *partial operators*: they are defined on a subspace (in an appropriate *lattice of subspaces* of $\mathcal{M}(M)$), and send smooth families in such a subspace to smooth families. Axiomatizing this, we define *conformal fields* as arbitrary families of partial operators in $\mathcal{M}(M)$ which satisfy both these properties.

The “variety” M related to standard vertex operators is formed by rational functions of one variable $z \in Z = \mathbb{P}^1$, changing the variety Z one obtains different examples of M , and multidimensional analogues of vertex operators. One can cover $\mathcal{M}(M)$ by “big smooth subsets”; these subsets are parameterized by appropriate projective bundles over Hilbert schemes of points on Z . We deduce conformal associativity relation for conformal fields, and conformal commutation relations for Laurent coefficients of commuting conformal fields from geometric properties of the Hilbert schemes.

One of the main stumbling blocks for a mathematician when dealing with conformal fields is the very first sentence they see: “Consider two holomorphic functions $a(s)$ and $b(t)$ such that $a(s)b(t)$ has a pole when $s = t$.” We *start* with examples of such functions (with values in partially defined operators in spaces of meromorphic functions), thus make this problem nonextant.

We also provide geometric description of boson-fermion correspondence, and relations of this correspondence to geometry of the set of meromorphic functions.

CONTENTS

0. Introduction	3
0.1. Loops spaces and OPEs	3
0.2. OPEs and linear algebra	4
0.3. Finite dimensional algebraic geometry	5
0.4. Schemes and loop spaces	5
0.5. Geometry of classical vertex operators and moving poles	6
0.6. OPEs for classical vertex operators	8

Date: May 1998 (Revision II: August 2002), archived at [arXiv:math.AG/980509](https://arxiv.org/abs/math/980509). Printed: February 8, 2008.

0.7.	Locality condition and singularities	9
0.8.	Skyscraper variations	10
0.9.	Birational geometry and partial operators	10
0.10.	Flavors of the theories	11
0.11.	On $\text{Func}(\mathcal{L}^r X)$	13
0.12.	Partial operators in $\text{NaiveFunc}(\mathcal{L}^r X)$	14
0.13.	Basic and generalized functions	15
0.14.	The plan	16
0.15.	Alternative approaches	17
0.16.	Arithmetic analogue	18
1.	Field of meromorphic functions on $\mathcal{M}(Z)$	18
1.1.	Meromorphic functions	18
1.2.	Families of meromorphic functions on $\mathcal{M}(Z)$	21
1.3.	Action of automorphisms of $\mathcal{M}(Z)$	23
1.4.	Multiplication in the field, smooth families, and conformal operators	24
1.5.	Families of operators $\mathbf{M}_\eta$	25
1.6.	Conformal fields	26
1.7.	Conformal field theory	27
1.8.	Homogeneous variant	28
1.9.	Residues	28
1.10.	Fusions	32
1.11.	Localization	33
2.	Examples of conformal families	35
2.1.	Fields based on $\mathbb{P}^1$	35
2.2.	Fusions of $\mathbf{F}_\eta$	37
2.3.	A different reduction	38
2.4.	Further reductions	39
2.5.	Vertex operators	42
2.6.	Analogy with flag space	44
3.	Hilbert schemes and codominant conformal fields	44
3.1.	Big families	44
3.2.	Versality	48
3.3.	V -conformal operators	51
3.4.	Codominant families	53
3.5.	Conformal algebras	57
3.6.	Fractions of formal series	58
3.7.	Laurent coefficients of conformal fields	60
3.8.	Laurent coefficients of a composition	61
4.	Chiral algebras and commutation relations	64
4.1.	Chiral algebras	65
4.2.	Conformal commutation relations	66
4.3.	The family $\mathcal{G}$	68

4.4. Generators X_{ij}	69
5. Boson-fermion correspondence	71
5.1. Bosonic space	71
5.2. Boson creation operator	73
5.3. Naive boson-fermion correspondence	73
5.4. Fermion creation operator	74
5.5. Fock space I	74
5.6. Fock space II	76
5.7. Bosons: geometric <i>vs</i> algebraic	77
References	79

0. INTRODUCTION

All the geometry and linear algebra considered in this paper happens over an arbitrary algebraically closed field $\mathbb{K}$. However, the main application is when $\mathbb{K} = \mathbb{C}$, thus one can feel free to substitute $\mathbb{C}$ instead of $\mathbb{K}$.

Note however that in a case of a topological field (such as $\mathbb{C}$) our weak axioms will not work in the case of an analytic manifold, to implement field theory over an analytic manifold one may need much more rigid axioms. We confine ourself to the case of algebraic manifolds and algebraic functions, so one does not need to worry about significant singularities and non-dense open sets.

0.1. Loops spaces and OPEs. Fix a variety Z . Given a variety X , a *loop in* X is a mapping $\varphi: Z \rightarrow X$ (smooth, or regular, or formal, or rational, etc., as specified by context). Denote the set of loops on X by $\mathcal{L}X$ or $\mathcal{L}_Z X$; later we equip this set with additional structures, and call it the *loop space*. Assume that these additional structures on $\mathcal{L}X$ lead to a particular algebra $\text{Func}(\mathcal{L}X)$ of functions on $\mathcal{L}X$ (continuous, or regular, or rational, etc., as needed). To avoid confusion with the loops (which are functions themselves), call elements of $\text{Func}(\mathcal{L}X)$ *functionals on* $\mathcal{L}X$.

Loops spaces are the first examples of infinite-dimensional varieties with properties dramatically different from those familiar from finite-dimensional algebraic geometry. *Vertex* flavors of algebro-geometric constructions are convenient tools helping handling objects from this still-unchartered territory.

The simplest manifestation of this new geometry is the appearance of *operator product expansions (OPEs)* in the rule of composition of families of geometric linear operators acting on $\text{Func}(\mathcal{L}X)$. Recall that given a manifold M , a *geometric operator* in $\text{Func}(M)$ is a composition of an operator $\mathbf{F}$ of multiplication by $F \in \text{Func}(M)$ and of the operator h^* for a diffeomorphism $h \in \text{Diff}(M)$ of M . Recall also that given two smooth families A_t , $t \in T$, and B_s , $s \in S$, of linear operators acting in a vector

space V , the product family $A_t B_s$ is *not necessarily*¹ smooth on the whole parameter space $T \times S$. We say that these families have an *operator product expansions* if the product is smooth outside of a hypersurface $\Sigma \subset T \times S$; moreover, $A_t B_s$ has *at most a pole* at Σ ; in other words, there is a number $k \in \mathbb{Z}$ such that the product $\Phi^k(t, s) A_t B_s$ allows a smooth continuation from $T \times S \setminus \Sigma$ to $T \times S$; here Φ is defined by $\Sigma = \{\Phi(t, s) = 0\}$.

0.2. OPEs and linear algebra. The aim of this introduction is to analyze the relationship of OPEs with infinite-dimensional geometry of loop spaces in more detail. The bulk of this paper develops a linear algebra framework in which “good” families of linear operators in a vector space V *automatically* satisfy OPEs. It turns out that the notion of “goodness” can be expressed in terms of two compatible structures on V : on the notion of *smoothness* on V (i.e., a description of which mappings $M \rightarrow V$ are “smooth”; here M is an arbitrary manifold), and on a *lattice*² of vector subspaces in V parameterized by a union of algebraic varieties.

The principal results of this paper claim that

- (1) it makes sense to consider the set of functionals on (an appropriate flavor of) the loop space; this set is a vector space (even an algebra) V ;
- (2) this vector space V carries a smoothness structure and a compatible lattice of vector subspaces;
- (3) it makes sense to consider smooth families of (partial³) linear operators in V which are compatible with two structures on V introduced above;
- (4) any two such families satisfy an OPE;
- (5) the usual “physical parlance” lexicon and toolset of conformal fields is *literally* applicable to the families of the above form;
- (6) the “standard” families of linear operators appearing in the theory of vertex operators act in a vector subspace of V ; they can be “continuously” extended to the families of partial operators in V compatible with two structures on V ;
- (7) moreover, these “standard vertex operators” have a geometric nature, corresponding to the action of certain groups in certain line bundles on the loop space.

The corresponding flavor of the loop space is the space $\mathcal{L}^r X$ of *rational loops*, i.e., of rational mappings $Z \rightarrow X$. (Here we assume that Z is a projective manifold.)

This introduction serves a special purpose: in the principal part of this paper we mostly ignore the last part of the program above, and introduce the vector space V in

¹It would be smooth if the multiplication mapping $\text{End}(V)^2 \rightarrow \text{End}(V)$ were continuous (in an appropriate sense). However, it is not, with a few exceptions, such as $\dim V < \infty$, or the case of Banach V . Later, in Section 0.9, we discuss the actual mechanism of appearance of singularities; however, right now it is enough to understand that there is no *a priori* reason to assume $A_t B_s$ to be smooth.

²I.e., a collection of vector subspaces of V closed w.r.t. finite sums and intersections.

³In other words, linear operators $V_0 \rightarrow V$, here $V_0 \subset V$.

purely algebraic terms. Here, in the introduction, we discuss in details the geometric realization of V . Given that the corresponding infinite-dimensional geometry is quite hairy, sometimes the discussion becomes superficial, e.g., providing only circumstantial evidence, parables and analogies. We are going to discuss the the geometry in details in [12].

0.3. Finite dimensional algebraic geometry. The geometry of an affine finite-dimensional algebraic variety M (possibly equipped with an action of a group G) is encoded in commutator rules for geometric linear operators (defined in Section 0.1) in $\text{Func}(M)$. First of all, the commutative algebra $A = \text{Func}(M)$ of functions on M (the corresponding commutator rule is $aa' = a'a$ for $a, a' \in A$) determines M . Moreover, given a group G of symmetries of M , the algebra $\tilde{A} = k[G] \ltimes A$ acts on $\text{Func}(M)$ by geometric operators. The structure of the group G is encoded into the generators and relations of $k[G]$; the structure of the action is encoded in relationships of the form $ga = a'g$; here $a \in A$ and $g \in G$ are given, and a' is an appropriate element of A . In other words, given a pair $\tilde{A} \supset A$ of algebras with commutation relations as described above, one can reconstruct M and G .

In a non-affine situation an extra ingredient is added; the *locality* along M enters the play via consideration of locally-defined functions. The action of G mixes the domains of definition; the situation may be simplified again in the quasiprojective case, when one can replace $A = \text{Func}(M)$ with the algebra $A = \text{MerFunc}(M)$ of meromorphic (alternatively named *rational*) functions on M . After such a translation G acts again on the algebra A .

Suppose that the action π of G on M is extended to an action Π on a line bundle $\mathcal{L}$ over M . Trivialize $\mathcal{L}$ over a Zariski open subset $U \subset M$; this associates a meromorphic function a_σ on U to any locally defined section σ of $\mathcal{L}$. Now the action Π of G on locally defined (or meromorphic) sections of $\mathcal{L}$ is a geometric operator, $\Pi_g a = \mu_g \cdot \pi_g^* a$; here μ_g is a meromorphic function on M (a *Schur multiplier*) describing how the action of G on $\mathcal{L}$ changes the trivialization of $\mathcal{L}$.

In other words, the action of $g \in G$ on the vector space of sections of $\mathcal{L}$ can be expressed in terms of operators of two types: the action of diffeomorphisms of M on (meromorphic) functions of M , and the operator of multiplication by a (meromorphic) function on M . In the introduction we discuss analogues of these operators when M is a loop space.

0.4. Schemes and loop spaces. Note that the appearance of meromorphic functions on M is an artifact of our usage of a trivialization of $\mathcal{L}$. However, in the case of loop spaces this is going to be an immanent feature: it turns out that the “manifolds” $M = \mathcal{L}^r X$ (consisting of *rational* loops on X) has no regular functions, even locally defined regular functions:

Example 0.1. Suppose $Z = \mathbb{P}^1$, $X = \mathbb{A}^1$. Let $R_k \subset \mathcal{L}^r X$ consists of rational loops of degree k . One can parameterize R_1 via $\varphi_P(z) = \frac{az+b}{cz+d}$; here $P = (a : b : c : d) \in \mathbb{P}^3$.

On the other hand, the loops φ_P with $P = (\alpha\beta : \alpha'\beta : \alpha\beta' : \alpha'\beta')$ are just constant loops (from R_0) with the value $(\beta : \beta') \in \mathbb{P}^1$. In other words, R_1 is the image of $\mathbb{P}^3$ with all the projective lines $(\alpha B : \alpha' B : \alpha B' : \alpha' B')$ shrunk to a point (here B, B' are given numbers, and $(\alpha : \alpha') \in \mathbb{P}^1$ is a parameter on the line).

On the other hand, it is obvious that $\mathbb{P}^3$ with a line l shrunk to a point has few locally defined functions; indeed, let $L \in Q$ be the image of the line l on the quotient space Q ; consider a neighborhood U of L . Its preimage U' on $\mathbb{P}^3$ contains l , thus it contains any projective line l' which is close to l ; in particular, any regular function on U' is constant, since its pull back must be constant on any (projective!) line l' .

Similarly, R_1 can have no regular functions in the neighborhood of $R_0 \subset R_1$; same for a neighborhood of R_k in R_{k+1} . Since any point of $\mathcal{L}^r X$ is in R_k for a large enough k , this shows that $\mathcal{L}^r X$ can have no regular function.

Remark 0.2. It turns out that the principal reason for this phenomenon is that in the example we allow a “moving singularity”: the position of the pole of the loop $\varphi_P(z)$ depends on the parameter P ; however, we assume that the family $\{\varphi_P\}_{P \in \mathbb{P}^3}$ is a “smooth family” in $\mathcal{L}^r X$. In turn, the reason for such an assumption (and the reason to consider $\mathcal{L}^r X$!) is the observation (see Sections 0.5 and 0.6) that the “classical vertex operators” $\exp \sum_{n>0} \frac{s^{-n}}{n} \partial_n$ and $\sum_{n \geq 0} s^{-n-1} \partial_n$ “create” a pole of a loop at a moving point (actually at the point $s \in Z = \mathbb{P}^1$).

The example above shows that it does not make a lot of sense to describe the “space” $\mathcal{L}^r X$ (as well as the action of vertex operators) using the language of schemes; indeed, this language is based on the idea that the locally defined functions separate points. This may explain why the scheme-theoretic approaches of [6] and [9] to vertex algebras and loop spaces lead to so complicated constructions.

0.5. Geometry of classical vertex operators and moving poles. In a lot of applications of families of linear operators with OPEs, the families are given by explicit formulae involving the following series depending on parameters t and s :

$$\begin{aligned} & \exp \sum_{n \geq 0} t^n \mathbf{y}_n, & \exp \sum_{n > 0} \frac{s^{-n}}{n} \partial_n \\ & \sum_{n \geq 0} t^n \mathbf{x}_n, & \sum_{n \in \mathbb{Z}} t^n \mathbf{x}_n, & \sum_{n \geq 0} s^{-n-1} \partial_n, & \sum_{n \in \mathbb{Z}} s^{-n-1} \partial_n \end{aligned}$$

Here x_n (or y_n) are free generators of $A = \mathbb{C}[x_n]_{n \in \mathbb{Z}}$ or $A = \mathbb{C}[y_n]_{n \geq 0}$ correspondingly, and $\mathbf{x}_n$ (or $\mathbf{y}_n$) are operators in A of multiplication by the corresponding symbol x_n (or y_n); ∂_n is the operator of taking the derivative w.r.t. the corresponding symbol x_n or y_n . To show how these operators are related to loop spaces and moving singularities of meromorphic loops, consider A as an algebra of functions on a particular loop space of the additive (for operators with x_n) or multiplicative (for operators with y_n) group.

The flavor of a loop space on which this relation is simplest to illustrate corresponds to Z being a punctured disk, $Z = \text{PDisk}(R) \stackrel{\text{def}}{=} \{0 < |z| < R\} \subset \mathbb{C}$. First, consider the additive group $X = \mathbb{A}^1$. Then any loop $\varphi \in \mathcal{L}X$ can be written as $\varphi(z) = \sum_{n \in \mathbb{Z}} x_n z^n$; this gives a coordinate system (x_n) on $\mathcal{L}X$. Consider any “reasonable” definition of $\text{Func}(\mathcal{L}X)$; then the evaluation functional $\text{Eval}_t: \varphi \mapsto \varphi(t)$ must be an element of $\text{Func}(\mathcal{L}X)$ for $|t| < R$. Denote by $\mathbf{E}_t$ the operator in $\text{Func}(\mathcal{L}X)$ of multiplication by Eval_t . Similarly, consider the mapping

$$G_s^{(\varepsilon)}: \mathcal{L}\mathbb{A}^1 \rightarrow \mathcal{L}\mathbb{A}^1: \varphi \rightarrow \varphi', \quad \varphi'(z) = \varphi(z) + \frac{\varepsilon}{z-s}.$$

These mappings form a well-defined one-parameter group of automorphisms of $\mathcal{L}\mathbb{A}^1$ for $|s| > R$; denote by $\mathbf{d}_s$ the corresponding vector field on $\mathcal{L}\mathbb{A}^1$. Again, for any reasonable definition of $\text{Func}(\mathcal{L}\mathbb{A}^1)$, this vector field induces a linear operator on $\text{Func}(\mathcal{L}\mathbb{A}^1)$; denote it by the same symbol.

In the coordinates x_n one can write the function Eval_t on $\mathcal{L}\mathbb{A}^1$ as $\sum_{n \in \mathbb{Z}} t^n x_n$. Moreover, since the vector field $\mathbf{d}_s$ is translation-invariant on $\mathcal{L}\mathbb{A}^1$, it must be a linear combination of the operators ∂_n . Writing the Taylor coefficients of $1/(z-s)$, one obtains

Theorem 0.3. *The operators $\mathbf{E}_t$ and $\mathbf{d}_s$ can be written as*

$$\mathbf{E}_t = \sum_{n \in \mathbb{Z}} t^n \mathbf{x}_n, \quad \mathbf{d}_s = - \sum_{n \geq 0} s^{-n-1} \partial_n.$$

here $\mathbf{E}_t$ is a well-defined⁴ linear operator in $\text{Func}(\mathcal{L}_{\text{PDisk}(R)}\mathbb{A}^1)$ for $0 < |t| < R$; $\mathbf{d}_s$ is a well-defined linear operator in $\text{Func}(\mathcal{L}_{\text{PDisk}(R)}\mathbb{A}^1)$ for $|s| \geq R$.

One can also consider a non-punctured disk $\text{Disk}(R)$ in similar terms. In this case the coordinate systems is $(x_n)_{n \geq 0}$, and the sum for the operator $\mathbf{E}_t$ has only the terms for $n \geq 0$.

Now consider a similar loop space $\{\varphi: \text{Disk}(R) \rightarrow \mathbb{C}^*\}$ of the multiplicative group. Consider a coordinate system $(y_n)_{n \geq 0}$ on $\mathcal{L}\mathbb{C}^*$ by writing $\varphi(z) = y_0 \exp \sum_{n > 0} y_n z^n$. Abusing notations, denote ∂_{y_n} as ∂_n .

As above, consider the operator $\mathbf{E}_t$; consider also the mapping

$$H_s: \mathcal{L}\mathbb{C}^* \rightarrow \mathcal{L}\mathbb{C}^*: \varphi \rightarrow \varphi', \quad \varphi'(z) = \frac{s\varphi(z)}{s-z}.$$

This mapping is well-defined for $|s| > R$; denote by $\mathbf{H}_s$ the corresponding operator on $\text{Func}(\mathcal{L}\mathbb{C}^*)$. Again, since H_s preserves y_0 and is a parallel translation in coordinates y_n , $n > 0$, one can write $\mathbf{H}_s$ as $\exp(v)$; here v is a translation-invariant vector field in coordinates y_n . The calculation of the Taylor series for $\log \frac{s}{s-z}$ gives

⁴Since these sums are infinite, one needs to describe the topology on $\text{Func}(\mathcal{L}_{\text{PDisk}(R)}\mathbb{A}^1)$ first. However, this statement holds with all the natural choices of topology on this space, so we skip this issue now.

Theorem 0.4. *The operators $\mathbf{E}_t$ and $\mathbf{H}_s$ can be written as*

$$(0.1) \quad \mathbf{E}_t = \mathbf{y}_0 \exp \sum_{n>0} t^n \mathbf{y}_n, \quad \mathbf{H}_s = \exp \sum_{n>0} \frac{s^{-n}}{n} \partial_n;$$

here $\mathbf{E}_t$ is a well-defined linear operator in $\text{Func}(\mathcal{L}_{\text{Disk}(R)}\mathbb{C}^*)$ for $|t| < R$; $\mathbf{H}_s$ is a well-defined linear operator in $\text{Func}(\mathcal{L}_{\text{Disk}(R)}\mathbb{C}^*)$ for $|s| \geq R$.

Remark 0.5. For the mapping H_s the point $0 \in \text{Disk}(R)$ plays a special role. However, one can consider the mapping $h_s: \varphi \mapsto \tilde{\varphi}$, $\tilde{\varphi}(z) = \varphi(z)/(z-s)$, instead of H_s . The corresponding linear operator $\mathbf{h}_s$ is just $(-s)^{y_0 \partial_0} \mathbf{H}_s$; hence it can be written by a formula similar to (0.5.10).

Remark 0.6. It is clear that the operators above should be written by the same formulae no matter which flavor of $\mathcal{L}X$ (here $X = \mathbb{C}$ or $X = \mathbb{C}^*$) and which flavor of the space $\text{Func}(\mathcal{L}X)$ of functional on $\mathcal{L}X$ one considers. Indeed, it is enough to check the equality of the right-hand side and the left-hand side on a “dense subset” of $\text{Func}(\mathcal{L}X)$, and this dense subset can be chosen independently of the flavor.

0.6. OPEs for classical vertex operators. Consider a manifold M ; denote by μ_F the operator in $\text{Func}(M)$ of multiplication by a function $F \in \text{Func}(M)$; consider the operator $\mathbf{V}$ in $\text{Func}(M)$ induced by a vector field V on M . Then, obviously, $[\mathbf{V}, \mu_F] = \mu_{V \cdot F}$. Hence the operators $\mathbf{d}_s$ and $\mathbf{E}_t$ have the commutation relation $[\mathbf{d}_s, \mathbf{E}_t] = \frac{1}{s-t} \mathbf{E}_t$. This exposes the first OPE-like relationship of this paper. Similarly, $\mathbf{h}_s \mathbf{E}_t = \frac{1}{t-s} \mathbf{E}_t \mathbf{h}_s$. Since these expressions are not exactly of the OPE form, comment on their relations to OPEs.

In the former case the right-hand side depends on s via the denominator only; thus the left-hand side *must* have a pole at $s = t$. In the latter case $\mathbf{E}_t \mathbf{h}_s$ sends the constant functional 1 on $\mathcal{L}X$ to Eval_t ; hence $\mathbf{E}_t \mathbf{h}_s$ has no zero at $t = s$. Thus, again, $\mathbf{h}_s \mathbf{E}_t$ *must* have a singularity of an OPE type when $t = s$.

However, these poles are impossible to see when one considers the vector space of functionals on analytic flavors of loops, such as $\text{Func}(\mathcal{L}_{\text{Disk}(R)}\mathbb{C}^*)$. Indeed, the domains of definition $\{|s| > R\}$ and $\{|t| < R\}$ of the families in question do not intersect, thus the divisor of poles (the diagonal) does not intersect the product of domains of two families (e.g., $\mathbf{E}_t$ and $\mathbf{d}_s$). Moreover, in the discussion above we did not completely specify the nature of families $\mathbf{E}_t$ etc.: all we defined was $\mathcal{L}_{\text{PDisk}(R)}\mathbb{A}^1$ etc., while we also need to specify $\text{Func}(\mathcal{L}_{\text{PDisk}(R)}\mathbb{A}^1)$ and which families of operators in $\mathcal{L}_{\text{PDisk}(R)}\mathbb{A}^1$ are smooth families. Until we define these notions, we can only discuss the properties of commutators, $[\mathbf{d}_s, \mathbf{E}_t]$ and $\mathbf{h}_s \mathbf{E}_t \mathbf{h}_s^{-1}$, not the properties of products of elements of the family, as the OPEs must.

Indeed, the crucial operation in the theory of vertex operators is the calculation of the *fusions*; fusions are the Laurent coefficients of the composition $A_s \circ B_t$ near the divisor of poles. For a lot of calculations it is convenient to write these Laurent coefficients as the corresponding Cauchy integrals. To be able to do this, one needs

to have $A_s \circ B_t$ defined in a neighborhood of the divisor of poles; it is not enough to have it defined in “one half-space” near this divisor; however, this is the situation with the domain $\{(s, t) \mid |s| > R, |t| < R\}$.

Later, when we switch to the different flavor $\mathcal{L}X = \mathcal{L}^r X$ of loop spaces, we will be able to define the necessary notions (the vector space $\text{Func}(\mathcal{L}X)$ and the notion of smooth families of operators on $\text{Func}(\mathcal{L}X)$) in such a way that the domains of the families $\mathbf{E}_t$ and $\mathbf{h}_s$ will intersect. We will see, for example, that for this flavor $\mathbf{E}_t \circ \mathbf{h}_s$ is a smooth family; thus $\mathbf{h}_s \circ \mathbf{E}_t$ is going to have a pole of the first order when $s = t$.

Anyway, a theory of loop spaces which allows the consideration of OPEs for classical vertex operators must allow the consideration of the dependence of the family of mappings h_s on the parameter s , similarly for the vector fields $\mathbf{d}_s$ and mappings $G_s^{(\varepsilon)}$. On the other hand, application of the group generated by mappings h_s (or by mappings $G_s^{(\varepsilon)}$) to an arbitrary loop allows a creation of a family of loops with arbitrary *moving poles*. This justifies the parameterizations of Example 0.1 via Remark 0.2.

0.7. Locality condition and singularities. As we have seen it above, a simultaneous consideration of “basic” functionals on loop spaces and “basic” diffeomorphisms of loop spaces leads to OPEs and OPE-like singularities. We also saw that the calculus of classical vertex operators is equivalent to the calculus involving such basic functionals and basic transformations.

It turns out that there is another reason to consider OPE-like singularities; it is connected with another crucial feature of loop spaces: a special role played by “local functionals”; this phenomenon is completely lacking in the situation $\dim < \infty$. In addition to “locality” on $\mathcal{L}X = \{Z \rightarrow X\}$ (e.g., consideration of open subsets of $\mathcal{L}X$) one must consider *locality along a loop* (i.e., consideration of open subsets of Z). As an illustration, consider the language of calculus of variations: when one considers a functional $F(\varphi)$ on the space of loops $Z \xrightarrow{\varphi} X$, the partial derivatives of F are written as $\delta F / \delta \varphi(z)$; they depend on a *continuous* index z ; it is crucial to analyze the degree of smoothness of the dependence of these data on z .

A functional of the form $F(\varphi) = \int_Z L(z, \varphi(z), \varphi'(z), \dots, \varphi^{(n)}(z)) dz$ (here φ is a loop $\varphi(z)$) is called a *local functional* of φ (here L is an arbitrary fixed function). When we consider two local variation of the loop φ , one near $z_1 \in Z$, another near $z_2, z_2 \neq z_1$, the changes of F w.r.t. these two variations are additive; in other words, the mixed variational derivative $\Delta(z, z') = \frac{\delta^2 F}{\delta \varphi(z) \delta \varphi(z')}$ vanishes outside of the diagonal $\{z = z'\}$: for an appropriate N one has $(z - z')^N \Delta(z, z') = 0$; here $\Delta(z, z')$ is a generalized function on Z^2 .

The appearance of δ -function singularity of $\Delta(z, z')$ on the diagonal is very similar to appearances of OPEs in the theory of loop spaces. This singularity is not a pole, however, as we will see it in the next section, there is a standard technique to construct related objects with poles.

0.8. Skyscraper variations. Here we construct a bridge between the construction of the previous section, which lead to δ -function singularities, and the constructions of Section 0.5, which lead to singularities of the OPE type.

In the language of variational calculus one analyses the dependence of the functional on the ultimate local variation of a loop; it is $g_s^{(\varepsilon)} : \varphi \mapsto \varphi + \varepsilon \delta_s$; here δ_s is the skyscraper δ -function with support at $s \in Z$, and, at least in the case $X = \mathbb{A}^n$, ε is a tangent vector to X . Assume $X = \mathbb{A}^1$, then ε is a number. Consider $g_s^{(\varepsilon)}$ as a 1-parameter subgroup of $\text{Diff}(\mathcal{L}X)$; taking $\varepsilon \rightarrow 0$ gives a vector field δ_s on $\mathcal{L}X$. Define a linear operator $\mathbf{D}_s$ on $\text{Func}(\mathcal{L}X)$ as the action of δ_s .

We did not specify which flavor of the loop space $\mathcal{L}X$ we work with, but the most useful examples do not allow δ -like loops; thus one needs to consider $g_s^{(\varepsilon)}$ formally, as a generalized function. For any smooth measure $\Phi(z)$ on Z , the integral $\int_Z \Phi(z) \delta_z$ is a well-defined vector field on $\mathcal{L}X$, thus $\mathbf{D}_{[\Phi]} \stackrel{\text{def}}{=} \int_{S^1} \Phi(z) \mathbf{D}_z$ is a well-defined operator on $\text{Func}(\mathcal{L}X)$. Using this description, it is easy to verify that $[\mathbf{D}_s, \mathbf{E}_t] = \delta(s - t) \mathbf{E}_t$ (here both sides are considered as generalized functions of s).

Similarly to Section 0.5, the consideration of the products $\mathbf{D}_s \mathbf{E}_t$ and $\mathbf{E}_t \mathbf{D}_s$ separately is much more delicate.

To bridge this discussion with one of Section 0.5, it is convenient to consider the kernels $\Phi(z)$ of a special form. Assume that $Z = S^1$; embed S^1 in $\mathbb{C}$ as $\{|z| = 1\}$; let $\Phi_s = \frac{dz}{z-s}$; then $\mathbf{d}_s = \mathbf{D}_{[\Phi_s]}$; here $\mathbf{d}_s$ was introduced in Section 0.5. On the other hand, the importance of δ -function variations is related to them spanning the vector space of all functions; however, the functions Φ_s also span this vector space. This is easiest to see via the relation

$$\delta(x - s) = \frac{1}{2\pi i} \bar{\partial}_s \Phi_s;$$

in other words, at least formally, the family $\mathbf{D}_s$ can be reconstructed as

$$(0.2) \quad \mathbf{D}_s = \frac{1}{2\pi i} \bar{\partial}_s \mathbf{d}_s.$$

This formula allows one to replace considerations of $\mathbf{D}_s$ by considerations of $\mathbf{d}_s$. Since $\mathbf{d}_s$ is a bona fide vector field on $\mathcal{L}X$ (at least for $|s| \neq 1$), it is much more convenient to work with. In what follows we do not work with the family of vector fields $\mathbf{D}_s$; however, make one more remark about this family.

Our treatments of the classical vertex operators in Section 0.5 skipped one family out of six: $\sum_{n \in \mathbb{Z}} s^{-n-1} \partial_n$ was not related to either one of operators $\mathbf{E}_t$, $\mathbf{d}_s$ and $\mathbf{h}_s$ on different flavors of loop spaces. However, the usual “physical” interpretation of $\sum_{n \in \mathbb{Z}} s^{-n-1} z^n$ is that it describes the δ -function at the point s . This leads to identification of $\sum_{n \in \mathbb{Z}} s^{-n-1} \partial_n$ with the operator $\mathbf{D}_s$. We do not know how to relate this heuristic argument with Equation (0.2).

0.9. Birational geometry and partial operators. As we saw it in Section 0.4, the set $\mathcal{L}^r X$ of rational loops is not a variety: it does not makes sense to consider

regular functions on $\mathcal{L}^r X$. This is a crucial phenomenon; one of the principal aims of this paper is to show that the set $\mathcal{L}^r X$ has a well-defined “birational geometry”: it makes sense to consider *meromorphic* functionals of a loop. During the introduction, we call such sets “*birational manifolds*”.

The principal feature⁵ of the birational geometry of infinite-dimensional manifolds is that there is no notion of “finite codimension”; show how such a notion would lead to a contradiction in the case of $\mathcal{L}\mathbb{P}^1$. Fix $z \in Z$; the 0-divisor D_0 of Eval_z is given by the equation $\varphi(z) = 0$ on φ ; if there is a theory of codimension, this set *must* have codimension 1. Since the inversion ι of $\mathcal{L}\mathbb{P}^1$ (induced by $x \mapsto x^{-1}$, $x \in \mathbb{P}^1$) must be a diffeomorphism of $\mathcal{L}\mathbb{P}^1$, the ∞ -divisor D_∞ of Eval_z must be also of codimension 1. However, the “diffeomorphism” h_z of Section 0.5 identifies $\mathcal{L}\mathbb{P}^1 \setminus D_0$ with D_∞ . Thus either we do not consider the invertible mapping h_z as a diffeomorphism, or we have no viable notion of codimension. Since the former kills the geometric description of vertex operators of Section 0.5, we are stuck with the later.

In particular, a diffeomorphism of an infinite-dimensional birational manifold M can send it “into” a “divisor of poles” of a meromorphic function (or even into the “indeterminacy subset” of such a function). Thus the action of a diffeomorphism on the meromorphic functions on M is not always defined. This is one of the key ingredients of our approach: the action of group G on M is encoded not in a family of linear operators on $\text{Func}(M)$, but in a family of *partial linear operators* on $\text{Func}(M)$ (here Func stands for the vector space of meromorphic functions).

One of the aims of this paper is to demonstrate that rules of composition of partial linear operators fully reflect the properties of products in vertex operator algebras. Moreover, in appropriate vector spaces any two smooth families of partial linear operators satisfy OPE relations.

Note that products of two smooth families of partial operators in a finite-dimensional vector space automatically possess properties *similar* to OPE: for the simplest example consider a partial operator P and a family Q_t ; then the dimension of the domain $\text{Dom}(R_t)$ of $R_t \stackrel{\text{def}}{=} P \circ Q_t$ is the same for generic t , but has jumps where $\text{Image}(Q_t)$ take special position w.r.t. $\text{Dom}(P)$. In other words, the composition $P \circ Q_t$ has singularities for special values of t ; this property is similar to OPE.

In Section 1.6 we formulate conditions on special families of partial operators in an infinite-dimensional vector space (which we call *codominant conformal fields*); one of the principal results of this paper is that compositions of such families satisfy *exactly* the OPEs. This gives the interpretation of the OPEs purely in the terms of the properties of the *vector space* in which the families of the operators act.

0.10. Flavors of the theories. If one needs more convincing examples of conformal fields, one needs more precise definitions of the building blocks: $\mathcal{L}X$, $\text{Func}(\mathcal{L}X)$, and of smooth families of operators in $\text{Func}(\mathcal{L}X)$. The examples of families $\mathbf{E}_t$, $\mathbf{d}_s$ and

⁵It is this feature which is the principal geometric reason for appearance of OPE for geometric operators.

$\mathbf{h}_s$ in Section 0.5 show that it is very convenient to be able to analytically extend the loops from $S^1 \subset \mathbb{C}$.

This is a basis of the *analytic approach*. It assumes that X and Z are complex-analytic; for an example, let $Z = A_\varepsilon = \{1 < |z| < 1 + \varepsilon\}$. This gives a loop space $\mathcal{L}_\varepsilon X$ consisting of loops $\varphi(z)$ in X , $z \in A_\varepsilon$. Then the family $\mathbf{E}_t$ is well defined and analytic⁶ for $t \in A_\varepsilon$; the operators $\mathbf{d}_s$ and $\mathbf{h}_s$ are well-defined and analytic for $t \notin A_\varepsilon$. As we saw it already in Section 0.6, in this approach the composition $\mathbf{h}_s \mathbf{E}_t$ is defined on too small a subset of $\mathbb{C} \times \mathbb{C}$ to define fusions via Cauchy integrals. Even if one allows ε to vary, one can define $\mathbf{h}_s \mathbf{E}_t$ only for $|s| > |t|$. In physics-speak, this leads to a *time-ordered* product—time being $|z|$.

This difficulty, having the operator product well-defined only on small subset of the desired set, is immanent to all approaches to the loop spaces based on regular functions on $\mathcal{L}X$. Another example of such approach is the *formal approach*; in this approach $\mathcal{L}^f X$ consists of *formal loops* in X , i.e., the mappings $\varphi \in X((z))$ of a “punctured formal disk” into X ; this set is well-defined if $X = \mathbb{A}^m$ (or $X \subset \mathbb{A}^m$). A loop can be written as a formal Laurent series $\sum_{n \geq -N} x_n z^n$, $x_n \in \mathbb{A}^m$. The loops with $N = 0$ are *holomorphic formal loops*, which form a subset $\mathcal{L}_+^f X \subset \mathcal{L}^f X$.⁷

Since the punctured formal disk plays the role of the limit of annuli $\{r < |z| < r + \varepsilon\}$ when $r, \varepsilon \rightarrow 0$, the considerations for the analytic case show that the family of operators $\mathbf{h}_s$ is defined on a large set $s \neq 0$. However, the family $\mathbf{E}_t$ is defined “nowhere”; it depends on the parameter t as a formal Laurent series in t . The coefficients of this formal series are well-defined operators in $\text{Func}(\mathcal{L}^f X)$. Thus to discuss the “poles of the composition” $\mathbf{h}_s \mathbf{E}_t$ one needs to invent an appropriate purely-algebraic language.

The great advantage of the formal approach is that all the objects are direct and/or inverse limits of finite-dimensional manifolds; thus the task of definition of $\text{Func}(\mathcal{L}^f X)$ becomes purely algebraic. E.g., the vector space $\text{Func}(\mathcal{L}^f X)$ consists of *polynomials* in variables x_n from Section 0.5 (see [3] for details). Thus the formal approach avoids the principal difficulty: that the infinite-dimensional differential geometry is currently very far from being a satisfactory theory (recall that—with the exception of Banach manifolds—infinite dimensional analysis is still in its cradle).

Finally, in the *rational approach* of this paper, Z is a projective manifold, X is a quasiprojective manifold, and $\mathcal{L}^r X$ consists of rational mappings $Z \rightarrow X$; in other words, of mappings $Z_0 \rightarrow X$ with Z_0 being dense open in Z ; two mappings are the same if their restrictions to some open dense subset Z_{00} coincide. As Example 0.1 shows, with reasonable definitions, $\mathcal{L}^r X$ has no non-constant regular function. However, $\mathcal{L}^r X$ has many “rational functions” (e.g., Eval_t , $t \in Z$); it is the interplay of the

⁶After an appropriate definition of $\text{Func}(\mathcal{L}_\varepsilon X)$.

⁷The set $\mathcal{L}_+^f X$ is well-defined for an arbitrary manifold X ; while it is hard to define $\mathcal{L}^f X$ in such a case, the formal neighborhood $\mathcal{L}^{ff} X$ of $\mathcal{L}_+^f X$ in $\mathcal{L}_+^f X$ is well-defined. One can describe this neighborhood by the condition that x_n with $n < 0$ are confined to an infinitesimal neighborhood of 0. See [9] for details.

functions not being defined everywhere, and of diffeomorphisms of $\mathcal{L}^r X$ (e.g., $g_t^{(\varepsilon)}$) mixing “divisors” and “complements to divisors” which leads to appearance of OPE in the algebraic geometry of $\mathcal{L}^r X$.

The advantages of the rational approach are the following. First, (same as in the analytic approach), the crucial families of operators (such as $\mathbf{h}_s, \mathbf{E}_t$) are well-defined families depending on points $s \in S, t \in T$ (see Section 2). Second, the compositions (such as $\mathbf{h}_s \mathbf{E}_t$) are well-defined on large enough subsets of $S \times T$ so that the fusions can be defined via Cauchy integrals (see Section 4). Third, the crucial definitions (of the set $\mathcal{L}^r X$, of the vector space $V = \text{Func}(\mathcal{L}^r X)$, and of which families of operators in V are smooth) turn out to be purely algebraic (see Sections 1.1 and 1.6).

In particular, the analysis on the set $\mathcal{L}^r X$ considered as an “infinite dimensional smooth manifold” does not require introduction of topology on this set! Out of the data encoded into the definition of a smooth manifold, we use only the notion of “smooth families”; in other words, which mappings $M \rightarrow \mathcal{L}^r X$ are smooth (for finite-dimensional manifolds M). In other words, essentially we consider $\mathcal{L}^r X$ as a functor $\text{Manifolds} \xrightarrow{\text{Families}} \text{Sets}$; the set $\mathcal{L}^r X$ is reconstructed as the particular value $\text{Families}(M_0)$ of the functor when M_0 is a point.

Note the surprising analogy of this approach with the so-called “convenient settings” [11] in analysis. In the latter approach one *starts* with a topology of a would-be-manifold L , and defines which mappings $M \rightarrow L$ (with $\dim M < \infty$) are smooth via an atlas on L ; however, after this initial step, one more or less forgets about the initial topology, constructing all data basing on this “structure of the functor” on Y .

0.11. On $\text{Func}(\mathcal{L}^r X)$. In this section we give a heuristic sketch of results of [12] which reduces the birational geometry of $\mathcal{L}^r X$ to purely algebraic terms. We motivate the definition of $\text{Func}(\mathcal{L}^r X)$ from Section 1.1 via more detailed consideration of the family $\text{Eval}_s, s \in Z$, of functionals on $\mathcal{L}^r X$.

First of all, in which sense is the functional $\text{Eval}_s: \varphi \mapsto \varphi(s)$ a meromorphic functional of φ ? Answer: the restriction of this functional to “natural” finite-dimensional families in $\mathcal{L}^r X$ is meromorphic. List some of such families.

Fix an ample line bundle $\mathfrak{L}$ on Z . Recall that one can define *the degree* of a loop $\varphi \in \mathcal{L}^r \mathbb{A}^1, Z \xrightarrow{\varphi} \mathbb{A}^1$, as the minimum of k such that $\varphi = S/S'$ with $S, S' \in \Gamma(Z, \mathfrak{L}^k)$. One can proceed similarly for $\mathcal{L}^r X$ for other quasiprojective manifolds X . Now $\mathcal{L}^r X$ is a union of subsets R_k consisting of loops of degree k or less. For large enough k , Eval_s is defined on R_k (for small k it may happen that R_k is in the subset of indeterminacy of Eval_s , if s is in the base set of the linear system for $\mathfrak{L}$). Moreover, one can parameterize R_k via the mapping $\tilde{R}_k \stackrel{\text{def}}{=} \mathbb{P}(\Gamma(Z, \mathfrak{L}^k)^2) \rightarrow R_k: (S, S') \mapsto S/S'$. This is a bijection on a dense open subset of $\tilde{R}_k$. Clearly, Eval_s corresponds to a meromorphic function $\varepsilon_{s,k}$ on $\tilde{R}_k$; one could call Eval_s a “pro-meromorphic” function. This leads to the following:

Definition 0.7. A meromorphic functional F on $\mathcal{L}^r X$ is a collection of meromorphic functions F_k on $\tilde{R}_k$ “compatible” with natural inclusions $R_k \rightarrow R_{k+1}$.

The problem of this definition is that it combines two objects: R_k and $\tilde{R}_k$; it turns out that the “correct” notion of “compatibility” is delicate.⁸ Indeed, F is not everywhere defined on R_{k+1} .

Instead of the mappings $R_k \rightarrow R_{k+l}$, consider their liftings $\tilde{R}_k \rightarrow \tilde{R}_{k+l}$. The lifting is not uniquely defined; for any $T \in \Gamma(Z, \mathfrak{L}^l)$, the mapping $(S, S') \xrightarrow{\alpha_T} (TS, TS')$ is a lifting. Thus the set of liftings is parameterized by $\mathbb{P}\Gamma(Z, \mathfrak{L}^l)$. A particular mapping α_T may⁹ send $\tilde{R}_k$ into the indeterminacy set of F_{k+l} . Thus the compatibility condition should be formulated as this: for a generic $T \in \mathbb{P}\Gamma(Z, \mathfrak{L}^l)$ the set $\alpha_T \tilde{R}_k \subset \tilde{R}_{k+l}$ is not contained in the indeterminacy set of the meromorphic function F_{k+l} on $\tilde{R}_{k+l}$; moreover,¹⁰ $\alpha_T^* F_{k+l} = F_k$. It turns out that this condition is not enough for functoriality: one needs also require that $\alpha_T^{-1} \mathcal{I}_{F_{k+l}} = \mathcal{I}_{F_k}$; here $\mathcal{I}_G$ denotes the indeterminacy set of a meromorphic function G .

It is obvious that the defined above functions $\varepsilon_{s,k}$ (induced by Eval_s) satisfy these compatibility conditions. Similarly to $\text{Eval}_s: \varphi \mapsto \varphi(s)$, one can send φ to the K -th partial derivative of φ at s ; here K is a multiindex. This gives another meromorphic function $\text{Eval}_{s,K}$ on $\mathcal{L}^r X$; as one can easily check, functions $\text{Eval}_{t,K}$ are algebraically independent. Since $\text{MerFunc}(\mathcal{L}^r X)$ is a field, one can consider its subfield freely generated by $\text{Eval}_{t,K}$. Call the latter field $\text{NaiveFunc}(\mathcal{L}^r X)$.

One of the principal conclusions of [12] is that in the case $Z = \mathbb{P}^1$ one has $\text{NaiveFunc}(\mathcal{L}^r X) = \text{MerFunc}(\mathcal{L}^r X)$. One can conjecture that a similar identity holds for an arbitrary Z . No matter what is the fortune of this conjecture, in this paper we *define* a “meromorphic function” on $\mathcal{L}^r X$ as an element of $\text{NaiveFunc}(\mathcal{L}^r X)$. The motivation for such a definition is that it quickly leads to a rich theory with all the manifestations of OPEs present from the start.

0.12. Partial operators in $\text{NaiveFunc}(\mathcal{L}^r X)$. Now, when we finally defined the vector space $\text{NaiveFunc}(\mathcal{L}^r X)$, we can analyze the operators $\mathbf{E}_t$, $\mathbf{d}_s$ and $\mathbf{h}_s$ in this space. First of all, the operator $\mathbf{E}_t$ are bona fide linear operators in $\text{NaiveFunc}(\mathcal{L}^r X)$. The operator $\mathbf{E}_t$ multiplies a rational expression of symbols $\text{Eval}_{z,K}$ (as any element of $\text{NaiveFunc}(\mathcal{L}^r X)$ is) by the symbol Eval_t . The result is a rational expression of the same form.

On the other hand, the operators $\mathbf{d}_s$ and $\mathbf{h}_s$ are of a different nature. E.g., obviously $\text{Eval}_t \circ \mathbf{h}_s = \frac{1}{t-s} \text{Eval}_t$. This can be taken as a definition of the action of $\mathbf{h}_s$ on

⁸It is this detail which leads to the fact that diffeomorphism only act as partial linear operators on meromorphic functions (thus to OPEs); hence the complication we discuss in this section is actually a central point of the geometry of $\mathcal{L}^r X$.

⁹As already happens with the functional Eval_s .

¹⁰Note that this condition applied to a locally defined *regular* functions on $\tilde{R}_k$, $k \geq 0$, can be satisfied by constant functions only. See Example 0.1 for details.

NaiveFunc ($\mathcal{L}^r X$) : $\mathbf{h}_s(\text{Eval}_t) = \frac{1}{t-s} \text{Eval}_t$, with obvious modifications (given by Leibniz identity) for the action on $\text{Eval}_{t,K}$ (e.g., $\mathbf{h}_s(\text{Eval}_{t,1}) = \frac{1}{t-s} \text{Eval}_{t,1} - \frac{1}{(t-s)^2} \text{Eval}_{t,0}$), and extending to rational expressions as an automorphism of an algebra. (Similarly for the action of $\mathbf{d}_s$, which acts as a derivation in the algebra.)

However, the formula for $\mathbf{h}_s$ does not make sense when $t = s$. Instead of trying to redefine this formula in this case, accept this as a fact: the operators $\mathbf{h}_s$ and $\mathbf{d}_s$ are only *partial operators* (i.e., operators defined on a vector subspace only). We already discussed the geometric reasons for this in Section 0.9; we are going to consider this in details in a foregoing paper [12].

A closely related statement from geometry of $\mathcal{L}^r X$ is that $\mathcal{L}^r X$ has no *regular functions*, see Example 0.1. Provide a different illustration of this phenomenon:

Example 0.8. Consider a constant loop $\varphi(z) \equiv 0 \in R_0 \subset \mathcal{L}^r X$. Then $\text{Eval}_0(\varphi) = 0$. Include $\varphi = \varphi_0$ into a one-parameter family $\varphi_s(z) = \frac{s}{s-z}$. Then $\text{Eval}_0(\varphi_s) \equiv 1$; thus the value of Eval_0 on φ_0 is not compatible with the value of Eval_0 on the family $\{\varphi_s\}$. To restore the compatibility, we need two modifications: first, we need to include φ into a larger family (e.g., 3-parameter parameterizing R_1 ; for simplicity, consider a smaller family $\varphi_{s,r} = \frac{s}{r-z}$); second, out of many representatives $\varphi_{0,r}$ for φ in this larger family we need to choose those with $r \neq 0$. After this, the compatibility is restored: indeed, if $r_0 \neq 0$, then $\lim_{(s,r) \rightarrow (0,r_0)} s/r$ exists and is 0.

Remark 0.9. Compare the need to consider an open subset of values r with the compatibility relation of Section 0.11.

This example shows a geometric reason for a diffeomorphism not necessarily acting on a meromorphic function if $\dim = \infty$: the meromorphic function is defined via its “restrictions” to a special exhausting collection of finite-dimensional families (such as R_d) inside our infinite-dimensional manifold; these restrictions are compatible. However, one should not expect that a restriction to *any* finite-dimensional family is compatible with this exhausting collection of families; some families are “too skew” to be compatible.

Now a diffeomorphism can map any “good” family to a skew¹¹ one; then the action of this diffeomorphism on the meromorphic function is not defined. E.g., h_s maps any “good” family to a family on which Eval_s has a pole. *This* is the reason why $\mathbf{h}_s(\text{Eval}_t)$ has a pole at $s = t$.

0.13. Basic and generalized functions. The principal reason why OPEs appear in the formal theory “in a formal way” only is the fact (see Section 0.10) that the family $\mathbf{E}_t$ of operators is defined only as a formal series in t . Comparing this with Theorem 0.3, one can see that the reason for this deficiency is that $\text{Func}(\mathcal{L}^f X)$ is “too small”; multiplication by Eval_t “moves” a functional “outside” of the set

¹¹W.r.t. the given meromorphic function.

$\text{Func}(\mathcal{L}^f X)$. Our approach increases the vector space of function on the loop space so that the space remains invariant under multiplication by Eval_t .

To show how consideration of $\mathcal{L}^r X$ is related to such an increase, recall the usual duality between basic and generalized functions. While for finite-dimensional vector spaces increasing V increases the dual space V^* as well, for topological vector spaces increasing V (in such a way that V is dense in the result) *decreases* V^* . To make the space of generalized functions as large as possible, one considers the space of base functions which is as small as possible. The similar effect holds in non-linear situation, when one considers $\text{Func}(X)$ for an infinite-dimensional manifold X instead of the set of linear functionals on an infinite-dimensional vector space V . If X is dense in $X' \supset X$, then $\text{Func}(X')$ is a subset of $\text{Func}(X)$.

Thus the reason why the operators $\mathbf{E}_t$ act in $\text{Func}(\mathcal{L}^r X)$ as honest operators is that $\text{Func}(\mathcal{L}^r X)$ is very large due to $\mathcal{L}^r X$ being very small (comparing to $\mathcal{L}^f X$). In fact $\mathcal{L}^r X$ is the orbit in $\mathcal{L}^f X$ of the action of diffeomorphisms $G_s^{(\varepsilon)}$ and/or H_s (from Section 0.5). Thus one should expect that the vector space $\text{Func}(\mathcal{L}^r X)$ is the largest possible space of functions in which the operators $\mathbf{E}_t$ and $\mathbf{h}_s, \mathbf{d}_s$ act as honest (partial) operators.

0.14. The plan. In this paper we *start* at the situation described in the previous sections. The principal objects we work with are families of partial linear operators in the vector space $V = \text{NaiveFunc}(\mathcal{L}^r X)$. The families $\mathbf{E}_t, \mathbf{d}_s$ and $\mathbf{h}_s$ become just particular examples of such families.

For the coarse theory of this paper the target manifold X of the loop does not matter much, we consider the case of the multiplicative group $X = \mathbb{C}^*$ only. In this case the set $\mathcal{L}_Z X$ of meromorphic loops becomes the set $\mathcal{M}(Z)$ of meromorphic functions on Z . To shorten the notation, denote $\text{NaiveFunc}(\mathcal{L}_Z \mathbb{C}^*)$ on $\mathcal{M}(Z)$ by $\mathfrak{MM}(Z)$.

In Sections 1.1 and 1.2 we equip the vector space $V = \mathfrak{MM}(Z)$ with two structures: a lattice of vector subspaces of V , and notion of a *smooth family* of vectors in V . To define the lattice, define the support of $\text{Eval}_{t,K} \in V$ as $\{t\} \subset Z$; support of a rational expression is the union of the supports of the symbols $\text{Eval}_{t,K}$ which appear in this expression. Now to each Zariski open subset $U \subset Z$ one can associate the vector subspace $V(U)$ consisting of $v \in V$ such that $\text{Supp}(v) \subset U$.

In Section 1.6 we introduce the notion of a *codominant conformal field*; this is a family of partial linear operators in V which behaves nicely with respect to two structures (lattice and smoothness) in V . Although $\mathbf{E}_t, \mathbf{d}_s$ and $\mathbf{h}_s$ (and other family coming from algebraic geometry of $\mathcal{L}^r X$) are examples of such families, in what follows we completely ignore the algebraic geometry of $\mathcal{L}^r X$: it turns out that the two specified above structures on V is *all* that is needed to treat conformal fields. In particular, *any* two codominant conformal fields satisfy the OPEs.

In the rest of Section 1 we define the rest of the principal objects of this paper: the field $\mathfrak{MM}(Z)$, conformal operators, conformal families, residues, fusion and

localization. In Section 2 we study a particular case $Z = \mathbb{P}^1$ and show that on localization at $0 \in \mathbb{P}^1$ one obtains the standard vertex operators (this enhances the discussion of Section 0.5). In Section 3 we show that appropriate projective bundles over Hilbert schemes of points in Z provide good finite-dimensional models of the infinite-dimensional vector space $\mathfrak{MM}(Z)$, and deduce that after applying a natural restriction, conformal fields automatically satisfy conformal associativity relations and vertex commutation relations. In Section 4 we show how chiral algebras are related to our definition of conformal fields, calculate vertex commutation relations in the case $\dim Z = 1$, and show how one of examples of Section 2 gives rise to algebras $\mathfrak{gl}_\infty$ and $\mathfrak{g}(\infty)$ when localized to Laurent series. The results of this section show that the standard “physical degree of strictness” arguments deducing Lie-algebraic identities comparing OPEs and cohomologies of a plane with 3 removed lines can be made honestly for the conformal fields of the type considered in this paper.

In Section 5 we provide an independent description of boson-fermion correspondence (this correspondence differs a little from the standard one), and show why it provides heuristics for considering the space $\mathfrak{MM}(\mathbb{P}^1)$ as a Fock space (in $1 + 1$ -dimensional theory).

I am deeply grateful to R. Borcherds, I. M. Gelfand, A. Givental, V. Goryunov, I. Grojnowsky, R. Hartshorn, Y.-Z. Huang, M. Kapranov, M. Kontsevich, A. Kresch, F. Malikov, N. Reshetikhin, M. Spivakovsky, V. Serganova and A. Vaintrob for fruitful discussions during the work over these concepts. The revision of the introduction was made possible by hospitality of IHES, MPIM-Bonn and MSRI. This paper was partially supported by Alfred P. Sloan research fellowship.

Revision history. The revisions of this paper are archived as [arXiv:math.AG/9805096](https://arxiv.org/abs/math.AG/9805096) on the math preprint server archive [arXiv.org/abs/math](https://arxiv.org/abs/math). In addition to cosmetic changes, Revision II of this paper contains a complete rework of the introduction (with the exception of Section 0.16). The numeration of statements outside the introduction did not change.

0.15. Alternative approaches. Two properties of the lattice $V(U)$ in $V = \mathfrak{MM}(Z)$ introduced above are most important for our approach: one is the *fatness*: for any element $v \in V$ and any $U \subset Z$ there is a family v_t , such that $v_0 = v$, and $v_t \in V(U)$ for $t \neq 0$. Compare this with Remark 1.11. Another is a related property of *stability*: given the existence of moduli spaces of subvarieties of Z , say that U_t is a *smooth one-parameter family of Zariski open subsets* of Z if the complement family $\mathcal{C}U_t$ is such. This leads to a one-parameter family of vector subspaces $V(U_t) \subset V$. Due to the description of the tangent space to the Grassmannian, any smooth variation W_t of a vector subspace $W \subset V$ leads to a *variation operator* $W \rightarrow V/W$. Now it is easy to see that the variation operator for the family $V(U_t)$ is actually 0. This is the stability property: $V(U_t)$ is a non-trivial family with vanishing infinitesimal variation. Conjecturally, the fatness and stability properties are what substitutes the

factorization property of the Drinfeld–Beilinson approach to the loop spaces ([6] and [9]).

Since in the case $Z = \mathbb{P}^1$ the families of operators $\mathbf{E}_t$, $\mathbf{d}_s$ and $\mathbf{h}_s$ coincide with classical vertex operators, in general case they are multidimensional analogues of vertex operators. Recall that the general theory of [12] shows that at least in the case $Z = \mathbb{P}^1$ one can describe $\mathfrak{M}\mathcal{M}(Z)$ as the field of meromorphic functions on $\mathcal{M}(Z)$ (see Section 0.11). Since for general Z this is not clear yet, it may be too early to identify multi-dimensional theory of vertex operators with birational geometry of sets $\mathcal{M}(Z)$; however, this is a very natural conjecture.

A different approach to multidimensional vertex operators is proposed in [1]. Note that Hilbert schemes in relation to vertex operators appear also in [10] and [13], but the setting of these papers do not look directly related to what we discuss here.

0.16. Arithmetic analogue. Consider here the result of formally substituting $Z = \text{Spec } \mathbb{Z}$ into the geometric considerations of the previous sections. One should consider $\mathbb{Q}$ as the set $\mathcal{M}(\text{Spec } \mathbb{Z})$ of meromorphic functions on Z . Results of Section 3.2 show that it is natural to suppose that meromorphic functions on $\mathcal{M}(\text{Spec } \mathbb{Z})$ may be defined via restrictions from $\mathcal{M}(\text{Spec } \mathbb{Z})$ to $\Gamma(S, \mathcal{O})$ for finite subschemes $S \subset \text{Spec } \mathbb{Z}$. Thus a meromorphic function on $\mathcal{M}(\text{Spec } \mathbb{Z})$ may be described by an ideal $N\mathbb{Z} \subset \mathbb{Z}$ of S and a function on $\mathbb{Z}/N\mathbb{Z}$. In other words, the obvious substitution for $\mathfrak{M}\mathbb{Q}$ is the space of locally constant functions on integer adeles $\mathbf{A}_{\mathbb{Z}}$.

There is a partially defined action of $\mathbb{Q}^\times$ on $\mathfrak{M}\mathbb{Q}$ given by the following rule: an element F of $\mathfrak{M}\mathbb{Q}$ is a function on $\mathbb{Z}/N\mathbb{Z}$ for some $N > 0$. Element $p/q \neq 0$ acts on F iff N can be taken prime to q , then it acts as multiplication by the corresponding invertible residue modulo N . This defines a partial multiplication in $\mathcal{C} = \mathbb{Q}^\times \ltimes \mathfrak{M}\mathbb{Q}$, and a partial action of $\mathcal{C}$ on $\mathfrak{M}\mathbb{Q}$. This partial multiplication is the analogue of conformal field theory, and the partial action gives analogues of vertex operators.

The previous interpretation has an advantage of $\mathfrak{M}\mathbb{Q}$ being an algebra. Sacrificing this property, one may consider locally constant partially defined functions on adeles $\mathbf{A}_{\mathbb{Q}}$ with compact *domain of definition* as a substitution for $\mathfrak{M}\mathbb{Q}$, i.e., functions with values in the union of numbers and a new symbol v (for “undefined”). In such a case one has a well-defined action of the group $\mathbb{Q}^\times$ on $\mathfrak{M}\mathbb{Q}$. However, one still has no *fair* algebra structure on $\mathcal{C} = \mathbb{Q}^\times \ltimes \mathfrak{M}\mathbb{Q}$, only a partial structure.

1. FIELD OF MEROMORPHIC FUNCTIONS ON $\mathcal{M}(Z)$

In this section we define principal objects (meromorphic functions on $\mathcal{M}(Z)$, conformal operators, conformal fields, residues, fusions and localizations) of this treatise.

1.1. Meromorphic functions. Consider an irreducible quasiprojective variety Z over a field $\mathbb{K}$ with a line bundle $\mathcal{L}$. Any two global sections n, d of $\mathcal{L}$ which are not simultaneously identically zero give a meromorphic function n/d on Z . This provides a family of meromorphic functions on Z , this family is parameterized by $X_{Z, \mathcal{L}} \stackrel{\text{def}}{=}$

$\mathbb{P}(\Gamma(Z, \mathcal{L}) \times \Gamma(Z, \mathcal{L}))$. (Since most of the time we consider a fixed quasiprojective variety Z , usually we reduce this notation to $X_{\mathcal{L}}$.)

We will need to consider $X_{\mathcal{L}}$ as an algebraic variety, thus $\Gamma(Z, \mathcal{L})$ should be a finite-dimensional vector space. This is automatically true if Z is projective. To *force* this to be true for an arbitrary quasiprojective Z , *define* a line bundle on such a Z to consist of a projective completion $\bar{Z}$ of Z , and a usual line bundle $\bar{\mathcal{L}}$ on $\bar{Z}$. Define $\Gamma(Z, \mathcal{L})$ as $\Gamma(\bar{Z}, \bar{\mathcal{L}})$.

Fix a point $z \in Z$. Suppose that z is not a base point of $\mathcal{L}$. Define the *evaluation function* E_z on $X_{\mathcal{L}}$ by

$$n/d \mapsto n(z)/d(z) \in \mathbb{P}^1 = \mathbb{K} \cup \{\infty\}.$$

It is a meromorphic function on $X_{\mathcal{L}}$, it is regular on an open subset $U_{\mathcal{L}, z} \subset X_{\mathcal{L}}$ given by $n(z) \neq 0$ or $d(z) \neq 0$.

Similarly, if z is smooth, and $v_1, \dots, v_n$ are meromorphic vector fields on Z which have no pole at z , then one can also define a *partial derivative* $E_z^{(v_1 \dots v_n)}$ of the evaluation function, given by $\xi \mapsto E_z(v_1 \dots v_n \xi)$. When we need to denote an arbitrary derivative, we will use notation $E_z^\bullet$. For a fixed z meromorphic functions $E_z^\bullet$ span a vector space with the same size as the symmetric power of the tangent space $T_z Z$. We will assume that $\bullet$ is an index from a basis of this symmetric power.

One can see that the vector space spanned by $E_z^\bullet$ for a fixed smooth z is the topological dual space $\widehat{\mathcal{O}}_z^\vee$ to the space of formal functions $\widehat{\mathcal{O}}_z$ near z . This description works for a non-smooth z as well, define indices $\bullet$ for $E_z^\bullet$ appropriately in the case of non-smooth z as well.

On the other hand, for a fixed multiindex $(v_1, \dots, v_n)$ one can consider $E_z^{(v_1 \dots v_n)}$ as a meromorphic function on $X_{\mathcal{L}}$ which depends on z . It is well defined on the complement to the set of base points of $\mathcal{L}$. Taking different values of z and/or $(v_1, \dots, v_n)$, one obtains many meromorphic functions on $X_{\mathcal{L}}$. Consider the field generated by these meromorphic functions.¹²

Consider how these functions depend on the choice of the line bundle $\mathcal{L}$. For any global section $\beta \not\equiv 0$ of another line bundle $\mathcal{M}$ on Z one can construct a mapping $X_{\mathcal{L}} \xrightarrow{\iota_\beta} X_{\mathcal{L} \otimes \mathcal{M}}$ given by $(n : d) \mapsto (n\beta : d\beta)$. Given a meromorphic function F on $X_{\mathcal{L} \otimes \mathcal{M}}$ such that the set of the base points of this meromorphic functions does not contain $\text{Im } \iota_\beta$, one can construct a meromorphic function $\iota_\beta^* F \stackrel{\text{def}}{=} F \circ \iota_\beta$ on $X_{\mathcal{L}}$.

Consider an arbitrary algebraic expression $\tilde{F}$ constructed from several symbols $\tilde{E}_z^\bullet$, $z \in Z$, with application of operations of addition, multiplication and division. Given $\mathcal{L}$, substitute the symbols $\tilde{E}_z^\bullet$ by the functions $E_z^\bullet$ on $X_{\mathcal{L}}$, separately in the numerator and denominator of $\tilde{F}$. Unless both the numerator and denominators become 0, one obtains a function $F^{(X_{\mathcal{L}})}$ on $X_{\mathcal{L}}$. One can try to perform the same procedure for

¹²Note that this field is generated by δ -functions on Z and their derivatives. Most approaches to the theory of generalized functions involve consideration of $\mathcal{D}$ -modules on Z . This may be a reason why $\mathcal{D}$ -modules appear in definitions of chiral algebras of [6] and [14].

$\mathcal{L} \otimes \mathcal{M}$ taken instead of $\mathcal{L}$. If both $F^{(X_{\mathcal{L}})}$ and $F^{(X_{\mathcal{L} \otimes \mathcal{M}})}$ are defined, and $F^{(X_{\mathcal{L} \otimes \mathcal{M}})}$ is compatible with ι_{β} in the above sense, then $\iota_{\beta} F^{(X_{\mathcal{L} \otimes \mathcal{M}})} = F^{(X_{\mathcal{L}})}$. Investigate for which values of $\mathcal{L}$ and $\mathcal{M}$ these assumptions hold.

Consider what is involved in the calculation of F at a given element $(n : d) \in X_{\mathcal{L}}$. Suppose for simplicity that F includes no partial derivatives of E_z . After all the operations are performed, $F((n : d))$ is given by an expression of the form

$$(1.1) \quad \frac{G(n(z_1), n(z_2), \dots, n(z_k), d(z_1), d(z_2), \dots, d(z_k))}{G'(n(z_1), n(z_2), \dots, n(z_k), d(z_1), d(z_2), \dots, d(z_k))},$$

here z_i are different points of Z , G and G' are polyhomogeneous polynomials of homogeneity degrees $(\delta_1, \dots, \delta_{2k})$, $(\delta'_1, \dots, \delta'_{2k})$. (These degrees satisfy $\delta_l + \delta_{l+k} = \delta'_l + \delta'_{l+k}$, thus the value of the above ratio is not “an element of an abstract 1-dimensional vector space”, but of $\mathbb{K}$.)

For this expression to be correctly defined, one needs $G(n(z_1), \dots, d(z_1), \dots)$ and $G'(n(z_1), \dots, d(z_1), \dots)$ to be not simultaneously identically zero as functions of $(n : d) \in X_{\mathcal{L}}$. Taking $\mathcal{L}$ positive enough, one can ensure that $n(z_1), n(z_2), \dots, n(z_k)$ are algebraically independent when n runs over $\Gamma(Z, \mathcal{L})$, same for d . This ensures correctness of F .

If F contains partial derivatives, then in the above expression one also needs to consider partial derivatives of $n(z_i)$ w.r.t. some (possibly meromorphic) connection on $\mathcal{L}$. After this all the arguments still work.

Thus it is clear that for any such formal expression $\tilde{F}$ one can pick up a corresponding line bundle $\mathcal{L}$ on Z such that $F^{(X_{\mathcal{L}})}$ is correctly defined, and in fact this remains true for any line bundle of the form $\mathcal{L} \otimes \mathcal{M}$ with an ample $\mathcal{M}$. The above arguments show that the manifolds $X_{\mathcal{L}}$ are objects of an appropriate (ordered) category, and any formal expression $\tilde{F}$ defines compatible meromorphic functions on “tails” of this category, i.e., on objects which are “greater” than some fixed object. (These functions are not necessarily compatible w.r.t. *all* possible morphisms between two given objects, but only w.r.t. an open dense subset of the set of morphisms, one can look up all the nasty details in [12].)

The motivations above lead to the following

Definition 1.1. Let Z be a quasiprojective variety. Let $\mathcal{M}(Z)$ be a set of meromorphic functions on Z . Consider a field $\mathfrak{M}\mathcal{M}(Z)$ freely generated by symbols $E_z^\bullet$, $z \in Z$. Call elements of this field¹³ *meromorphic functions* on $\mathcal{M}(Z)$.

Given any fixed meromorphic function F on $\mathcal{M}(Z)$, and any given meromorphic function ξ on Z one cannot guarantee that $F(\xi)$ is a well-defined number. However, one has seen that if ξ changes in sufficiently big families, then $F(\xi)$ is a well-defined meromorphic function on the base of such a family.

¹³Note that in analytic theories, $E_z^\bullet$ “span” the vector space of generalized functions on Z . Thus $\mathfrak{M}\mathcal{M}(Z)$ is the field spanned by generalized functions on Z .

Remark 1.2. Note that studying the interplay between different families $X_{Z,\mathcal{L}}$ of meromorphic functions, i.e., studying the algebraic geometry of $\mathcal{M}(Z)$, leads to a *very* complicated theory (see [12]). Moreover, the final result of this more complicated theory coincides with the above definition, at least in the simplest—but most important—case $Z = \mathbb{P}^1$.

Thus we accept the above ad hoc point of view, with *postulating* what is a meromorphic function on $\mathcal{M}(Z)$, consider $\mathcal{M}(Z)$ as a set only, and postulate which families of elements of $\mathcal{M}(Z)$ are “smooth”.

The collection $\{\mathfrak{M}\mathcal{M}(U)\}_{U \subset Z}$ for open $U \subset Z$ behaves as sections of a sheaf with compact supports.¹⁴ Denote it by $\mathfrak{M}\mathcal{M}$.

Collection $\{\mathfrak{M}\mathcal{M}(U)\}_{U \subset Z}$ forms a lattice of subfields of the field $\mathfrak{M}\mathcal{M}(Z)$. This lattice of vector subspaces becomes the playing field for conformal fields, they will act as operators defined on small enough spaces of this lattice.

1.2. Families of meromorphic functions on $\mathcal{M}(Z)$. We *defined* meromorphic functions on $\mathcal{M}(Z)$ to be rational expressions of symbols $E_{z_i}^\bullet$. Similarly, *define* families of such functions by allowing both the coefficients of the rational expressions and the points z_i to change in families. Before giving formal definitions consider some examples of what we want to define.

For an arbitrary regular mapping $T \xrightarrow{\iota} Z$ consider $E_{\iota(t)}$, $t \in T$. Obviously, one wants this family to be allowed.

Second, in the case $T = Z = \mathbb{A}^1$ one can consider the family $F_t = (E_t - E_{-t})/t$ if $t \neq 0$, obviously it “should be” extensible to $t = 0$ with the value $F_0 = 2E_0^{(d/dz)}$. Since F_t is even in t , $G_t \stackrel{\text{def}}{=} F_{\sqrt{t}}$ should be also allowed. The following definition is good enough if Z is complete, as Section 3.2 will show:

Definition 1.3. Consider an algebraic variety T . Say that a family $T \xrightarrow{F_\bullet} \mathfrak{M}\mathcal{M}(Z)$ is *admissible* if there are mappings $z_i: T \rightarrow Z$, $i = 1, \dots, I$, meromorphic vector fields v_{ij} , $i = 1, \dots, I$, $j = 1, \dots, J_i$, on Z and a meromorphic function $\tilde{F}(t, f_1, \dots, f_I) \not\equiv 0$ on $T \times \mathbb{A}^I$ such that F_t can be written as $\tilde{F}\left(t, E_{z_1(t)}^{\mathcal{V}_1}, \dots, E_{z_I(t)}^{\mathcal{V}_I}\right)$, here $\mathcal{V}_i = v_{i1} \dots v_{iJ_i}$, and v_{ij} has no pole at $z_i(t)$ for any i, j, t .

Say that a family $T \xrightarrow{F_\bullet} \mathfrak{M}\mathcal{M}(Z)$ is *strictly meromorphic on T* if it is admissible when lifted to a finite covering of an open dense subset of T , and $F_t(\xi)$, $\xi \in X_{\mathcal{L}}$, is a meromorphic function on $T \times X_{\mathcal{L}}$ for any ample line bundle $\mathcal{L} \geq \mathcal{L}_0$ on Z , $\mathcal{L}_0$ being some fixed line bundle on Z .

For a complete Z say that a family $T \xrightarrow{F_\bullet} \mathfrak{M}\mathcal{M}(Z)$ is *strictly smooth* if

- (1) it is strictly meromorphic;

¹⁴Recall that this means that the canonical mappings go in *opposite* direction comparing to usual sheaves. (Thus dual vector spaces $\mathfrak{M}\mathcal{M}(U)^\vee$ form a sheaf.)

- (2) intersections of the divisor of poles of the meromorphic function $F_t(\xi)$ on $T \times X_{\mathcal{L}}$ with fibers $\{t\} \times X_{\mathcal{L}}$ have codimension 1¹⁵ for any $\mathcal{L} \geq \mathcal{L}_0$.

To generalize this definition to the case of quasiprojective Z , note that for any meromorphic vector field v on T and a meromorphic family $F_{\bullet}$ the derivative $vF_{\bullet}$ is a well-defined meromorphic family. This family is smooth near $t_0 \in T$ if $F_{\bullet}$ is smooth and v is regular near t_0 .

Definition 1.4. Consider a projective variety $\bar{Z}$ and an open subset $Z \subset \bar{Z}$. Say that a family $T \xrightarrow{F_{\bullet}} \mathfrak{MM}(Z)$ is *strictly smooth* if it is strictly smooth when considered as a family in $\mathfrak{MM}(\bar{Z})$, and for any fixed $t_0 \in T$ and any meromorphic vector fields $v_1, \dots, v_k$ on T which are regular near t_0 the smooth families $v_1 \dots v_k F_{\bullet}$ in $\mathfrak{MM}(\bar{Z})$ are in fact families in $\mathfrak{MM}(Z)$.

Remark 1.5. The additional constraints of the latter definition may be easily seen in the case $T = \bar{Z} = \mathbb{P}^1$, $Z = \mathbb{A}^1$. In this case the family $F_t = \frac{E_t}{1+tE_t}$ sends any $t \in \mathbb{P}^1$ to $\mathfrak{MM}(\mathbb{A}^1)$. However, the derivative E_{∞} at $t = \infty$ is not in $\mathfrak{MM}(\mathbb{A}^1)$.

As with any ad hoc definition, a lot of problems disappear, but some other problems jump to existence. Say, with the above definition it is hard to show that a subfamily of a strictly smooth family is strictly smooth. On the other hand, it is obvious that strictly meromorphic families form a commutative algebra with division, strictly smooth mappings form a commutative algebra.

To avoid the problem with subfamilies, use the following trick:

Definition 1.6. Say that a family $T \xrightarrow{F_{\bullet}} \mathfrak{MM}(Z)$, $F_{\bullet} \neq \infty$, parameterized by a quasiprojective variety T is *smooth* if there is a quasiprojective variety T' , a strictly smooth mapping $T' \xrightarrow{G_{\bullet}} \mathfrak{MM}(Z)$, and a regular mapping $T \rightarrow T'$ such that the composition is $F_{\bullet}$.

Say that a family $T \xrightarrow{F_{\bullet}} \mathfrak{MM}(Z)$ is *meromorphic* if there is a blow-up $T' \xrightarrow{\pi} T$ such for any point $t \in T'$ there is a neighborhood $U \ni t$ and a regular scalar function $f \neq 0$ on U such that $f \cdot (F \circ \pi)$ gives a smooth mapping $U \rightarrow \mathfrak{MM}(Z)$. Say that $F_{\bullet}$ is *defined* at $t \in T$ if it is smooth in a neighborhood of t .

Now by definition the restriction of a smooth mapping to a subvariety is smooth or $\equiv \infty$, and meromorphic families form a commutative algebra with division, strictly smooth mappings form a commutative algebra. Of course, the above trick is just for simplicity, and

Fact . Any meromorphic family is strictly meromorphic, any smooth family is strictly smooth.

We will not use this statement in what follows. It follows from the results of Section 3.2.

¹⁵In algebro-geometric parlance, this divisor is *flat* over T .

Example 1.7. Suppose that Z is a point O . In such a case $\mathcal{M}(Z) = \mathbb{P}^1$, and $\mathfrak{MM}(Z)$ has one generator E_O , thus is $\mathbb{K}(E_O)$. Let $T = \mathbb{A}^2$ with coordinates (t, t') . Then any element of $\mathbb{K}(E_O, t, t')$ defines a meromorphic family of meromorphic functions on $\mathcal{M}(Z)$. Write this element as P/Q , $P, Q \in \mathbb{K}[E_O, t, t']$. Suppose that P and Q are mutually prime, write $Q = \sum_{k=0}^K a_k(t, t') E_O^k$. The family P/Q is smooth iff polynomials $a_k(t, t')$, $k = 0, \dots, K$, have no common zeros.

In particular, both $F_{t,t'} = t/t'$ and $G_{t,t'} = 1/(t + t'E_O)$ are meromorphic, but not smooth. $F_{t,t'}$ becomes smooth either after a multiplication by t' , or after a blow up of $(0, 0) \in \mathbb{A}^2$. $G_{t,t'}$ becomes smooth after multiplication by any scalar polynomial $P(t, t')$ such that $P(0, 0) = 0$ followed by a blow-up of $(0, 0) \in \mathbb{A}^2$.

A fundamental property of smooth and meromorphic families is their relation with the lattice $\{\mathfrak{MM}(U)\}$, U running through open subsets of Z . If F_t , $t \in T$, is a meromorphic family defined at t_0 , and $F_{t_0} \in \mathfrak{MM}(U)$, then $F_t \in \mathfrak{MM}(U)$ for T in some opens neighborhood of t_0 . Moreover, say that U_s , $s \in S$, is a *family of open subsets* of Z if there is an open subset $\mathbf{U} \subset Z \times S$ such that $\mathbf{U} \cap (Z \times \{s\}) = U_s \times \{s\}$. Then for any meromorphic family F_t and any family of open subsets $U_s \subset Z$ the subset

$$\{(t, s) \in T \times S \mid F_t \in \mathfrak{MM}(U_s)\}$$

is open.

Call this property a *locality property*. Heuristically, it states that a function $F \in \mathfrak{MM}(Z)$ has 0-dimensional support in Z . Strictly speaking, it would be better to define $\mathfrak{MM}$ as a sheaf on the Hilbert scheme of subschemes of Z of dimension 0, but we want to avoid all the algebro-geometric complications until Section 3.

Remark 1.8. Here is another trivial but important observation: for any quasiprojective variety S with an open dense subset U any smooth family on U extends to a meromorphic family on S .

1.3. Action of automorphisms of $\mathcal{M}(Z)$. We are going to define some partially defined mappings of vector spaces $\mathfrak{MM}(Z)$.

Consider a fixed meromorphic function $\eta \neq 0$ on Z . Multiplication by η gives an invertible mapping $\mathbf{m}_\eta: \mathcal{M}(Z) \rightarrow \mathcal{M}(Z)$. Consider how this mapping acts on meromorphic functions on $\mathcal{M}(Z)$. Given such a meromorphic function $F(\xi) \in \mathfrak{MM}(Z)$, try to find a meromorphic function $\mathbf{M}_\eta F$ which satisfies $(\mathbf{M}_\eta F)(\xi) \stackrel{\text{def}}{=} F(\eta\xi)$. For $\mathbf{M}_\eta F$ to exist, the mapping $\mathbf{m}_\eta$ and families $X_\mathcal{L}$ should satisfy some compatibility conditions.

First, one can assume that $\eta = n_0/d_0$, n_0, d_0 being global sections of a line bundle $\mathcal{M}$ on Z . Multiplication by η sends $X_\mathcal{L}$ into $X_{\mathcal{L} \otimes \mathcal{M}}$. If $\mathcal{L}$ is positive enough, then F is correctly defined on $X_{\mathcal{L} \otimes \mathcal{M}}$. However, the image $\mathbf{m}_\eta X_\mathcal{L} \subset X_{\mathcal{L} \otimes \mathcal{M}}$ may lie inside the base set of the meromorphic function $F|_{X_{\mathcal{L} \otimes \mathcal{M}}}$, where it is not defined, thus it is possible that $F(\eta\xi)$ is not defined for $\xi \in X_\mathcal{L}$ for any $\mathcal{L}$.

Suppose that there is $\mathcal{L}_0$ such that for $\mathcal{L} \geq \mathcal{L}_0$ the above situation will not happen, i.e., $\mathbf{m}_\eta X_\mathcal{L} \subset X_{\mathcal{L} \otimes \mathcal{M}}$ intersects with the base set of $F|_{X_{\mathcal{L} \otimes \mathcal{M}}}$ along a subvariety of

codimension two¹⁶ in $\mathbf{m}_\eta X_\mathcal{L}$. (Recall that the base set is the intersection of the divisor of zeros and the divisor of infinity, and it has codimension two.) Then $\mathbf{m}_\eta^* F$ is a well-defined meromorphic function on $X_\mathcal{L}$.

It is clear that obtained in such a way functions on $X_\mathcal{L}$ (for all $\mathcal{L} \geq \mathcal{L}_0$) are compatible in the same sense as algebraic combinations of $E_z^\bullet$ are. Thus $\mathbf{m}_\eta^*$ gives a (partially defined) mapping from the set of meromorphic functions on $\mathcal{M}(Z)$ into itself. Denote this mapping by $\mathbf{M}_\eta$.

Moreover, if $\mathbf{M}_\eta F$ and $\mathbf{M}_{\eta'} \mathbf{M}_\eta F$ are both well-defined, then so is $\mathbf{M}_{\eta' \eta} F$, and $\mathbf{M}_{\eta'} \mathbf{M}_\eta F = \mathbf{M}_{\eta' \eta} F$. In particular, one can see that any mapping $\mathbf{M}_\eta$ is automatically invertible (on the domain of definition).

Example 1.9. Suppose that η has no pole or zero at $z \in Z$. Then $\mathbf{M}_\eta E_z$ is well-defined, and equals $C \cdot E_z$, $C = \eta(z)$. It is clear that if η has zero at z , then one should require $C = 0$, which contradicts invertibility of $\mathbf{M}_\eta$.

Yet stronger breach of contract comes in the case when z is in the base set of η , when C is not defined at all.

From the above description one can easily conclude that $\mathbf{M}_\eta$ is well defined on $\mathfrak{MM}(U)$ if η has no zero and no pole on $U \subset Z$, and that it sends smooth families in $\mathfrak{MM}(U)$ to smooth families. Note also that the operators $\mathbf{h}_s$ and $\mathbf{H}_s$ of Section 0.5 are of this form.

1.4. Multiplication in the field, smooth families, and conformal operators. Consider a different construction of mappings which send meromorphic functions on $\mathcal{M}(Z)$ into themselves. The study of these mappings will justify the definition of general conformal fields.

The generators E_z of $\mathfrak{MM}(Z)$ form a smooth family parameterized by $z \in Z$. For any element $F \in \mathfrak{MM}(Z)$ one can consider a smooth family $\mathbf{E}_z F \stackrel{\text{def}}{=} E_z \cdot F$, here $\cdot$ is the multiplication in $\mathfrak{MM}(Z)$. This defines a family of operators $\mathbf{E}_z$. The difference with the family $\mathbf{M}_\xi$ is the fact that $\mathbf{E}_z$ is defined on the whole $\mathfrak{MM}(Z)$, but does not send $\mathfrak{MM}(U)$ into $\mathfrak{MM}(U)$ if $z \notin U$.

Note that to make $\mathbf{M}_\xi$ defined on $\mathfrak{MM}(U)$ the open subset U should be small enough, but to make $\mathbf{E}_z$ act in $\mathfrak{MM}(U)$ the open subset U should be big enough. To make the situation more uniform, introduce the notion of V -conformal operator.

Definition 1.10. Consider partially defined mappings $F_k: \mathfrak{MM}(U_k) \rightarrow \mathfrak{MM}(Z)$, $k = 1, 2$, $U_{1,2} \subset Z$. Say that two such mappings are *equivalent* if they become equal when restricted to some $\mathfrak{MM}(U)$, $U \subset Z$.

Say that such a mapping is *smooth* if it sends smooth families in its domain of definition to smooth families. A *conformal operator* on $\mathfrak{MM}(Z)$ is a class of equivalence of smooth mappings.

¹⁶The motivation of codimension 2 is functoriality w.r.t. composition of mappings $\mathbf{m}_\eta$. The details are outside of the scope of this paper, see [12] instead. For the problems which appear otherwise see Example 1.9.

Let V be an open subset of Z . Say that a conformal operator is V -conformal if for any $U \subset U_0$ the corresponding mapping sends $\mathfrak{MM}(U)$ into $\mathfrak{MM}(U \cup V)$, and smooth families in $\mathfrak{MM}(U)$ to smooth families in $\mathfrak{MM}(U \cup V)$.

Obviously, operators $\mathbf{M}_\eta$ are $\emptyset$ -conformal, and $\emptyset$ -conformal operators form a monoid with unity. The operators $\mathbf{M}_\eta$ form a commutative subgroup of this monoid.

Remark 1.11. Note that two equivalent smooth mappings $\mathbf{c}, \mathbf{c}'$ coincide on the intersection of their domains. Indeed, they are both defined on some $\mathfrak{MM}(U)$, and coincide on this subset. However, any element F of $\mathfrak{MM}(Z)$ can be included in a smooth family F_s , $s \in S$, which intersects $\mathfrak{MM}(U)$. Now families $\mathbf{c}F_s$ and $\mathbf{c}'F_s$ are defined on a common open subset of S , and coincide there. Since smooth, they coincide where both are defined. Thus two images of F coincide if both mappings are defined on F .

In particular, any conformal operator can be maximally extended: it is represented by a smooth mapping with maximal possible domain of definition. In what follows we always consider such families only.

The particular definition of smoothness we used here is tuned to make the proofs of principal theorems (associativity relation and commutation relations) easier. One should expect that a definition written completely in terms of the geometry of the vector space $\mathfrak{MM}(Y)$ and the lattice of vector subspaces may work as well.

1.5. Families of operators $\mathbf{M}_\eta$. In this section we provide an abstract linear algebra framework formalizing properties of operators $\mathbf{h}_s$ from Section 0.5.

Instead of one meromorphic function η on Z consider a family $\mathbf{H}_\bullet$ of meromorphic functions $\mathbf{H}_s$ parameterized by $s \in S$ (in other words, there is a mapping $S \xrightarrow{\iota} X_{\mathcal{M}}$, and $\mathbf{H}_s$ is n/d if $i(s) = (n : d) \in X_{\mathcal{M}}$, here $\mathcal{M}$ is a line bundle on Z). For every fixed value of parameter $s \in S$ one obtains a meromorphic function $\mathbf{H}_s$ on Z , thus a linear operator $\mathbf{M}_{\mathbf{H}_s}$ defined on $\mathfrak{MM}(U_s)$ for a corresponding subset U_s .

Definition 1.12. For $z \in Z$ let

$$S^{(z)} = \{s \in S \mid \mathbf{H}_s \text{ has a pole or a zero at } z\}.$$

Say that $\mathbf{H}_s$, $s \in S$, *dominates* $U \subset Z$ if S cannot be covered by a finite number of sets $S^{(z)}$, $z \in U$.

Example 1.13. An arbitrary meromorphic function η on Z can be considered as a family parameterized by a 1-point set. It dominates the complement to its divisor of zeros and poles.

Consider a family $\mathbf{H}_s$, $s \in S$, which dominates U . Then for every meromorphic function $F \in \mathfrak{MM}(U)$ there is a non-empty open subset $S^{(F)}$ of S such that $\mathbf{M}_{\mathbf{H}_s}F$ is well-defined for $s \in S^{(F)}$. One can see an immediate advantage of families over individual functions: families dominate bigger open subsets than individual functions, and can even dominate a whole projective variety:

Example 1.14. Consider a projective variety Z , and a very ample line bundle $\mathcal{M}$ on it. There is a tautological meromorphic function H_s parameterized by $s \in S = X_{\mathcal{M}}$. Since $\mathcal{M}$ has no base points, subsets $S^{(z)}$ have smaller dimension than S , thus H_s dominates Z .

1.6. Conformal fields. The following property singles out family of conformal operators which behave similar to $\emptyset$ -conformal operators when considered together.

Definition 1.15. Consider a family $\mathbf{C}_s$, $s \in S$, of partially defined mappings each acting from a subset of $\mathfrak{MM}(Z)$ into $\mathfrak{MM}(Z)$. Suppose that for any $s \in S$ the mappings $\mathbf{C}_s$ is a conformal operator. Say that this family is *codominant* if

- (1) for any non-empty open subset V of Z there is a non-empty open dense subset $V^{(S)}$ of S such that the mapping $\mathbf{C}_s$ is a V -conformal operator in $\mathfrak{MM}(Z)$ if $s \in V^{(S)}$.
- (2) If V changes in a family, then $V^{(S)}$ can be chosen to change in a family.

In other words, a codominant family may increase support, but it will increase it by an arbitrary small amount, thus codominant families are closest analogues of $\emptyset$ -conformal operators. Obviously, the family $\mathbf{E}_z$, $z \in S \subset Z$ is codominant iff S is dense.

Consider a family H_s which dominates U . If $F \in \mathfrak{MM}(U)$, then $\mathbf{M}_{H_s}F$ is well-defined for s in a non-empty open subset $S^{(F)}$ of S , moreover, it can be written as a rational expression of $E_z^\bullet$ with regular functions on $S^{(F)}$ as coefficients. Consider this as a hint for a definition of a smooth family of conformal operators.

Another hint to understand what one wants from a family of V -conformal operators to be smooth is given by the locality property of a smooth family E_z of meromorphic functions on $\mathcal{M}(Z)$. These result in

Definition 1.16. Consider a family $\mathbf{C}(s)$, $s \in S$, each member of which is a V_s -conformal operator for a corresponding subset $V_s \subset Z$. Call this family a *conformal family* or a *conformal field on S acting on $\mathfrak{MM}(Z)$* if

- (1) There is a non-empty open dense subset $\tilde{S} \subset S \times Z$ such that $\mathbf{C}(s)$ is defined on $\mathfrak{MM}(U)$ if $\{s\} \times U \subset \tilde{S}$;
- (2) For any smooth family F_t in $\mathfrak{MM}(Z)$, $t \in T$, let $\mathbf{U}$ the subset of $S \times T$ where $\mathbf{C}(s)F_t$ is well defined. Then $\mathbf{U}$ is open, and the family $\mathbf{C}(s)F_t$ is smooth on $\mathbf{U}$.

Say that $\mathbf{C}(\bullet)$ *dominates* an open subset $U \subset Z$ if $\tilde{S}$ can be chosen to project onto U , $\mathbf{C}(\bullet)$ is *dominant* if it dominates Z . (With appropriate modification in the case of reducible S .)

Say that $\mathbf{C}(\bullet)$ is *defined* on $\tilde{S}$. Say that $\mathbf{C}(\bullet)$ is *smooth* if $\tilde{S}$ can be chosen to project onto S .

Example 1.17. If a family $H_\bullet$ of meromorphic functions dominates U , then $\mathbf{M}_{H_\bullet}$ is a $\emptyset$ -conformal field which dominates U .

Obviously, there is a maximal open $U \subset Z$ which is dominated by a conformal field $\mathbf{C}$, and it is dense. For any $F \in \mathfrak{MM}(U)$ there is a non-empty open subset $S^{(F)}$ of S where $\mathbf{C}(s)F$ is defined. Suppose that the family F_t from the last condition of the definition is in $\mathfrak{MM}(U)$. Then the subset U is not empty, thus the family $\mathbf{C}(s)F_t$ is meromorphic on $S \times T$.

Thus dominant conformal fields send smooth families to meromorphic ones! This is a first harbinger of the relation with the OPE expansions of the field theory.

The heuristic for the definition of smoothness is the following one: example of $\mathbf{M}_\eta$ shows that one cannot expect all the interesting operators to be defined on the whole vector space $\mathfrak{MM}(Z)$. The condition of smoothness ensures that the domain of definition does not jump down to $\{0\}$ when the parameter of the family changes. (It is not clear whether for such families the domain of definition *actually* always changes in a smooth family.)

1.7. Conformal field theory. The principal property of conformal fields is that they may be composed, but the composition of smooth fields is not necessarily smooth:

Theorem 1.18. *Suppose that $\mathbf{C}(s)$, $s \in S$, $\mathbf{C}'(s')$, $s' \in S'$, are conformal fields acting on $\mathfrak{MM}(Z)$, and $\mathbf{C}'$ is codominant. Then $\mathbf{C}(s) \circ \mathbf{C}'(s')$ is a conformal field on an open dense subset of $S \times S'$, and it is dominant if $\mathbf{C}$ and $\mathbf{C}'$ are dominant, and codominant if $\mathbf{C}$ is codominant.*

Proof. Suppose that $\mathbf{C}$ dominates U , $\mathbf{C}'$ dominates U' . Consider an arbitrary $F \in \mathfrak{MM}(U')$. Then $\mathbf{C}'(s')F$ is defined for s' in an open dense subset T' of S' , and depends smoothly on s' .

Since $\mathbf{C}'$ is codominant, one can decrease T' and U' in such a way that $\mathbf{C}'(s')F$ is in $\mathfrak{MM}(U)$ if $s' \in T'$ and $F \in \mathfrak{MM}(U')$. Hence $\mathbf{C}(s)\mathbf{C}'(s')F$ is defined for s in an open dense subset of S for any given $s' \in T'$.

Moreover, since the family $\mathbf{C}'(s')F$ is smooth, the family $\mathbf{C}(s)\mathbf{C}'(s')F$ is smooth on an open dense subset of $S \times S'$. If one lets F vary in some family F_t , $t \in T$, inside $\mathfrak{MM}(U')$, then the same is true for the family $\mathbf{C}(s)\mathbf{C}'(s')F_t$, which verifies the second condition of Definition 1.16.

Let $\mathbf{C}(s)$ be defined on $\mathfrak{MM}(V_s)$, here V_s give a family of open subsets of Z . Since $\mathbf{C}'$ is codominant, one can pick up $V_s^{(S')}$ to be the subset of S' from the definition of codominantness, let $\mathbf{V}$ be the corresponding total subset of $S \times S'$. One can conclude that $\mathbf{C}(s)\mathbf{C}'(s')F$ is defined if $F \in \mathfrak{MM}(V_s \cap U')$ and $(s, s') \in \mathbf{V}$. This verifies the first condition from Definition 1.16. If $\mathbf{C}$ is dominant, $\{V_s\}$ covers Z , which proves that $\mathbf{C}(s)\mathbf{C}'(s')$ dominates U' .

To show that the composition is codominant if $\mathbf{C}$ is is straightforward. $\square$

Note that one cannot expect the composition of smooth families to be smooth:

Example 1.19. Consider the family $\mathbf{M}_\xi$ parameterized by $\xi \in X_{\mathcal{M}}$ and the family $\mathbf{E}_z$ parameterized by $z \in Z$. If $\mathcal{M}$ is ample, both families are smooth, dominant and codominant. Moreover, there is a commutation relation

$$(1.2) \quad \mathbf{M}_\xi \circ \mathbf{E}_z = \langle \xi, z \rangle \mathbf{E}_z \circ \mathbf{M}_\xi,$$

here $\langle \xi, z \rangle$ is a scalar meromorphic function on $X_{\mathcal{M}} \times Z$ of taking the value of ξ at z . Since $\langle \xi, z \rangle$ has poles on $S = X_{\mathcal{M}} \times Z$, the set $\tilde{S}$ for $\mathbf{E}_z \circ \mathbf{M}_\xi$ cannot project on the points of S which are on the divisor of poles of $\langle \xi, z \rangle$.

Since the calculation of the above relationship is confusion-prone, let us outline it here. Given $F(\xi_1)$, $\mathbf{E}_z F(\xi_1) = \xi_1(z) F(\xi_1)$, thus

$$\mathbf{M}_\xi \mathbf{E}_z (F(\xi_1)) = \mathbf{M}_\xi (\xi_1(z) F(\xi_1)) = (\xi \xi_1)(z) F(\xi \xi_1) = \xi(z) \xi_1(z) F(\xi \xi_1).$$

Similarly, $\mathbf{E}_z \mathbf{M}_\xi (F(\xi_1)) = \xi_1(z) F(\xi \xi_1)$.

1.8. Homogeneous variant. The vector spaces in the lattice on which conformal operators act are *huge*, and the theory can be simplified a lot by reducing them. One of the variants is to consider only the semisimple part with respect to the natural action of the multiplicative group $\mathbb{G}_m = \text{GL}(1)$ on $\mathcal{M}(Z)$.

Let $\lambda \in \mathbb{G}_m$ acts on $X_{\mathcal{M}}$ by $(n : d) \rightarrow (\lambda n : d)$. Consider only those meromorphic functions on $X_{\mathcal{M}}$ which are homogeneous w.r.t. this action. In other words, for some $\delta \in \mathbb{Z}$

$$F(\lambda n : d) = \lambda^\delta F(n : d)$$

(this gives some additional restriction on degrees of polyhomogeneity of G and G' in (1.1)). Similarly, consider only conformal fields of degree $\delta' \in \mathbb{Z}$, i.e., operators which send homogeneous functions of degree δ into functions of degree $\delta + \delta'$.

Example 1.20. The degree of $\mathbf{M}_\bullet$ is 0, the degree of $\mathbf{E}_\bullet$ is 1. Composition adds degrees. In particular, the family $\mathbf{E}_z \circ \mathbf{E}_{z'}^{-1}$, $(z, z') \in Z \times Z$, is of degree 0.

When one considers conformal families of homogeneity degree 0, the representation of $\text{GL}(1)$ in the vector space (or lattice of vector spaces) is split into a direct sum of homogeneous components. Thus it is reasonable to restrict attention to one homogeneous component.

Note that if Z has a distinguished point z_0 , then multiplication by E_{z_0} identifies all the different homogeneous components. Thus it is enough to consider the action in the subspace of homogeneity degree 0.

1.9. Residues. Recall that given a meromorphic function G on an algebraic variety S , and an irreducible hypersurface $D \subset S$, one may consider *Leray residues* $\text{Res}_D^{(n)} G$ of G along D . If S is a direct product $D_0 \times C$ with C being a curve, and $D = D_0 \times \{c_0\}$, $c_0 \in C$, then $\text{Res}_D^{(n)} G|_{d \in D}$ is $-n$ -th Laurent coefficients of $G|_{\{d\} \times C}$ along C at c_0 . In general, some additional data make Laurent residues into meromorphic functions on D , and one can easily describe how to transform these functions if these additional data are changed.

Say, let α be a meromorphic function on S which has a zero of the first order at D , $\omega = d\alpha$. ω is a meromorphic 1-form on S which has no zero or pole at D , and vanishes when pulled back to D . Let v be a meromorphic vector field on S such that $\langle \omega, v \rangle = 1$, and v has no pole or zero on D . Using these data, one can describe the Leray residue of G as the residue along v of the differential form $\omega(s) \alpha(s)^{n-1} G$ at generic points of the divisor D . Fix such data (α, v) , in the future consider residues w.r.t. these data only. Note that one can also use these data to define the residue on $D \times T$ of a function on $S \times T$ for any irreducible variety T .

Residues of positive order vanish if G is regular at generic points of D . In this case residues of negative order are Taylor coefficients of G in the direction of v . Thus residues with $n \geq 1$ describe the singular part of G at D . Abusing notations, call residues with positive and negative n taken together *Laurent coefficients* of G at D (numbered with opposite numbers).

Note that $\text{Res}_D^{(0)} G$ coincides with restriction to D if G is smooth at generic points of D . However, $\text{Res}_D^{(0)} G$ is defined even if G has a pole at D . The biggest contrast with restriction is that if D' is a different hypersurface, then for any irreducible component D'' of $D \cap D'$ one can consider $\text{Res}_{D''}^{(k)} \text{Res}_D^{(l)} G$ and $\text{Res}_{D''}^{(l)} \text{Res}_{D'}^{(k)} G$. These two functions coincide¹⁷ if no pole of G contains D'' , however, this is true no more if G is not regular on D'' . This observation is the core of *conformal commutation relations*.

In particular, if $T \subset S$ is not contained in D , this does not imply that

$$\left(\text{Res}_D^{(n)} G \right) |_T = \text{Res}_{D \cap T}^{(n)} (G|_T).$$

This identity holds only if $D \cap T$ is not contained in the divisor of poles of G .

Example 1.21. Consider a function $G = \frac{1}{x-z}$ on the plane, let $T = \{x = 0\}$, $D = \{z = 0\}$. Then $\text{Res}_D^{(1)} G = 0$, but $G|_T$ has a pole at the origin, thus $\text{Res}_0^{(1)} (G|_T) = -1$.

Multiplying G by a power of α , one can see that

Amplification 1.22.

$$\left(\text{Res}_D^{(n)} G \right) |_T = \text{Res}_{D \cap T}^{(n)} (G|_T)$$

if G is defined and finite on $D \cap T$, or if the only component of the divisor of poles of G which contain $D \cap T$ is D .

Consider a meromorphic family F_s in $\mathfrak{MM}(Z)$ parameterized by S . Let D be an irreducible hypersurface in S . Then F_s induces a meromorphic function on $S \times X_{\mathcal{L}}$ for $\mathcal{L} \geq \mathcal{L}_0$. Thus for any $n \in \mathbb{Z}$ one can consider the Leray residue $\text{Res}_{D \times X_{\mathcal{L}}}^{(n)} (F_s)$. This residue is a meromorphic function on $D \times X_{\mathcal{L}}$.

¹⁷Assume that the vector fields v, v' associated to D, D' are tangent to the other hypersurface of a pair (D, D') .

Consider a generic inclusion $X_{\mathcal{L}} \rightarrow X_{\mathcal{L}'}$ for $\mathcal{L} < \mathcal{L}'$. This gives an inclusion $S \times X_{\mathcal{L}} \rightarrow S \times X_{\mathcal{L}'}$, which is compatible with families F_s on these spaces. To show that the residues are compatible too, note that this is true if F_s is defined on $D \times X_{\mathcal{L}}$ for $\mathcal{L} \geq \mathcal{L}_0$. Otherwise F_s has a pole of the *same* order d on $D \times X_{\mathcal{L}} \subset S \times X_{\mathcal{L}}$ for $\mathcal{L} \geq \mathcal{L}_0$, and multiplication of F by α^d finishes the argument.

Thus $\text{Res}_{D \times X_{\mathcal{L}}}^{(n)}(F_s)$ provides a meromorphic family $\text{Res}_D^{(n)} F_s$ on $\mathfrak{MM}(Z)$ parameterized by D .

Definition 1.23. Call this family $\text{Res}_D^{(n)} F_s$ the *residue* of F_s at $D \subset S$.

Consider a conformal field $\mathbf{C}(s)$ on S acting on $\mathfrak{MM}(Z)$. Let D be an irreducible hypersurface in S . Suppose the $\mathbf{C}$ dominates $U \subset Z$. For any $F \in \mathfrak{MM}(U)$ one can consider $\text{Res}_D^{(n)}(\mathbf{C}(s)F)$.

Definition 1.24. The n -th *fine residue* $\text{RES}_D^{(n)} \mathbf{C}$ of $\mathbf{C}$ on D is the family of mappings $\mathfrak{MM}(U) \rightarrow \mathfrak{MM}(Z)$ parameterized by D , and defined by $F \mapsto \text{Res}_D^{(n)}(\mathbf{C}(s)F)$.

The big problem with this notion is that $\text{Res}_D^{(n)}(\mathbf{C}(s)F)$ does not depend smoothly on F , compare Example 1.28 below. One needs a coarser notion that that: below we will find a restriction of this operator to a vector subspace which is smooth.

For any smooth family F_t , $t \in T$, one can consider the residue of $\mathbf{C}(s)F_t$ on $D \times T$. Note that if $T_1 \subset T$, then the residue on $D \times T$ restricted to $D \times T_1$ does not necessarily coincide with the residue of the restriction of $\mathbf{C}(s)F_t$ on $S \times T_1$. However, they coincide if $D \times T_1$ is not contained in the divisor of poles of $\mathbf{C}(s)F_t$.

Next step is to study for which T and T_1 the operations of taking a residue and of restriction commute. Suppose that $\mathbf{C}$ is defined on $\tilde{S} \subset S \times Z$. Let $\tilde{R} = (S \times Z) \setminus \tilde{S}$, let $\tilde{R}_0$ be the union of irreducible components of $\tilde{R}$ which differ from $D \times Z$. Consider an open subset $\tilde{D}_0 = (D \times Z) \setminus \tilde{R}_0$ of $D \times Z$. Let D_0 be the projection of $\tilde{D}_0$ to D , Z_0 be the projection of $\tilde{D}_0$ to Z .

If $F_{t_0} \in Z_0$ for $t_0 \in T_1$, then for a generic $s_0 \in S$ the family $\mathbf{C}(s)F_t$, $s \in S$, $t \in T_1$, is meromorphic, and the only poles of this family near (s_0, t_0) are poles at $D \times T_1$ (if any). Since $t_0 \in T$ as well, same is true for the family $\mathbf{C}(s)F_t$ on $S \times T$. Amplification 1.22 implies that in this case taking the residue along D and restriction to T_1 commute indeed.

Definition 1.25. Define a parameterized by D family of partially defined linear operator $\text{Res}_D^{(n)} \mathbf{C}$ in the lattice $\{\mathfrak{MM}(U)\}_{U \subset Z}$ by

$$\left(\text{Res}_D^{(n)} \mathbf{C}\right) F = \text{Res}_D^{(n)}(\mathbf{C}F), \quad F \in \mathfrak{MM}(Z_0).$$

The following statement is obvious:

Proposition 1.26. Suppose that $\mathbf{C}$ is defined on $\tilde{S} \subset S \times Z$. Let $\tilde{R} = (S \times Z) \setminus \tilde{S}$, let $\tilde{R}_1$ be the union of irreducible components of $\tilde{R}$ which differ from $D \times Z$. Consider

an open subset $\tilde{D}_0 = (D \times Z) \setminus \tilde{R}_0$ of $D \times Z$. Let D_0 be the projection of $\tilde{D}_0$ to D , Z_0 be the projection of $\tilde{D}_0$ to Z .

Then $\text{Res}_D^{(n)} \mathbf{C}$ is a conformal field on D acting in $\mathfrak{MM}(Z)$ which is defined on $\tilde{D}_0$, dominates Z_0 , and is smooth on D_0 .

Remark 1.27. Heuristically, the above description of the conformal field $\text{Res}_D^{(n)} \mathbf{C}(s)$ can be boiled down to: define it by $\left(\text{Res}_D^{(n)} \mathbf{C}(s)\right) F = \text{Res}_D^{(n)} (\mathbf{C}(s) F)$ for generic F , and extend smoothly to non-generic F . The necessity of such double limit comes from the fact that $\text{Res}_D^{(n)} (\mathbf{C}(s) F)$ does not depend smoothly on F .

Example 1.28. Consider $\mathbf{C}(s) = \mathbf{M}_{\frac{1}{z-s}}$, $z, s \in \mathbb{A}^1$. Then $\text{Res}_{s=0}^{(1)} \left(\mathbf{M}_{\frac{1}{z-s}} E_{z_0}\right)$ is 0 for generic z_0 , -1 if $z_0 = 0$. Yet worse, $\text{Res}_{s=0}^{(0)} \left(\mathbf{M}_{\frac{z+s}{z-s}} E_{z_0}\right)$ is 1 for generic z_0 , -1 if $z_0 = 0$, and $\text{Res}_{s=s_0}^{(n)} \left(\mathbf{M}_{\frac{z+s}{z-s}} E_{z_0}\right) = 0$ for any s_0 and $n > 0$.

Remark 1.29. Let $\mathbf{C}$ is smooth on $S_0 \subset S$. Then $D_0 \supset S_0 \cap D$. Note that one cannot deduce anything else about D_0 and Z_0 by looking at the set which $\mathbf{C}$ dominates, and at S_0 . Indeed, if S is $\mathbb{A}^2$ with coordinates s_1, s_2 , Z is $\mathbb{A}^1$ with coordinate z , and $\tilde{S}$ is given by $s_1 \neq 0$ and $s_2 = zs_1$, then the field is smooth at $s_1 \neq 0$, but the residue at $s_1 = 0$ is smooth on $s_2 \neq 0$.

Similarly, one cannot assume that the residue is codominant even if the initial field is.

Remark 1.30. Note that the system of axioms we use here ensures that for any given element of $\mathfrak{MM}(Z)$ (or even any family of elements)

$$\left(\text{Res}_D^{(n)} \mathbf{C}\right) F = 0 \text{ for } n \gg 0.$$

However, these axioms do not ensure the fact that the above equality is valid *uniformly* in F .

To improve the system of axioms to exclude the above pathologies one may need to consider as support not open subsets $U \subset Z$, but closed subschemes of $Z \setminus U$, similarly for the definition set of a conformal field. Yet better alternative is to consider the Hilbert scheme of 0-dimensional subschemes of Z (compare with Section 3).

Since we do not want to delve into such details, note that for all examples of conformal fields considered here, the above assumption is true.

Remark 1.31. Note that the above equality for fine residue

$$\left(\text{RES}_D^{(n)} \mathbf{C}\right) F = 0 \text{ for } n \gg 0$$

is manifestly non-uniform in F . Indeed, consider again $\mathbf{C}(s) = \mathbf{M}_{\frac{1}{z-s}}$, $Z = S = \mathbb{A}^1$, $D = \{0\}$. Let F be the k -th derivative of evaluation function $E_0^{(d/dz)^k}$. Then $(\text{RES}_D^{(n)} \mathbf{C}) F = 0$ iff $n > k + 1$.

Remark 1.32. Note that though conformal operators are linear, and residues may be written as integrals in the case $\mathbb{K} = \mathbb{C}$, conformal fields not necessarily commute with taking a residue. Indeed, fix $\xi_0 \in \mathcal{M}(Z)$, let ξ_0 has a simple pole at hypersurface $H \subset Z$. Restrict what we did in Example 1.19 to $\{\xi_0\}$: let $\mathbf{C} = \mathbf{M}_{\xi_0}$, consider a family E_z , $z \in Z$, of elements of $\mathfrak{MM}(Z)$. Then $\text{Res}_H^{(0)} E_z = E_z|_{z \in H}$, so $\mathbf{C}(\text{Res}_H^{(0)} E_z)$ is not defined. On the other hand,

$$\text{Res}_H^{(0)} (\mathbf{C} E_z) = \text{Res}_H^{(0)} \xi_0(z) E_z = \text{Res}_H^{(-1)} E_z = E_z^v|_{z \in H},$$

here v is an appropriate (meromorphic) vector field on Z . (Proposition 3.35 will provide a way to fix this non-commutativity.)

1.10. Fusions. Consider now two smooth conformal fields $\mathbf{C}(s)$, $s \in S$, and $\mathbf{C}'(s')$, $s' \in S'$, both acting on $\mathfrak{MM}(Z)$. The composition $\mathbf{C}(s) \circ \mathbf{C}'(s')$ is smooth on an open subset of $S \times S'$, after a blow up of the complement to this subset one may assume that this composition is smooth on a complement to a collection of divisors D_i , $i \in I$. Fixing a local equation α_i and a transversal vector field v_i for each D_i , one obtains families $\text{Res}_{D_i}^{(n)} (\mathbf{C}(s) \circ \mathbf{C}'(s'))$, $n \geq 1$. If i -th family is not smooth on D_i , one can consider further blow-ups and residues, and so on, until all the residues of high iteration will be smooth.

Definition 1.33. The *fusions* of conformal fields $\mathbf{C}$ and $\mathbf{C}'$ are the elements of the defined above collection of conformal fields.

Remark 1.34. Since composition of conformal fields is associative, one should be able to describe some “pseudo-associativity” restrictions on the fusions of fusions, which are what remains of associativity relation. The actual description of these relations (i.e., an axiomatic conformal field theory, compare [1]) is outside of the scope of this paper.

However, note that Section 3.5 shows that a wide class of conformal fields has pairwise fusions, but no “new” triple-product fusions, which is the base of conformal field theory.

Note also that we use the word *fusion* is a slightly more general way than it is used in the conformal field theory: their the families $\mathbf{C}$ and $\mathbf{C}'$ are defined on the same parameter space $S = S'$; moreover, $\dim S = 1$, and the divisor D coincides with the diagonal in S . Under this restriction our definition of fusions coincides with the classical one; moreover, the mentioned above absense of “new” triple-product fusions means that the singularities of a tripple product, such as $\mathbf{C}(s) \mathbf{C}'(s') \mathbf{C}''(s'')$, happen only on divisors $\{s = s'\}$, $\{s' = s''\}$, and $\{s = s''\}$. Moreover, one can easily check that this condition is equivalent to the conformal associativity relation.

1.11. Localization. The main difference between conformal fields and families of operators in a vector space is that the conformal fields are only *partially defined*, i.e., they act only on small subspaces of the lattice $\mathfrak{M}\mathcal{M}(U)$, $U \subset Z$, of subspaces of a vector space $\mathfrak{M}\mathcal{M}(Z)$. To bridge this gap, one may consider a direct limit over the family U_i elements of which become smaller and smaller. To do this, take a point $z_0 \in Z$, and consider a limit w.r.t. the lattice of open subsets of Z which contain z_0 . (To simplify notations, in what follows z_0 is supposed to be smooth.)

Note that this process is absolutely analogous to the definition of localization of Z at z_0 , however, we will see that our axioms lead to consideration of formal completion $\widehat{Z}$ of Z at z_0 instead. Indeed, one can localize any conformal field from Z to any open subset U of Z . However, this does not change the theory a lot, since $\mathcal{M}(U) = \mathcal{M}(Z)$, thus the only things which change are domains of definition of families of meromorphic functions on $\mathcal{M}(Z)$, and dominantness and codominantness.

On the other hand, $\cap_{U \ni z_0} \mathfrak{M}\mathcal{M}(U)$ contains only generators $E_{z_0}^\bullet$, and coincides with the field generated by these elements. While one can calculate the value of an element of $\mathfrak{M}\mathcal{M}(Z)$ at “almost any” meromorphic function on Z , it also makes sense to calculate values of elements of $\cap_{U \ni z_0} \mathfrak{M}\mathcal{M}(U)$ on “almost any” formal power series at z_0 . One can see that, first, we achieved the settings of the formal approach (see Section 0.10), thus the in the conformal field theory *localization* of Z at z_0 leads to study of functions on *completion* of Z at z_0 .

Second, the generators $E_{z_0}^\bullet$ of $\mathfrak{M}\mathcal{M}(\widehat{Z}) \stackrel{\text{def}}{=} \cap_{U \ni z_0} \mathfrak{M}\mathcal{M}(U)$ are enumerated by multiindices $K = (k_1, \dots, k_d)$, $d = \dim Z$, $k_i \geq 0$. Thus $\mathfrak{M}\mathcal{M}(\widehat{Z})$ consists of rational functions of symbols $E^{(K)}$, $K \in \mathbb{Z}_{\geq 0}^d$.

At last, the lattice of subsets of $\widehat{Z}$ is trivial, so is what remains of the lattice of subspaces in $\mathfrak{M}\mathcal{M}(Z)$ when one restricts attention to $\mathfrak{M}\mathcal{M}(\widehat{Z}) \subset \mathfrak{M}\mathcal{M}(Z)$. In particular, those conformal fields which may be restricted to $\mathfrak{M}\mathcal{M}(\widehat{Z})$ act there as usual families of linear operators. Note however, that not all conformal operators can be restricted to $\mathfrak{M}\mathcal{M}(\widehat{Z})$, the operator should be V -conformal for any $V \ni z_0$, and should dominate some neighborhood of z_0 . In particular, any dominant $\emptyset$ -conformal field acts in $\mathfrak{M}\mathcal{M}(\widehat{Z})$ by families of linear operators meromorphically depending on the parameter. For example, M_ξ acts in $\mathfrak{M}\mathcal{M}(\widehat{Z})$ if ξ has no zero or pole at z_0 .

Consider now an arbitrary codominant conformal field $\mathbf{C}(s)$, $s \in S$. For any $V \ni z_0$ there is a non-empty dense subset $S^{(V)}$ such that the operators $\mathbf{C}(s)$ are V -conformal operators if $s \in S^{(V)}$. A priori the intersection $\cap_{V \ni z_0} S^{(V)}$ may be empty, but it is possible to show that it is a non-empty closed subset $S^{\{z_0\}}$ of S if S is projective. One may conclude that operators $\mathbf{C}(s)$ and all their derivatives in s act on $\mathfrak{M}\mathcal{M}(\widehat{Z})$ if $s \in S^{\{z_0\}}$, and $\mathbf{C}$ is defined at (s, z_0) . The last condition gives an open subset of S ,

so in general one cannot conclude that the set of such s is non-empty, even if $\mathbf{C}$ is dominant. However, for the family $\mathbf{E}_z$ this holds.

In general, one can see that the restriction to $\mathfrak{MM}(\widehat{Z})$ of a codominant conformal field $\mathbf{C}$ on a projective variety S is defined as on an open subset $\widehat{U}$ of a formal completion $\widehat{S}^{\{z_0\}}$ in S of a closed subvariety $S^{\{z_0\}}$ of S . If $\mathbf{C}$ is dominant, the open subset is an intersection of non-empty open subset U of S with $\widehat{S}^{\{z_0\}}$. However, this intersection does not necessarily contain closed points of Z , but it is non-empty as a scheme. Indeed, functions on a formal completion are formal power series, when some closed subsets are removed, one should consider the subalgebra of fractions of such formal power series with denominators having zero at the thrown away divisors only. (Note that this complication does not appear if one considers the field $\mathbf{E}_z$.)

If S is not projective, consider any projective completion $\bar{S}$ of S . The family $\mathbf{C}$ can be extended to $\bar{S}$ (possibly with a loss of smoothness). However, smoothness was not used in the above arguments, thus, as above, a dominant codominant family $\mathbf{C}$ can be restricted to an appropriate localization of formal neighborhood of a closed subvariety $S^{\{z_0\}}$ of $\bar{S}$. If one needs to localize $\widehat{S}^{\{z_0\}}$ outside of $S^{\{z_0\}}$, one needs to consider formal Laurent series instead of formal Taylor series.

Consider operators $\mathbf{M}_\xi$ in more details. Assume that ξ has no zero or pole at z_0 , so can be written as a formal power series near z_0 . Thus the subgroup of $\{\mathbf{M}_\xi\}$ which acts in $\mathfrak{MM}(\widehat{Z})$ is generated by multiplications by formal power series $\xi = \sum_K \xi_K H^K$, here $H = (\eta_1, \dots, \eta_d)$, η_i are local coordinates near z_0 . (One should consider ξ which give meromorphic functions on Z only.)

Restrict ourselves to the homogeneous theory. Suppose that ξ has no zero at z_0 . Since $\mathbf{M}_\xi$ is of homogeneity degree 0, $\xi = \text{const}$ gives a trivial operator in grading 0, thus one may normalize ξ at any point, say, by $\xi(z_0) = 1$. This means that only the *divisor* of ξ is relevant, thus one obtains an action of the group of principal divisors on Z on $\mathfrak{MM}(\widehat{Z})$. This action is given by the product formula, which expresses Taylor coefficients of $\xi\xi'$ in terms of Taylor coefficients of ξ and ξ' . In this action each $E^{(K)}$ is mapped to a linear combination of $E^{(L)}$, $l_i \leq k_i$ for any i .

As far as the divisor does not pass through z_0 , this action preserves polynomials in $E^{(K)}$. However, the action of divisor which pass through z_0 is not defined.

Consider $\mathbf{E}_z$ now. Obviously, one cannot multiply an element of $\mathfrak{MM}(\widehat{Z})$ by E_z unless $z = z_0$. However, the action of $\mathbf{E}_z$ can be written in terms of the (formal) Taylor series of $\mathbf{E}_z$ near $z = z_0$:

$$E_z = \sum_K H^K(z) E^{(K)},$$

here a monomial H^K in $(\eta_1, \dots, \eta_d)$ is considered as a function of $z \approx z_0$. Multiplication by such an expression gives not a rational function of $E^{(K)}$, but a power series.

Particular Taylor coefficients of $\mathbf{E}_z$ in z near $z = z_0$ are just operators of multiplication by $E^{(K)}$, thus preserve polynomials in $E^{(K)}$.

Proposition 1.35. *The operators $\mathbf{E}_z$ considered as power series near $z = z_0$ act in the field of fractions of formal power series in letters $E^{(K)}$. Same if one considers homogeneous fractions of power series, with $\deg E^{(K)} = 1$.*

Similarly

Proposition 1.36. *The operators ξ_D are defined for any principal divisor of Z which does not pass through z_0 . They act in any algebra: polynomials in $E^{(K)}$, formal power series in $E^{(K)}$ and fractions thereof.*

2. EXAMPLES OF CONFORMAL FAMILIES

Consider the simplest possible example, when Z is a point z_0 . In such a case the only generator in $\mathfrak{MM}(Z)$ is $E \stackrel{\text{def}}{=} E_{z_0}$. The vector subspace of homogeneous functions of degree d is generated by E^d . Thus a homogeneous conformal operator C of degree δ is described by sequences of numbers C_d , $d \in \mathbb{Z}$, such that $C(E^d) = C_d E^{d+\delta}$. Restricted to functions of homogeneity degree 0, this becomes a number. Similarly, conformal fields are as dull as families of numbers, i.e., scalar functions.

Thus any real theory starts in dimension 1, with the simplest case being of genus $g = 0$. The biggest difference of this case with the higher-dimensional situation is the fact that the group of all divisors is generated by the subset of effective divisors of degree 1. Moreover, if genus $g = 0$, then the same is true for principal divisors. This allows one to consider $\mathbf{M}_\xi$ for $\xi \in X_{\mathcal{O}(1)}$ only, since the rest can be described by compositions of such operators.

In this section we investigate conformal fields in the particular case $Z = \mathbb{P}^1$.

2.1. Fields based on $\mathbb{P}^1$. Consider the simplest possible case of the family $\mathbf{M}_\xi$, with $\xi \in X_{Z, \mathcal{L}}$, and $Z = \mathbb{P}^1$, $\mathcal{L} = \mathcal{O}(1)$. In this case $X_{Z, \mathcal{L}}$ is 3-dimensional, and described by coordinates $(a : b : c : d)$ giving a meromorphic function $z \mapsto \frac{az+b}{cz+d}$. If one considers the homogeneous case only, then multiplication by a non-0 constant does not matter, so one can consider $\widehat{\xi} = (\xi_n, \xi_d) = ((a : b), (c : d)) \in \mathbb{P}^1 \times \mathbb{P}^1$ instead, here $\xi_n \ominus \xi_d$ is the divisor of ξ .

For a 0-cycle $\mathfrak{y} = \sum k_i z_i$ on Z with coefficients in $\mathbb{Z}$ one can define $\mathbf{E}_\mathfrak{y} = \prod \mathbf{E}_{z_i}^{k_i}$ (recall that $\mathbf{E}_z$ is the operator of multiplication by E_z , and different $\mathbf{E}_z$ commute). Consider the homogeneous case again, then one needs to consider 0-cycles of degree 0, and the simplest such a cycle is given by $z_+ \ominus z_-$, it is parameterized by $Z \times Z$. In the case $Z = \mathbb{P}^1$ one can see that the set of parameters $\mathbb{P}^1 \times \mathbb{P}^1$ is the same as in the case of the simplest family $\mathbf{M}_\xi$.

Obviously, $\mathbf{M}_\xi$, $\xi \in X_{\mathbb{P}^1, \mathcal{O}(1)}$, form a dominant and codominant conformal field, similarly for $\mathbf{E}_\eta$ if η runs through $Z \times Z$, as above. If one considers compositions of $\mathbf{M}_\xi$ with itself, then it is a smooth family, and compositions commute, the same for $\mathbf{E}_\eta$. Repeat again that though we restrict ourselves to the smallest-dimension families $X_{\mathcal{O}(1)}$ and $Z \times Z$, *any* operator M_ξ can be described as a composition of such simplest $\mathbf{M}_\xi$, similarly for any 0-cycle of degree 0.

Consider now compositions $\mathbf{M}_\xi \circ \mathbf{E}_\eta$ and $\mathbf{E}_\eta \circ \mathbf{M}_\xi$. As we have seen it already,

$$\mathbf{M}_\xi \circ \mathbf{E}_\eta = \langle \xi, \eta \rangle \mathbf{E}_\eta \circ \mathbf{M}_\xi, \quad \langle \xi, \eta \rangle \stackrel{\text{def}}{=} \xi(z_+) / \xi(z_-) \text{ if } \eta = z_+ \ominus z_-.$$

Indeed, $\mathbf{M}_\xi E_z \stackrel{\text{def}}{=} \xi(z) E_z$, $\mathbf{E}_{z_+ \ominus z_-} E_z \stackrel{\text{def}}{=} E_{z_+} E_{z_-}^{-1} E_z$. Thus

$$\mathbf{M}_\xi \circ \mathbf{E}_{z_+ \ominus z_-} (E_z) = \mathbf{M}_\xi (E_{z_+} E_{z_-}^{-1} E_z) = \xi(z) \xi(z_+) \xi^{-1}(z_-) E_{z_+} E_{z_-}^{-1} E_z,$$

$$\mathbf{E}_{z_+ \ominus z_-} \circ \mathbf{M}_\xi (E_z) = \xi(z) \mathbf{E}_{z_+ \ominus z_-} (E_z) = \xi(z) E_{z_+} E_{z_-}^{-1} E_z.$$

Moreover, one can see that the family $\mathbf{E}_\eta \circ \mathbf{M}_\xi$ is smooth. Since $\langle \xi, \eta \rangle = \frac{z_+ - \xi_n}{z_+ - \xi_d} \cdot \frac{z_- - \xi_d}{z_- - \xi_n}$, $\mathbf{M}_\xi \circ \mathbf{E}_\eta$ has fusions on the hypersurfaces $z_- = \xi_n$ and $z_+ = \xi_d$. Only the $n = 1$ fusions are not zero. For example, consider $z_- = \xi_n$. On this hypersurface the fusion is

$$(z_- - \xi_d) \frac{z_+ - z_-}{z_+ - \xi_d} \mathbf{E}_{z_+ \ominus z_-} \circ \mathbf{M}_\xi$$

However, this conformal field has a pole at the intersection $z_+ = \xi_d$ with the other divisor $z_+ = \xi_d$ with the residue

$$(z_- - z_+) (z_+ - z_-) \mathbf{E}_{z_+ \ominus z_-} \circ \mathbf{M}_\xi,$$

which is smooth, here $\xi(z) = \frac{z - z_-}{z - z_+}$. One gets the same result when one considers the different order of taking residues along divisors. Note that the factor $(z_- - z_+) (z_+ - z_-)$ is just the normalization by the determinant of conormal bundle to the intersection of divisors.

Note that any $\eta = z_+ \ominus z_-$ corresponds to a meromorphic function $\xi(\eta)$ (up to proportionality) with a pole at z_+ and a zero at z_- . Let $\mathbf{M}_\eta \stackrel{\text{def}}{=} \mathbf{M}_{\xi(\eta)}$, $\mathbf{F}_\eta = \mathbf{E}_\eta \circ \mathbf{M}_\eta$. Then $\mathbf{F}_\eta$ coincides with the above smooth double-fusion without the normalizing coefficient:

$$\mathbf{F}_{(z_+, z_-)} = \mathbf{E}_{z_+ \ominus z_-} \circ \mathbf{M}_{\frac{z_- - z_-}{z_- - z_+}}.$$

One can describe the action of $\mathbf{F}_{(z_+, z_-)}$ on a rational function $F(E_{z_1}, \dots, E_{z_k})$ via

$$E_z \mapsto \frac{z - z_-}{z - z_+} E_{z_+} E_{z_-}^{-1} E_z.$$

Note that the action on E_z^{-1} is

$$E_z^{-1} \mapsto \frac{z - z_+}{z - z_-} E_{z_+} E_{z_-}^{-1} E_z^{-1}.$$

Thus this family is defined on $z \neq z_+$, $z \neq z_-$ (this is another way to see that the family is smooth).

2.2. Fusions of F_η . Since any M_ξ may be represented as a composition of M_{ξ_i} with $\xi_i \in X_{\mathcal{O}(1)}$, and any E_η can be represented as composition of E_{η_i} , $\eta_i = z_{i+} \ominus z_{i-}$, similarly for families, the fusions of the fields which are compositions of M_ξ and E_η may be expressed via fusions of elementary fields. Thus the most interesting fusions are the fusions between F_η and $F_{\eta'}$.

One can write the action of $F_\eta F_{\eta'}$ on E_z as

$$E_z \mapsto \frac{z'_+ - z_-}{z'_+ - z_+} \cdot \frac{z'_- - z_+}{z'_- - z_-} \cdot \frac{z - z_-}{z - z_+} \cdot \frac{z - z'_-}{z - z'_+} E_{z_+} E_{z_-}^{-1} E_{z'_+} E_{z'_-}^{-1} E_z,$$

here $\eta = z_+ \ominus z_-$, $\eta' = z'_+ \ominus z'_-$. Similarly, the action on E_z^{-1} is

$$E_z^{-1} \mapsto \frac{z'_+ - z_-}{z'_+ - z_+} \cdot \frac{z'_- - z_+}{z'_- - z_-} \cdot \frac{z - z_+}{z - z_-} \cdot \frac{z - z'_+}{z - z'_-} E_{z_+} E_{z_-}^{-1} E_{z'_+} E_{z'_-}^{-1} E_z^{-1},$$

One can see that this conformal field is defined if

$$\frac{z'_+ - z_-}{z'_+ - z_+} \cdot \frac{z'_- - z_+}{z'_- - z_-}, \quad \frac{z - z_+}{z - z_-} \cdot \frac{z - z'_+}{z - z'_-}, \quad \frac{z - z_-}{z - z_+} \cdot \frac{z - z'_-}{z - z'_+}$$

are not infinity, thus if

$$z'_+ \neq z_+, z'_- \neq z_-, z \neq z_-, z \neq z_+, z \neq z'_-, z \neq z'_+.$$

Correspondingly, $F_\eta F_{\eta'}$ is smooth on

$$z'_+ \neq z_+, z'_- \neq z_-.$$

Thus $F_\eta F_{\eta'}$ has fusions at two divisor, both fusions are non-zero for $n = 1$ only. Again, these fusions have residues on the intersection with the other divisor, and the fusion of the second order is defined on the surface

$$z'_+ = z_+, z'_- = z_-.$$

of codimension 2. Plugging into the formula for $F_\eta F_{\eta'}$, one can describe this fusion as

$$E_z \mapsto \left(\frac{z - z_-}{z - z_+} \right)^2 E_{z_+}^2 E_{z_-}^{-2} E_z,$$

one can continue taking the fusions obtaining newer 2-parametric conformal fields as fusions of the older ones.

2.3. A different reduction. One has seen that fusions of conformal fields $\mathbf{M}_\xi$ and $\mathbf{E}_\eta$ can be described as a reduction of the smooth field $\mathbf{E}_\eta \circ \mathbf{M}_\xi$ on subvariety $\xi_n = z_-$, $\xi_d = z_+$. Consider now a different reduction $\xi_n = z_+$, $\xi_d = z_-$:

Definition 2.1. Let $\mathbf{G}_\eta = \mathbf{E}_\eta \circ \mathbf{M}_\xi$ with $\xi(z) = \frac{z-z_+}{z-z_-}$, $\eta = z_+ \ominus z_-$.

Obviously, $\mathbf{G}_\eta$ is a smooth dominant and codominant family sending

$$E_z \mapsto \frac{z-z_+}{z-z_-} E_{z_+} E_{z_-}^{-1} E_z.$$

The restriction of $\mathbf{G}_\eta$ to the diagonal $z_+ = z_-$ is the identity operator id . One can write the action of $\mathbf{G}_\eta \mathbf{G}_{\eta'}$ on E_z as

$$E_z \mapsto \frac{z'_+ - z_+}{z'_+ - z_-} \cdot \frac{z'_- - z_-}{z'_- - z_+} \cdot \frac{z - z_+}{z - z_-} \cdot \frac{z - z'_+}{z - z'_-} E_{z_+} E_{z_-}^{-1} E_{z'_+} E_{z'_-}^{-1} E_z,$$

here $\eta = z_+ \ominus z_-$, $\eta' = z'_+ \ominus z'_-$. Similarly, the action on E_z^{-1} is

$$E_z^{-1} \mapsto \frac{z'_+ - z_+}{z'_+ - z_-} \cdot \frac{z'_- - z_-}{z'_- - z_+} \cdot \frac{z - z_-}{z - z_+} \cdot \frac{z - z'_-}{z - z'_+} E_{z_+} E_{z_-}^{-1} E_{z'_+} E_{z'_-}^{-1} E_z^{-1},$$

One can see that this conformal field is defined if

$$\frac{z'_+ - z_+}{z'_+ - z_-} \cdot \frac{z'_- - z_-}{z'_- - z_+}, \quad \frac{z - z_-}{z - z_+} \cdot \frac{z - z'_-}{z - z'_+}, \quad \frac{z - z_+}{z - z_-} \cdot \frac{z - z'_+}{z - z'_-}$$

are not infinity, thus if

$$z'_+ \neq z_-, \quad z'_- \neq z_+, \quad z \neq z_-, \quad z \neq z_+, \quad z \neq z'_-, \quad z \neq z'_+.$$

Correspondingly, $\mathbf{G}_\eta \mathbf{G}_{\eta'}$ is smooth on

$$z'_+ \neq z_-, \quad z'_- \neq z_+.$$

Thus $\mathbf{G}_\eta \mathbf{G}_{\eta'}$ has fusions at two divisor, both fusions are non-zero for $n = 1$ only. Calculate the residue at $z'_+ = z_-$:

$$E_z \mapsto (z_- - z_+) \frac{z'_- - z_-}{z'_- - z_+} \cdot \frac{z - z_+}{z - z'_-} E_{z_+} E_{z'_-}^{-1} E_z.$$

This formula can be significantly simplified by a renormalization

$$\mathcal{G}(z_+, z_-) = -\frac{1}{z_+ - z_-} \mathbf{G}_{z_+ \ominus z_-}.$$

Then the fusion of $\mathcal{G}(z_+, z_-)$ with $\mathcal{G}(z'_+, z'_-)$ at $z'_+ = z_-$ is given by

$$E_z \mapsto \frac{1}{z'_- - z_+} \cdot \frac{z - z_+}{z - z'_-} E_{z_+} E_{z'_-}^{-1} E_z = \mathcal{G}(z_+, z'_-) E_z.$$

Note, however, that the conformal field $\mathcal{G}(z_+, z_-)$ is no longer smooth, it has a pole at $z_+ = z_-$ with the residue $\mathbf{G}_{z_+ \ominus z_+} = -\text{id}$.

One can conclude that the non-smoothness of the algebra generated by conformal fields $\mathcal{G}(z_+, z_-)$ is completely describable by relations

$$\begin{aligned} \text{Res}_{z_+=z_-} \mathcal{G}(z_+, z_-) &= -\text{id}, \\ \text{Res}_{z'_+=z_-} \mathcal{G}(z_+, z_-) \mathcal{G}(z'_+, z'_-) &= \mathcal{G}(z_+, z'_-), \\ \text{Res}_{z_+=z'_-} \mathcal{G}(z_+, z_-) \mathcal{G}(z'_+, z'_-) &= -\mathcal{G}(z'_+, z_-). \end{aligned}$$

The last two relationship look similar to commutation relations of matrix elements. The results of Section 4.3 will show that this is indeed true after taking Laurent coefficients in z_+ and z_- at 0.

Note also that the conformal fields $\mathcal{G}(z_+, z_-)$ and $\mathcal{G}(z'_+, z'_-)$ commute:

$$\mathcal{G}(z_+, z_-) \mathcal{G}(z'_+, z'_-) = \mathcal{G}(z'_+, z'_-) \mathcal{G}(z_+, z_-)$$

on the subset of the set of parameters where both sides are defined. We leave verification of this statement to the reader.

Remark 2.2. Note that renormalization from $\mathbf{G}$ to $\mathcal{G}$ is not invariant w.r.t. automorphisms of $\mathbb{P}^1$. To see the invariant form of fusion relation, note that the residue describes the singular part of a conformal field near a divisor.

Rewrite the above calculations as

$$\mathbf{G}_{z_+ \ominus z_-} \mathbf{G}_{z'_+ \ominus z'_-} = \frac{\frac{z'_+ - z_+}{z_+ - z'_-}}{\frac{z_- - z'_+}{z'_- - z_-}} \mathbf{E}_{z_-}^{-1} \mathbf{E}_{z'_+} \mathbf{G}_{z_+ \ominus z'_-}, = \lambda(z'_+, z'_-, z_+, z_-) \mathbf{E}_{z_-}^{-1} \mathbf{E}_{z'_+} \mathbf{G}_{z_+ \ominus z'_-},$$

if $z_- \approx z'_+$ this is

$$\mathbf{G}_{z_+ \ominus z_-} \mathbf{G}_{z'_+ \ominus z'_-} \approx \lambda(z'_+, z'_-, z_+, z_-) \mathbf{G}_{z_+ \ominus z'_-},$$

here $\lambda(a, b, c, d)$ is the double ratio of 4 points on $\mathbb{P}^1$.

One should understand the above equality in the following sense: for any $F \in \mathfrak{MM}(\mathbb{P}^1)$ the left-hand side and the right-hand side have same residues at (an open subset of) $z_- = z'_+$ when applied to F , same for $z_+ = z'_-$. Note that the double ratio has poles exactly at these two divisors.

Remark 2.3. Results Section 4.3 will show that the Laurent coefficients of the family $\mathcal{G}(s, r)$ (and its smooth part $\mathcal{G}(s, r) - \frac{1}{r-s}$) give rise to standard descriptions of the Lie algebra $\mathfrak{gl}(\infty)$ and its central extension $\mathfrak{g}(\infty)$.

2.4. Further reductions. Consider a further reduction of the family $\mathbf{E}_\eta M_\xi$. Restrict $\mathfrak{h} = (z_+, z_-)$ to have $z_- = 0$, and consider only ξ which have a pole at ∞ . The second restriction makes this family dominant on $\mathbb{A}^1 \subset \mathbb{P}^1$, the first one codominant outside of 0 only.

In fact it is better to abandon the restriction that the homogeneity degree is 0, and consider $\mathbf{E}_0^{-1}$ as a normalization of degree only. Thus consider the family $\Phi(z_+, z_-) =$

$\mathbf{E}_{z_+} \mathbf{M}_{1/(z-z_-)}$. It is now dominant and codominant smooth family of homogeneity degree 1 on $\mathbb{A}^1$. Note that

$$\mathbf{M}_{1/(z-z_-)} \mathbf{E}_{z_+} = \frac{1}{z_+ - z_-} \mathbf{E}_{z_+} \mathbf{M}_{1/(z-z_-)},$$

and one can easily calculate fusions of $\Phi(z_+, z_-)$ and $\Phi(z'_+, z'_-)$ using this relationship.

Make a reduction $z_+ = z_-$ in $\Phi(z_+, z_-)$ (or consider the residue of $\mathbf{M}_{1/(z-z_-)} \mathbf{E}_{z_+}$):

$$\varphi(z_0) \stackrel{\text{def}}{=} \Phi(z_0, z_0) = \mathbf{E}_{z_0} \mathbf{M}_{1/(z-z_0)}.$$

It is still a dominant codominant smooth family on $\mathbb{A}^1$. One can easily see that $\varphi(z) \varphi(z')$ is not smooth, thus φ has a fusion with itself of order $n = 1$, which is proportional to $\mathbf{E}_{z_0}^2 \mathbf{M}_{1/(z-z_0)}^2$, and that

$$\varphi(z) \varphi(z') = -\varphi(z') \varphi(z).$$

One may also calculate fusions of φ with itself of higher order, and see that one gets higher and higher degrees of $\mathbf{E}$ and $\mathbf{M}$. Thus the field φ is a 1-parametric analogue of the field from Section 2.2.

Two other fields are defined by reductions of $\mathbf{E}_\eta \mathbf{M}_\xi$, $\eta = z_+ \ominus z_-$, with either $z_+ = 0$, or ξ having a pole at ∞ . They are again dominant on $\mathbb{A}^1$ only:

$$\psi(z_0) = \mathbf{E}_{z_0} \mathbf{M}_{z-z_0}, \quad \psi^+(z_0) = \mathbf{E}_{z_0}^{-1} \mathbf{M}_{1/(z-z_0)}.$$

These fields have the same commutation relations as φ :

$$\begin{aligned} \psi(z) \psi(z') &= -\psi(z') \psi(z), \\ \psi(z) \psi^+(z') &= -\psi^+(z') \psi(z), \\ \psi^+(z) \psi^+(z') &= -\psi^+(z') \psi^+(z), \end{aligned}$$

moreover, the families $\psi(z) \psi(z')$ and $\psi^+(z) \psi^+(z')$ are smooth, and $\psi(z) \psi^+(z')$ has the only fusion

$$\psi(z) \psi^+(z') \approx \frac{1}{z' - z} \text{id}$$

Thus the fields ψ and ψ^+ are 1-parametric analogues of the field from Section 2.3.

Let us spell out the calculation of the products $\psi(r) \psi^+(s)$ and $\psi^+(s) \psi(r)$.

Suppose that $F \in \mathfrak{MM}(\mathbb{A}^1)$, $\xi \in \mathcal{M}(\mathbb{A}^1)$, let z runs through $\mathbb{A}^1$. Then

$$(2.1) \quad (\psi^+(s) F)(\xi) = \xi(r) \frac{1}{(s-r)\xi(s)} F\left(\frac{1}{z-s}(z-r)\xi(z)\right),$$

and

$$\begin{aligned} (\psi(r) \psi^+(s) F)(\xi) &= \xi(r) \frac{1}{(s-r)\xi(s)} F\left(\frac{1}{z-s}(z-r)\xi(z)\right), \\ (\psi^+(s) \psi(r) F)(\xi) &= \frac{1}{(r-s)\xi(s)} \xi(r) F\left((z-r) \frac{1}{z-s} \xi(z)\right), \end{aligned}$$

thus $\psi(r) \psi^+(s) = -\psi^+(s) \psi(r)$, if $r \neq s$. It is also easy to calculate the pole of $\psi(r) \psi^+(s)$ at $r \approx s$ for a fixed F :

$$(\psi(r) \psi^+(s) F)(\xi) = \frac{1}{(s-r)} F(\xi) + [\text{a smooth function}].$$

Indeed, fix $\xi(z) \in \mathcal{M}(\mathbb{A}^1)$, consider $F\left(\frac{z-r}{z-s}\xi(z)\right)$ as a function of r and s . Show that for a generic ξ this function is a well-defined meromorphic function of r and s . Moreover, it has no pole at $r \approx s$.

It is easy to deduce this given an explicit formula for F as a rational expression of $E_z^\bullet$, $z \in \mathbb{A}^1$, but one can also obtain it from the heuristic for $\mathfrak{MM}(Z)$ as being compatible collections of meromorphic functions on $X_{\mathcal{L}}$, $\mathcal{L}$ running through ample sheaves on Z (see Section 1.1 and [12]).

As in Section 1.1, in the case $Z = \mathbb{A}^1$ one should consider manifolds $X_{\mathcal{O}(k)}$, $k > 0$, and mappings between them given by multiplication of numerator and denominator by $P_l(z)$, $P_l(z)$ being a polynomial in z of degree l . The “function” F is modeled by meromorphic functions $F^{(k)}$ on $X_{\mathcal{O}(k)}$ compatible with mappings which correspond to a generic P . In particular, the divisor of poles of $F^{(k+1)}$ does not contain the image of $X_{\mathcal{O}(k)}$ in $X_{\mathcal{O}(k+1)}$ for $P_1 = x - s$ and a generic s . This means that for generic s the expression $F\left(\frac{z-r}{z-s}\xi(z)\right)$ has no pole at $r = s$ for a generic $\xi \in X_{\mathcal{O}(k)}$.

Hence

$$\psi(r) \psi^+(s) = \frac{1}{(s-r)} \mathbf{1} + O(1),$$

here $\mathbf{1}$ is the identity operator. Thus the fields ψ and ψ^+ are 1-parametric analogues of the field from Section 2.3.

Similarly,

$$\begin{aligned} \varphi(r) \varphi(s) &= \frac{1}{(s-r)} \varphi_2 + O(1), \\ (\varphi_2(r) F)(\xi) &= \xi^2(r) F\left(\frac{\xi(z)}{(z-r)^2}\right), \quad \xi \text{ is rational.} \end{aligned}$$

One can interpret these formulae as saying that $\mathbf{1}$ is a *fusion* of ψ and ψ^+ , and φ_2 is a *fusion* of φ with itself.

In Section 5.6 we will show that ψ^+ and ψ give rise to standard fermion creation and destruction operators.

2.5. Vertex operators. Consider how the transition to the formal completion $\widehat{Z}$ of $Z = \mathbb{P}^1$ at 0 can be written in an explicit form. Recall that in the case of a completed manifold the lattice disappears, thus conformal fields become well-defined families of operators on the vector space $\mathfrak{MM}(\widehat{Z})$. As in the case of general Z , $\mathfrak{MM}(\widehat{Z})$ is generated by $E_0^\bullet$, but in 1-dimensional case the index $\bullet$ takes values in $\mathbb{Z}_{\geq 0}$. (Similarly, in the homogeneous situation one may normalize an element $\xi \in \mathcal{M}(\widehat{Z})$ to satisfy $\xi(0) = 1$, thus $E_0 \equiv 1$, so the set of indices of $E_0^\bullet$ can be restricted to be $\mathbb{N} = \mathbb{Z}_{>0}$.) Denote these generators of $\mathfrak{MM}(\widehat{Z})$ by $e^{(k)}$, $k \in \mathbb{Z}_{\geq 0}$.

The action of M_ξ is defined on $\mathfrak{MM}(\widehat{Z})$ only if ξ has no pole or zero at 0. Let ξ has a Taylor coefficients ξ_k , $\xi = \sum \xi_k z^k$, then M_ξ acts as a field automorphism of $\mathfrak{MM}(\widehat{Z})$ generated by

$$M_\xi: e^{(k)} \mapsto \sum_{l=0}^k e^{(l)} \xi_{k-l}.$$

In the particular case $\xi(z) = -s/(z-s)$, one gets

$$M_{-s/(z-s)}: t_k \mapsto \sum_{l=0}^k s^{l-k} e^{(l)},$$

However, to define the action of operators E_z “outside” of $z = 0$ one needs to consider formal power series, since only the operators $E_0^\bullet$ act in $\mathfrak{MM}(\widehat{Z})$. So consider

$$E_z = \sum_k \frac{1}{k!} z^k E_z^{(k)}, \quad E_z = \sum_k \frac{1}{k!} z^k E_z^{(k)}$$

as formal power series in z , and see that the coefficients of the latter series act in the field $\mathfrak{MM}(\widehat{Z})$, $E_z^{(k)}$ acts as multiplication by $e^{(k)}$.

Both these actions preserve the subalgebra $\mathcal{V}_{\geq 0} \subset \mathfrak{MM}(\widehat{Z})$ of polynomials in $e^{(k)}$. Assigning to each $e^{(k)}$ grading 1, one obtains a grading $\mathcal{V}_{\geq 0} = \bigoplus_{l \geq 0} \mathcal{V}_l$. Allowing negative powers of $e^{(0)}$, one may define $\mathcal{V}_l$ with $l < 0$. Let $\mathcal{V} = \bigoplus_{l \in \mathbb{Z}} \mathcal{V}_l$.

The usual way to obfuscate the simple formulae given above is to choose a different set of generators in the algebra $\mathcal{V}$. The above set of generators $e^{(k)}$ corresponds to taking Taylor coefficients $e^{(k)}$ at 0 of an element $\xi(z)$ of $\mathcal{M}(\mathbb{P}^1)$ without a pole at 0. These Taylor coefficient generate some field of functions on $\mathcal{M}(\mathbb{P}^1)$.

Up to this moment one could work in any characteristic, since though the above formulae contain denominators, in fact $\frac{e^{(k)}}{k!}$ is well defined as a Taylor coefficient of given meromorphic function ξ at 0.

Now assume $\text{char } \mathbb{K} = 0$, and take different coordinates: for a fixed $\xi \in \mathcal{M}(\mathbb{P}^1)$, $\xi(0) \in \mathbb{K}^\times$, write

$$\xi(z) = T \exp \sum_{k>0} t_k z^k, \quad T, t_k \in \mathbb{K}.$$

Considered as functions of ξ , the symbols $T, t_1, \dots$ are meromorphic functions on $\mathcal{M}(\mathbb{P}^1)$ (since they can be expressed as polynomials in $e^{(k)}$), and $e^{(k)}$ can be expressed as polynomials in $T, t_1, \dots$. Thus $T, t_1, \dots$ form a different collection of independent generators of $\mathcal{V}_{\geq 0}$. The degree of T is 1, of t_k is 0, $\mathcal{V}_l = T^l \mathbb{C}[t_1, \dots]$.

One can easily write the considered above operators in terms of these new coordinates:

Proposition 2.4. (1) *The vector subspace $\mathcal{V}$ is invariant w.r.t. operators $\mathbf{M}_{z-s}$, $s \neq 0$. The action of $\mathbf{M}_{z-s}$ can be written as*

$$\mathbf{M}_{z-s} T = -sT, \quad \mathbf{M}_{z-s} t_k = t_k - \frac{1}{ks^k}, \quad \mathbf{M}_{z-s}(ab) = (\mathbf{M}_{z-s}a)(\mathbf{M}_{z-s}b).$$

(2) *The vector subspace $\mathcal{V}$ is invariant w.r.t. operators $\mathbf{E}_s$, $s \approx 0$. In other words, if one writes the Taylor series for $\mathbf{E}_s$:*

$$\mathbf{E}_s v \approx \sum_{l \geq 0} \varepsilon_l(v) s^l,$$

then the operators ε_k preserve the subspace $\mathcal{V}$. The action of $\mathbf{E}_s$ on $\mathcal{V}$ can be described as

$$\mathbf{E}_s T = e^{\sum_l t_l s^l} T^2, \quad \mathbf{E}_s t_k = T e^{\sum_l t_l s^l} t_k, \quad \mathbf{E}_s(ab) = (\mathbf{E}_s a)b = a(\mathbf{E}_s b).$$

Writing $\mathbf{M}_{z-s}|_{\mathcal{V}_l}$, $\mathbf{E}_s|_{\mathcal{V}_l}$ as Laurent series in s , one gets

$$\mathbf{M}_{z-s} = (-s)^l \exp \left(- \sum_{k>0} \frac{1}{ks^k} \frac{\partial}{\partial t_k} \right),$$

$$\mathbf{E}_s = T \exp \sum_{k>0} s^k t_k$$

(recall that in the Section 0 these operators were called $a^{-1}(s)$, $b(s)$). Taking into account definitions of φ , ψ , and ψ^+ , one can conclude that

Corollary 2.5. *When considered as Laurent series in s and restricted on the subspace $\mathcal{V}_l$, the operators $\mathbf{M}_{z-s}$, $\mathbf{E}_s$, $\varphi(s)$, $\psi(s)$, and $\psi^+(s)$ become the standard vertex*

operators:

$$\begin{aligned}\varphi(s) &= T(-s)^{-l} \exp\left(\sum_{k>0} s^k t_k\right) \exp\left(\sum_{k>0} \frac{1}{k s^k} \frac{\partial}{\partial t_k}\right), \\ \psi(s) &= T(-s)^l \exp\left(\sum_{k>0} s^k t_k\right) \exp\left(-\sum_{k>0} \frac{1}{k s^k} \frac{\partial}{\partial t_k}\right), \\ \psi^+(s) &= T^{-1}(-s)^{-l} \exp\left(-\sum_{k>0} s^k t_k\right) \exp\left(\sum_{k>0} \frac{1}{k s^k} \frac{\partial}{\partial t_k}\right).\end{aligned}$$

2.6. Analogy with flag space. Instead of interpreting standard vertex operators as restrictions of conformal fields to the space of local functionals $\mathcal{V}$, one can interpret smooth families $\mathbf{M}_\xi$, $\mathbf{E}_z$, φ , ψ , and ψ^+ as *geometric realizations* of the standard vertex operators. The relation of the geometric realization to the realization by differential operators (of infinite order) is parallel to the relation of geometric realization of representations of semisimple groups to representations in the category $\mathcal{O}$.

Indeed, take a representation of G in sections of a bundle $\mathcal{L}$ over the flag space $\mathcal{F}$. The Lie algebra $\mathfrak{g}$ acts by vector fields v_X on $\mathcal{F}$ (lifted to vector fields on $\mathcal{L}$), $X \in \mathfrak{g}$. Take an open Schubert cell U in $\mathcal{F}$. It is isomorphic to an affine space with coordinates T_k . One can trivialize the bundle $\mathcal{L}|_U$, so the vector fields v_X become differential operators of order 1 in variables T_k . These differential operators preserve the space of polynomials $k[T_k]$, and the resulting representation of $\mathfrak{g}$ is in category $\mathcal{O}$.

One can easily see that the set $\mathcal{M}_0(\mathbb{P}^1)$ of rational functions without a pole at 0 is an analogue of U , t_k are analogues of T_k . This makes the conformal fields into analogues of exponents of elements of Lie algebra (thus elements of the Lie group), and makes $\mathcal{M}(\mathbb{P}^1)$ into an analogue of the flag space.

3. HILBERT SCHEMES AND CODOMINANT CONFORMAL FIELDS

In this section we show that appropriate projective bundles over the Hilbert schemes of points in Z are providing important *models* of the space $\mathfrak{MM}(Z)$, and prove that codominant families automatically satisfy conformal associativity relations, and vertex commutation relations.

Blow-ups we consider are going to be blow-ups with a center in an arbitrary closed subscheme, not necessarily blow-ups with centers in smooth submanifolds.

3.1. Big families. Describe some families in $\mathfrak{MM}(Z)$ such that any other family can be *almost* pushed through one of these families. First, note that for any finite collection of points $A = \{z_1, \dots, z_m\} \subset Z$ any meromorphic function F on $\mathcal{M}(Z)$ can be determined by the restriction to the subset U_A of $\mathcal{M}(Z)$ consisting of functions without a pole at $z_i \in A$. Let Z_0 be any subscheme of Z of finite length with support at the points $z_i \in A$. Say that F can be pushed through Z_0 , if $F(\xi)$, $\xi \in U_A$,

is determined by $\xi|_{Z_0}$. Since any $E_{z_0}^{v_1 \dots v_d}$ can be pushed through a suitable finite subscheme of Z with support at $\{z_0\}$, any element $F \in \mathfrak{MM}(Z)$ can be pushed through a finite subscheme. Obviously, there is a minimal such subscheme which will be denoted $\text{Supp } F$.

Second, for any vector space V over $\mathbb{K}$ the symmetric algebra $S^\bullet V$ is filtered by $\mathfrak{P}_k V \stackrel{\text{def}}{=} \bigoplus_{i=0}^k S^i V$. The field of fractions of $S^\bullet V$ is filtered by $\bar{\mathcal{Q}}_k V \stackrel{\text{def}}{=} \mathfrak{P}_k V / (\mathfrak{P}_k V \setminus \{0\})$. Suppose $\dim V < \infty$, consider $\mathfrak{P}_k V$ as an affine algebraic variety over $\mathbb{K}$, denote the projectivization $\mathbb{P}(\mathfrak{P}_k V \oplus \mathfrak{P}_k V)$ by $\mathcal{Q}_k V$. It is a finite dimensional projective space, it contains projective subspaces $\mathcal{Q}_k^\infty V$ and $\mathcal{Q}_k^0 V$ which corresponds to $\mathbb{P}(\mathfrak{P}_k V \oplus 0)$ and $\mathbb{P}(0 \oplus \mathfrak{P}_k V)$, let $\bar{\mathcal{Q}}_k V = \mathcal{Q}_k V \setminus \mathcal{Q}_k^\infty V$. Points of $\bar{\mathcal{Q}}_k V$ over $\mathbb{K}$ map to elements of $\bar{\mathcal{Q}}_k V$.

For a vector space W denote the dual space by $W^\vee$, then $\bar{\mathcal{Q}}_k W^\vee$ may be identified with the set of rational functions on W of degree $\leq k$. Let Z_0 be a finite subscheme of length m in a quasiprojective variety Z . Then $\mathfrak{D}_{Z_0} \stackrel{\text{def}}{=} \Gamma(Z_0, \mathcal{O}_{Z_0})$ is an m -dimensional vector space, and given an element $\varphi \in \mathcal{Q}_k \mathfrak{D}_{Z_0}^\vee$, one can associate to it an element of $\mathfrak{MM}(Z)$ given by $\mathcal{M}(Z) \ni \xi \mapsto \varphi(\xi|_{Z_0})$ (defined if Z_0 does not intersect divisor of poles of ξ). This defines a family of meromorphic functions on $\mathcal{M}(Z)$ parameterized by $\mathcal{Q}_k \mathfrak{D}_{Z_0}^\vee$.

Example 3.1. Given distinct points $z_i \in Z$, $1 \leq i \leq m$, one can associate to them a subscheme $Z_0 = \bigcup_i \{z_i\}$. The vector space $\mathfrak{D}_{Z_0}^\vee$ has a natural basis E_{z_i} , and $\mathcal{Q}_k \mathfrak{D}_{Z_0}^\vee$ consists of rational functions of E_{z_i} , $1 \leq i \leq m$ of degree at most k . Let $U \subset Z^m$ consists of m -tuples of distinct points on Z , then one can associate a projective space $\mathcal{Q}_k \mathfrak{D}_{Z_0}^\vee$ to any $A \in U$. Let $Q_{k,m}$ be the total space of the corresponding projective bundle over U . There is a natural admissible family of meromorphic functions on $\mathcal{M}(Z)$ parameterized by $Q_{k,m}$.

To investigate what happens when some of the points z_i collide, note that there is an action of the symmetric group $\mathfrak{S}_m$ on $Q_{k,m}$ which preserves the family, so one should expect that the family may be pushed down to an appropriate “desingularization” of the quotient by the group action. It is indeed so if one takes the principal irreducible component $Z^{[m]}$ of the *Hilbert scheme* $\text{Hilb}^m(Z)$ of m points on Z as such a “desingularization”. Recall that points of $\text{Hilb}^m(Z)$ are finite subschemes of length m in Z , the principal irreducible component is one which contains m -tuples of distinct points¹⁸. It is known that $Z^{[m]}$ and $\text{Hilb}^m(Z)$ are projective varieties if Z is, and are quasiprojective in general case of quasiprojective Z .

For a given $Z_0 \in Z^{[m]}$ the vector space $\mathfrak{D}_{Z_0}$ is a fiber of a locally free *tautological structure sheaf* $\mathfrak{D}$ on $Z^{[m]}$. Applying $\mathcal{Q}_k$ to the dual sheaf, one obtains a projective bundle $\mathfrak{Q}_{k,m}$ over $Z^{[m]}$. Denote by the same symbol $\mathfrak{Q}_{k,m}$ the total space of this bundle. It is a quasiprojective variety, and is projective if Z is. Define $\bar{\mathfrak{Q}}_{k,m}$ similarly.

¹⁸Note that if $\dim Z \leq 2$ and Z is smooth, the Hilbert scheme is irreducible (and smooth).

Similarly, one can define corresponding bundles over $\text{Hilb}^m(Z)$, denote them $\mathfrak{Q}_{k,m}^{\text{Hilb}}$ and $\mathring{\mathfrak{Q}}_{k,m}^{\text{Hilb}}$.

We have seen that there is a natural family of meromorphic functions on $\mathcal{M}(Z)$ parameterized by $\mathfrak{Q}_{k,m}$. Denote this family by $\mathfrak{F}_{k,m}$.

Proposition 3.2. *The family $\mathfrak{F}_{k,m}$ is strictly meromorphic on $\mathfrak{Q}_{k,m}$ and strictly smooth on $\mathring{\mathfrak{Q}}_{k,m}$.*

Proof. First, the variety $Q_{k,m}$ from Example 3.1 is a finite covering of an open subset of $\mathfrak{Q}_{k,m}$. We have seen that the family becomes admissible when pulled back to $Q_{k,m}$.

To show that $\mathfrak{F}_{k,m}$ induces a meromorphic function on $\mathfrak{Q}_{k,m} \times X_{\mathcal{L}}$ for $\mathcal{L} \geq \mathcal{L}_0$ note that for any finite-dimensional vector space V the evaluation function $(f, v) \mapsto f(v)$ is well-defined and meromorphic on $Q_k V^\vee \times V$. Thus for any vector bundle B over S the evaluation function is well-defined on $Q_k B^\vee \times_S B$ and is meromorphic.

The next step of the proof requires the following fact. Represent the locally free sheaf $\mathfrak{D}$ on $Z^{[m]}$ as the sheaf of sections of a m -dimensional vector bundle $B \xrightarrow{\pi} Z^{[m]}$. The fibers of this bundle have a structure of algebra, and the multiplication mapping $*$: $B \times_{Z^{[m]}} B \rightarrow B$ is regular.

Fact . *One can find a regular scalar function φ on B and a regular mapping i : $B \rightarrow B$ over $Z^{[m]}$ such that $b \in B$ is invertible w.r.t. $*$ iff $\varphi(b) \neq 0$, and then $b^{-1} = \varphi(b)^{-1} i(b)$. Here $b \mapsto b^{-1}$ is taking the ring inverse in fibers of B .*

Proof. One may suppose that Z is affine. Fix a function f on Z . Define a function

$$\Sigma_l f: (z_1, \dots, z_m) \mapsto \sigma_l(f(z_1), \dots, f(z_m))$$

on Z^m , here σ_l is the l -th elementary symmetric function. This function on Z^m is invariant w.r.t. the action of the symmetric group $\mathfrak{S}_m$ on Z^m , thus it induces a function on *Chow fraction* $Z^m/\mathfrak{S}_m$ (compare [5]). Composing with the natural mapping $Z^{[m]} \rightarrow Z^m/\mathfrak{S}_m$, one obtains a regular function on $Z^{[m]}$, which will be denoted by the same symbol $\Sigma_l f: Z_0 \mapsto \Sigma_l f(Z_0)$.

If we fix a finite-dimensional vector space V of functions on Z , then $\Sigma_l: (f, Z_0) \mapsto \Sigma_l f(Z_0)$ is a regular function on $V \times Z^{[m]}$. On the other hand, for any fixed $Z_0 \in Z^{[m]}$ there is a natural mapping of restriction $r_{Z_0}: V \rightarrow \mathfrak{D}_{Z_0}$. Moreover, for a fixed $Z_{00} \in Z^{[m]}$ one can find an m -dimensional space V such that the above mapping is an isomorphism of vector spaces for $Z_0 \in U_{Z_{00}}$, $U_{Z_{00}}$ being a neighborhood of Z_{00} . Thus Σ_l can be lifted to a regular function on $\pi^{-1}U_{Z_{00}}$. It is clear that different choices of V result in the liftings which coincide on $\pi^{-1}(U_{Z_{00}} \cap U)$, thus everywhere (U is from Example 3.1).

This shows that Σ_l can be considered as a function on B . Note that if Z_0 contains points $z_1, \dots, z_m$ (counted with appropriate multiplicities), and $f \in \mathfrak{D}_{Z_0}$, then $\Sigma_m f = f(z_1) \dots f(z_m)$, thus $f \in \mathfrak{D}_{Z_0}$ is invertible in $\mathfrak{D}_{Z_0}$ iff Σ_m does not vanish on (Z_0, f) . Moreover,

$$f^{*m} - \Sigma_1(Z_0, f) f^{*m-1} + \dots \pm \Sigma_{m-1}(Z_0, f) f \mp \Sigma_m(Z_0, f) \mathbf{1} = 0$$

if $Z_0 \in U$, $f \in \mathfrak{D}_{Z_0}$, $*$ denotes multiplication in B , $\mathbf{1}$ the unit element of $\mathfrak{D}_{Z_0}$, and $\bullet^{*d}$ denotes taking d -th power w.r.t. multiplication in B . Since the left-hand side is a regular mapping from B to B , this is true for any $Z_0 \in Z^{[m]}$.

In other words,

$$\Sigma_m(Z_0, f) \mathbf{1} = f^*(\Sigma_{m-1}(Z_0, f) \mathbf{1} - \Sigma_{m-e}(Z_0, f) f + \cdots \pm \Sigma_1(Z_0, f) f^{*m-2} \mp f^{*m-1}).$$

Recall that for invertible $f \in \mathfrak{D}_{Z_0}$ one has $\Sigma_m(Z_0, f) \neq 0$, thus

$$f^{-1} = \Sigma_m(Z_0, f)^{-1} (\Sigma_{m-1}(Z_0, f) \mathbf{1} - \Sigma_{m-e}(Z_0, f) f + \cdots \pm \Sigma_1(Z_0, f) f^{*m-2} \mp f^{*m-1}).$$

□

Lemma 3.3. *The expression*

$$\mathcal{E}: (F, n, d) \rightarrow F(n/d)$$

defines a meromorphic function $\mathcal{E}_B$ on $\mathcal{Q}_k B^\vee \times_{Z^{[m]}} B \times_{Z^{[m]}} B$.

Proof. Indeed, $(n, d) \rightarrow n/d$ is a regular dominant mapping $B \times B^{\text{inv}} \rightarrow B$, here B^{inv} is an open subset of invertible elements of B . Thus $F(n/d)$ gives a meromorphic function on an open subset of $\mathcal{Q}_k B^\vee \times_{Z^{[m]}} B \times_{Z^{[m]}} B$, thus on $\mathcal{Q}_k B^\vee \times_{Z^{[m]}} B \times_{Z^{[m]}} B$. □

For a line bundle $\mathcal{L}$ on Z and a finite subscheme Z_0 of Z there is a natural *trivialization* mapping $\Gamma(Z, \mathcal{L}) \rightarrow \mathfrak{D}_{Z_0}$ defined up to multiplication by an invertible element of $\mathfrak{D}_{Z_0}$. Say that $\mathcal{L}$ *exhausts* Z_0 if this mapping is surjective, say that $\mathcal{L}$ is *m-ample* if it exhausts all the finite subschemes of length m . Now Lemma 3.3 shows that

Corollary 3.4. *If $\mathcal{L}$ is m-ample, then $\mathfrak{F}_{k,m}$ induces a meromorphic function on $\mathfrak{Q}_{k,m} \times X_{\mathcal{L}}$.*

Amplification 3.5. *If $\mathcal{L}$ is m-ample, then for any $q \in \mathring{\mathfrak{Q}}_{k,m}$ the divisor of poles of the above meromorphic function on $\mathfrak{Q}_{k,m} \times X_{\mathcal{L}}$ does not contain $\{q\} \times X_{\mathcal{L}}$.*

Proof. It is enough to prove a similar statement about the meromorphic function $\mathcal{E}_B$ from Lemma 3.3. Suppose that $q = (Z_0, q_{Z_0}) \in \mathcal{Q}_k B^\vee$ is such that $\{q\} \times \mathfrak{D}_{Z_0} \times \mathfrak{D}_{Z_0}$ is in the divisor of poles of $\mathcal{E}_B$. In the notations of Lemma 3.3 put $d = \mathbf{1}_{Z_0}$. One can see that q considered as a meromorphic function on $\mathfrak{D}_{Z_0}$ is ∞ , thus q is on the infinity $\mathcal{Q}_k^\infty B^\vee = \mathcal{Q}_k B^\vee \setminus \mathring{\mathcal{Q}}_k B^\vee$ of $\mathcal{Q}_k B^\vee$. □

To finish the proof of Proposition 3.2, note that

Lemma 3.6. *Let $\mathcal{L}_1$ be a very ample bundle on Z . Then any $\mathcal{L} \geq \mathcal{L}_1^m$ is m-ample.*

Proof. It is enough to prove that $\mathcal{O}(m)$ is m-ample on $\mathbb{P}^l$ for any l . This in turn follows from the (well-known and easy-to-prove, see [4]) fact that any ideal of $\mathbb{K}[[t_1, \dots, t_l]]$ of codimension m contains $\mathfrak{m}^m$, $\mathfrak{m}$ being the maximal ideal of $\mathbb{K}[[t_1, \dots, t_l]]$. □

Thus the divisor of poles of $\mathfrak{F}_{k,m}$ is flat over $\mathring{\mathfrak{Q}}_{k,m}$ indeed. □

Remark 3.7. Consider a blow-up $\tilde{\mathcal{Q}}_k B^\vee$ of $\mathcal{Q}_k B^\vee$ in $\mathcal{Q}_k^\infty B^\vee$. Let $\tilde{\mathcal{Q}}_k^\infty B^\vee$ be the preimage of $\mathcal{Q}_k^\infty B^\vee$ in $\tilde{\mathcal{Q}}_k B^\vee$. Then the family $\mathfrak{F}_{k,m}$ has a *pole* at $\tilde{\mathcal{Q}}_k^\infty B^\vee$ in the following sense: for a fixed $q_0 \in \tilde{\mathcal{Q}}_k^\infty B^\vee$ and any fixed scalar meromorphic function φ on $\tilde{\mathcal{Q}}_k B^\vee$ which vanishes on $\tilde{\mathcal{Q}}_k^\infty B^\vee$, and has no pole at q_0 , the expression $\varphi \mathfrak{F}_{k,m}$ gives a strictly meromorphic family which is strictly smooth in an appropriate neighborhood of q_0 . On the normal bundle $\mathcal{N}$ to $\tilde{\mathcal{Q}}_k^\infty B^\vee$ the family $\mathfrak{F}_{k,m}$ induces a homogeneous family of meromorphic functions of homogeneity degree -1 .

In particular, when studying homogeneous theory of $\mathfrak{MM}(Z)$, the line bundle $\mathcal{N}$ on $\tilde{\mathcal{Q}}_k^\infty B^\vee$ is an appropriate substitution of $\mathfrak{Q}_{k,m}$. It has an advantage of being smooth on a complete variety $\tilde{\mathcal{Q}}_k^\infty B^\vee$ (after a twist by $\mathcal{N}^\vee$).

In fact Proposition 3.2 may be partially extended to the other irreducible components of $\mathfrak{Q}_{k,m}^{\text{Hilb}}$, if one removes strictness conditions, and allows blowups of these components. This follows from the following

Lemma 3.8. *Consider a mapping $S \xrightarrow{\alpha} \text{Hilb}^m(Z)$ with an irreducible quasiprojective variety S . There is $M > 0$, a blow-up $\tilde{S} \xrightarrow{\pi} S$ and a mapping $\tilde{S} \xrightarrow{\alpha'} Z^{[M]}$ such that $\alpha(s) \subset \alpha'(\pi(s))$ if $s \in \tilde{S}$, here $\subset$ denotes inclusion of finite subschemes of Z .*

Proof. The technique of Section 3.2 will show that it is enough to prove the statement for an open subset $\tilde{S} \subset S$, instead of a blow-up. Associate to a finite subscheme $Z_0 \in \text{Hilb}^m(Z)$ the number $\text{rk}(Z_0)$ of distinct points of Z which support Z_0 (i.e., the length of $Z_{0,\text{red}}$). Let $\tilde{S}$ be the subset of S where $\text{rk}(\alpha(s))$ is maximal. Obviously, $\tilde{S}$ is open in S . Let $\mathfrak{m}_z$ be the maximal ideal of a point $z \in Z$, $\{z\}^k$ be the closed subscheme which correspond to the ideal $\mathfrak{m}_z^k$, $\text{mult}_z Z_0$ be the length of $Z_0 \cap \{z\}^m$. Define $\alpha'(s) = \bigcap_{z \in Z} \mathfrak{m}_z^{\text{mult}_z(Z_0)}$. Obviously, α' gives a smooth mapping of $\tilde{S}$ to $Z^{[M]}$ for an appropriate M . $\square$

3.2. Versality. As usual, assume that Z is a quasiprojective variety. Consider an admissible family of elements of $\mathfrak{MM}(Z)$ parameterized by S . By definition, there is a mapping $S \rightarrow \mathfrak{Q}_{k,m}$ (for appropriate k, m) such that the family is induced from the family $\mathfrak{F}_{k,m}$. For non-admissible families one gets the following statement:

Theorem 3.9. *For any meromorphic family $F_\bullet$ parameterized by an irreducible quasiprojective variety S there is a blow-up $\tilde{S}$ of S such that the pull-back of the family to $\tilde{S}$ is induced from a family $\mathfrak{F}_{k,m}$ on $\mathfrak{Q}_{k,m}$.*

If the family $F_\bullet$ is smooth, the same is true if we substitute $\mathfrak{Q}_{k,m}$ by $\mathring{\mathfrak{Q}}_{k,m}$.

Proof. Assume for the start that $F_\bullet$ is strictly meromorphic, and Z and S are projective. There is an open subset S_0 of S such that the family is admissible on S_0 , thus there is a mapping $\alpha: S_0 \rightarrow \mathfrak{Q}_{k,m}$ for appropriate k and m such that $F_\bullet|_{S_0}$ is induced from $\mathfrak{F}_{k,m}$.

Let $\alpha' = \alpha \times \text{id}: S_0 \rightarrow \mathfrak{Q}_{k,m} \times S_0$, S_1 be the closure of $\alpha'S_0$ in $\mathfrak{Q}_{k,m} \times S_0$. Then S_1 is a projective variety, thus is a blow-up $\tilde{S} = S_1 \rightarrow S$, and one can pull back the mapping $\alpha: S_0 \rightarrow \mathfrak{Q}_{k,m}$ from S_0 to $\tilde{S}$.

Lemma 3.10. *For any subvariety T of $\mathfrak{Q}_{k,m}$ the family $\mathfrak{F}_{k,m}$ induces a canonically defined strictly meromorphic family $\mathfrak{F}_{k,m}|_T$ on T . It is strictly smooth on $T \setminus \mathfrak{Q}_{k,m}^\infty$ and only there.*

Proof. The only thing one needs to prove is that when restricting $\mathfrak{F}_{k,m}$ to T , one will never need to resolve $0/0$. But we have seen that if $\mathcal{L}$ is m -ample, the restriction of $\mathfrak{F}_{k,m}$ to $X_{\mathcal{L}} \subset \mathcal{M}(Z)$ is well-defined, and has a pole only at $\mathfrak{Q}_{k,m}^\infty$, and zero only at $\mathfrak{Q}_{k,m}^0$. Since for any vector space V the projective subspaces $\mathcal{Q}_k^\infty V$ and $\mathcal{Q}_k^0 V$ do not intersect, this proves the lemma. $\square$

This gives two families of meromorphic functions on $\mathcal{M}(Z)$ parameterized by $\tilde{S}$, one induced from S , another one from $\mathfrak{Q}_{k,m}$. They coincide on the preimage of S_0 , thus everywhere. Minor modifications of this proof work in the cases of quasiprojective Z and S too.

Suppose that the family $F_\bullet$ is strictly smooth. Then the pull back to $\tilde{S}$ is also strictly smooth, thus one can see that $\alpha\tilde{S}$ does not intersect with $\mathfrak{Q}_{k,m}^\infty$. This proves the theorem in the case of strictly smooth $F_\bullet$.

Consider now the case of a smooth $F_\bullet$. There is an inclusion $S \rightarrow T$, and a blow-up $\pi: \tilde{T} \rightarrow T$ with $\tilde{T}$ which maps to $\mathfrak{Q}_{k,m}$. Again, taking completions, one may assume that S and T are projective. Let S_1 be the reduction of the scheme $\pi^{-1}S$. It is a variety that maps onto S . Consider any irreducible component S_2 of S_1 which maps onto S . Since S_2 is projective, cutting with hyperplanes, one can find a subvariety S_3 of S_2 such that the projection $S_3 \rightarrow S$ is finite of degree d over the generic point. Taking a finite covering of an open subset of S_3 , and taking a closure, one obtains a projective mapping $S_4 \rightarrow S$ which is finite Galois covering over an open subset of S , a mapping from S_4 to $\mathfrak{Q}_{k,m}$, and an action of a Galois group $G = \{g_1, \dots, g_d\}$ on S_4 .

We know that the pull-back of $F_\bullet$ to S_4 maps through $\mathfrak{F}_{k,m}$ via $S_4 \xrightarrow{\beta} \mathfrak{Q}_{k,m}$. For any $g \in G$ one can also consider a twisted map $\beta^g = \beta \circ g$. Consider

$$\bar{\beta} \stackrel{\text{def}}{=} \beta^{g_1} + \dots + \beta^{g_d}.$$

Let us explain what is the meaning of $+$ in the above expression. First of all, let γ be the composition of β with the projection $\mathfrak{Q}_{k,m} \rightarrow Z^{[m]}$. For $s \in S_4$ consider the finite subscheme $\bar{\gamma}s = \gamma^{g_1}s \cup \dots \cup \gamma^{g_d}s$ of length $m(s)$. Existence of a flattening subdivision [7] shows that for an open subset S_5 of S_4 the length of $\bar{\gamma}s$ does not depend on $s \in S_5$ (in fact by completeness of the Hilbert scheme it is $M = \max_{s \in S_4} m(s)$), and $\bar{\gamma}s$ depends smoothly on $s \in S_5$, i.e., $\bar{\gamma}$ extends to a regular mapping $S_5 \rightarrow \text{Hilb}^M Z$. Lemma 3.8 shows that decreasing S_5 and increasing M one can construct a mapping $S_5 \rightarrow Z^{[M]}$ instead.

Note that for any surjection of vector spaces $W \rightarrow V$ there is a natural mapping $\mathcal{Q}_k V \rightarrow \mathcal{Q}_k W$, thus each $\beta^{g_i} s$, $s \in S_5$, can be considered as an element of $\mathcal{Q}_k \mathfrak{D}_{\tilde{\gamma}s}^\vee$. This defines mappings $\tilde{\beta}^{g_i}: S_5 \rightarrow \mathfrak{Q}_{k,M}$. Since the images of these mappings do not intersect $\mathcal{Q}_k^\infty \mathfrak{D}_{\tilde{\gamma}s}^\vee$, the sum

$$\tilde{\beta}^{g_1} + \dots + \tilde{\beta}^{g_k}$$

is a well-defined element of $\mathcal{Q}_{dk} \mathfrak{D}_{\tilde{\gamma}s}^\vee$, thus defines a mapping $S_5 \rightarrow \mathfrak{Q}_{dk,M}$.

Taking a closure again, one obtains a mapping $\beta_6: S_6 \rightarrow \mathfrak{Q}_{dk,M}$ with S_6 being a projective variety with a mapping $S_6 \rightarrow S$ which is a finite Galois extension over an open subset of S . Moreover, the action of Galois group extends to S_6 and the mapping β_6 is Galois-invariant, thus induces a mapping $\beta_7: S_7 = S_6/G \rightarrow \mathfrak{Q}_{dk,M}$. Now S_7 is a blow-up of S , and the family on S_7 induced by β_7 coincides with d times the family induced by S . Dividing by d , we prove the theorem in the case of smooth $F_\bullet$ and $\text{char } \mathbb{K} = 0$.

If $\text{char } \mathbb{K} = p \neq 0$, suppose that degree of S_4/S in generic point is $d_1 p^r$ with $(d_1, p) = 1$. Instead of taking an arithmetic mean, split S_4/S into two extensions, S_4/S'_4 of degree d_1 , S'_4/S of degree p^r . For the extension of degree d_1 one can consider the arithmetic mean, as above, thus what remains is to consider the case of an extension of degree p^r .

In the case of such an extension consider the product of different β^{g_i} instead of the sum. Now the above arguments show that the family $F_\bullet^{p^r}$ is induced by β_7 .

Lemma 3.11. *The mapping $F \mapsto F^p$ induces a mapping $\mathfrak{Q}_{k,m} \xrightarrow{p} \mathfrak{Q}_{pk,m}$ with a closed image, and is a bijection with this image.*

Proof. Note that $\text{Supp } F_s^{p^r} = \text{Supp } F$ for any s , thus it is enough to show this in the case of the mapping $\mathcal{Q}_k V \rightarrow \mathcal{Q}_{pk} V$ for a finite-dimensional vector space V . In this case the lemma is obvious. $\square$

The lemma shows that F_s (for fixed any $s \in S_7$) may be written as a rational expression of appropriate $E_{z_i(s)}^\bullet$. Moreover, the p^r -th degree of coefficients c_l of these expressions are regular functions of $s \in S_7$. What remains is to show that the coefficients c_l themselves are regular functions of $s \in S_7$. On the other hand, fix sufficiently many test functions $\xi_i \in \mathcal{M}(Z)$, consider meromorphic functions $t_i = F_s(\xi_i)$ of $s \in S_7$. Each function t_i is a fraction of two linear combinations of c_l , coefficients of these linear combinations are meromorphic functions on S_7 . These provides a system of homogeneous linear equations on c_l with coefficients being meromorphic functions on S_7 . Obviously, it is possible to chose sufficiently many ξ_i so that this system determines c_l up to multiplication by a common meromorphic function on S_7 . Since for an appropriate value of this meromorphic function $c_l^{p^r}$ become regular, so are c_l . This finishes the proof for the smooth case.

Consider now the general case, when $F_\bullet$ is meromorphic. Substituting S by an appropriate blow-up, one may assume that $F_\bullet = \varphi \cdot G_\bullet$, with φ being a scalar meromorphic function on S , and $G_\bullet$ is induced by $\beta: S \rightarrow \mathfrak{Q}_{k,m}$. The mapping (φ, β)

maps on open subset of S to $\mathbb{P}^1 \times \mathfrak{Q}_{k,m}$, thus, replacing S by an appropriate blow-up, one can assume that $(\varphi, \beta) : S \rightarrow \mathbb{P}^1 \times \mathfrak{Q}_{k,m}$ is regular. On the other hand, for any vector space V there is a natural multiplication mapping

$$\mathbb{P}^1 \times \mathfrak{Q}_k V \setminus \delta \rightarrow \mathfrak{Q}_k V : ((\lambda_1, \lambda_2), (n, d)) \mapsto (\lambda_1 n, \lambda_2 d),$$

here $\delta = (\{0\} \times \mathfrak{Q}_k^\infty V) \cup (\{\infty\} \times \mathfrak{Q}_k^0 V)$. Since $\varphi \neq 0, \varphi \neq \infty$, one can see that there is a mapping from an open subset S_0 of S to $\mathfrak{Q}_{k,m}$. Now the same arguments with taking a closure and a blow-up finish the proof of the theorem. $\square$

Remark 3.12. Note that the values of k and m in the theorem are not uniquely determined. Moreover, for any $k_2 \geq k_1, m_2 > m_1$ there is a mapping from an appropriate blow-up of $\mathfrak{Q}_{k_1, m_1}$ to $\mathfrak{Q}_{k_2, m_2}$. Indeed, fix a point $z_0 \in Z$. For $Z_0 \in (Z \setminus \{z_0\})^{[m]}$ the scheme $Z_0 \cup \{z_0\}$ is finite of length $m+1$, and evaluation at z_0 gives a linear functional E_{z_0} on $\mathfrak{Q}_{Z_0 \cup \{z_0\}}$. Thus the mapping

$$(Z_0, n : d) \rightarrow (Z_0 \cup \{z_0\}, nE_{z_0}^l : dE_{z_0}^l)$$

maps an open subset of $\mathfrak{Q}_{k,m}$ to $\mathfrak{Q}_{k+l, m+1}$ if $l \geq 0$. This mapping extends to a mapping of an appropriate blowup of $\mathfrak{Q}_{k,m}$ to $\mathfrak{Q}_{k+l, m+1}$.

3.3. V -conformal operators. The description of families of elements of $\mathfrak{MM}(Z)$ from the previous sections allows a partial description of conformal operators. Recall that to each element of $\mathfrak{MM}(Z)$ one may associate the support, which is a finite subscheme of Z . Forgetting all the information about support except multiplicities of points in Z , one obtains a *support 0-cycle*, which is an element of $Z^m/\mathfrak{S}_m$ for an appropriate $m \in \mathbb{N}$.

Theorem 3.13. *Suppose that V is an affine open subset of a quasiprojective variety Z , and $\mathbf{C}$ is a V -conformal operator on $\mathfrak{MM}(Z)$ which dominates an open dense subset $U \subset Z$. Then there is a function $M : \mathbb{N}^2 \rightarrow \mathbb{N}$, blow-ups $Z^{(m,k)}$ of $Z^m/\mathfrak{S}_m$, and regular mappings*

$$\alpha_{m,k} : Z^{(m,k)} \rightarrow Z^{M(m,k)}/\mathfrak{S}_{M(m,k)}. \quad m, k \geq 1,$$

such that for any element F of $\mathfrak{MM}(Z)$ with the support 0-cycle $\sum a_i z_i$ there is $k > 0$ such that the support 0-cycle of $\mathbf{C}(F)$ is smaller than $\alpha_{m,k}(\sum a_i z_i)$ (here $m = \sum a_i$) if $\mathbf{C}(F)$ is defined.

Here $\alpha_{m,k}(\sum a_i z_i)$ means $\alpha_{m,k}(\eta)$, η being an arbitrary preimage of $\sum a_i z_i$ in $Z^{(m,k)}$.

Proof. Apply $\mathbf{C}$ to the family $\mathfrak{F}_{k,m}$ restricted to a big open subset $S \subset \mathfrak{Q}_{k,m}$. Let $T = U^{[m]}$, and $S = \pi^{-1}T$, here π is the projection of $\mathfrak{Q}_{k,m}$ to $Z^{[m]}$. Then $\mathbf{C}(\mathfrak{F}_{k,m}(s))$ is defined for any $s \in S$, and $\mathbf{C}(\mathfrak{F}_{k,m}(s))$ is a smooth family in $\mathfrak{MM}(V)$.

By Theorem 3.9, one can take a blow-up $\tilde{S}$ of a finite covering of S and a mapping $\alpha : \tilde{S} \rightarrow \mathfrak{Q}_{k', m'}$ which induces the family $\mathbf{C}(\mathfrak{F}_{k,m}(s))$. Compose α with the projection $\mathfrak{Q}_{k', m'} \rightarrow Z^{[m']}$ and the Chow mapping $Z^{[m']} \rightarrow Z^{m'}/\mathfrak{S}_{m'}$. By the above condition

and Definitions 1.4 and 1.10, one can pick up α (decreasing m' if necessary) in such a way that this composition β goes to an affine variety $V^{m'}/\mathfrak{S}_{m'}$.

Consider fibers of the projection from $\tilde{S}$ to S . They are projective varieties, thus each of them maps to a finite subset of $V^{m'}/\mathfrak{S}_{m'}$. Since a generic fiber of $\tilde{S}$ over S is a point, β induces a mapping from the normalization of S to $V^{m'}/\mathfrak{S}_{m'}$. Though fibers of $\pi: S \rightarrow U^{[m]}$ are not projective varieties, they are isomorphic to $\mathbb{P}^{2l-1} \setminus \mathbb{P}^{l-1}$, $l > 1$, thus contain many projective spaces. This means that any such fiber goes into a point when mapped to an arbitrary affine variety. Hence β induces a mapping from a normalization of $T = U^{[m]}$ to $V^{m'}/\mathfrak{S}_{m'}$. Finally, fibers of projection from $U^{[m]}$ to $U^m/\mathfrak{S}_m$ are projective, and a generic fiber is a point, thus β induces a mapping from normalization of $U^m/\mathfrak{S}_m$ to $V^{m'}/\mathfrak{S}_{m'}$. In other words, this is $\mathfrak{S}_n$ -invariant mapping from normalization of U^m to $V^{m'}/\mathfrak{S}_{m'}$, thus a mapping from $\tilde{U}^m/\mathfrak{S}_m$, $\tilde{U}$ being the normalization of U .

Moreover, $\tilde{S}$ and the mappings α and β can be completed in such a way that $\tilde{S}$ maps onto $\mathfrak{Q}_{k,m}$ and to $\mathfrak{Q}_{k',m'}$. This completion induces a mapping from a blow-up of $\tilde{Z}^m/\mathfrak{S}_m$ to $Z^{m'}/\mathfrak{S}_{m'}$, here $\tilde{Z}$ is the normalization of Z . On the other hand, $\tilde{Z}$ is a blow-up of Z , thus $\tilde{Z}^m/\mathfrak{S}_m$ is a blow-up of $Z^m/\mathfrak{S}_m$.

Since any element of $\mathfrak{MM}(Z)$ is in an appropriate $\mathfrak{Q}_{k,m}$, this finishes the proof. $\square$

Though the following statement does not add a lot of contents to Theorem 3.13, it will be very useful for study of codominant families. For two effective 0-cycles $\mathcal{A} = \sum a_i z_i$ and $\mathcal{A}' = \sum a'_j z_j$ let $\mathcal{A} \setminus \mathcal{A}'$ be the sum of terms $a_i z_i$ of $\mathcal{A}$ such that z_i does not appear in $\mathcal{A}'$.

Amplification 3.14. Suppose that V is an affine open subset of Z , and $\mathbf{C}$ is a V -conformal operator on $\mathfrak{MM}(Z)$ which dominates $U \subset Z$. Then there are functions $M, L: \mathbb{N}^2 \rightarrow \mathbb{N}$, blow-ups $Z^{(m,k)} \xrightarrow{\pi_{m,k}} Z^m/\mathfrak{S}_m$, and regular mappings

$$\alpha_{m,k}: Z^{m,k} \rightarrow Z^{M(m,k)}/\mathfrak{S}_{M(m,k)}. \quad m, k \geq 1,$$

such that for any element F of $\mathfrak{MM}(Z)$ with the support 0-cycle $\sum a_i z_i$ there is $k > 0$ such that the support 0-cycle of $\mathbf{C}(F)$ is smaller than

$$\alpha_{m,k} \left(\sum a_i z_i \right) + L(m, k) \cdot \left(\sum a_i z_i \right)$$

(here $m = \sum a_i$) if $\{z_i\} \subset U$. Here $\alpha_{m,k}(\sum a_i z_i)$ means $\alpha_{m,k}(\eta)$, η being an arbitrary preimage of $\sum a_i z_i$ in $Z^{(m,k)}$.

Moreover, let $U^{m,k}$ be the part of $Z^{(m,k)}$ which lies over $U^m/\mathfrak{S}_m$. Then for any $u \in U^{m,k}$

$$\alpha_{m,k}(u) \setminus \pi_{m,k}(u) \in V^L/\mathfrak{S}_L \quad \text{for a suitable } L \geq 0,$$

and for generic u supports of $\alpha_{m,k}(u)$ and $\pi_{m,k}(u)$ do not intersect.

Proof. For a cycle $\mathcal{A} = \sum a_i z_i$ denote $\sum a_i$ by $|\mathcal{A}|$. One may assume that Z is irreducible, since an appropriate blow-up of Z is. Obviously,

Lemma 3.15. *Suppose that Z is quasiprojective. Consider two regular mappings*

$$\alpha: T \rightarrow Z^M/\mathfrak{S}_M, \quad \alpha': T \rightarrow Z^N/\mathfrak{S}_N$$

with T being an irreducible variety. Then $|\alpha(t) \setminus \alpha'(t)|$ does not depend on $t \in U$ for an appropriate open dense subset $U \subset T$. Moreover,

$$t \mapsto \alpha(t) \setminus \alpha'(t)$$

gives a regular mapping $\beta: U \rightarrow Z^L/\mathfrak{S}_L$ for an appropriate L , thus for an appropriate blow-up $\tilde{T} \xrightarrow{\pi} T$ this mapping may be extended to $\beta: \tilde{T} \rightarrow Z^L/\mathfrak{S}_L$.

Moreover, for any $t \in \tilde{T}$ one has $\beta(t) \leq \alpha(\pi(t))$.

Consider α from Theorem 3.13, let α' be the projection of $Z^{(m,k)}$ to $Z^m/\mathfrak{S}_m$. Application of the previous lemma proves the first statement of the amplification. To prove the second statement, note that V -conformal operator sends an element of $\mathfrak{MM}(Z)$ with support at $A = \{z_1, \dots, z_l\} \subset Z$ to an element with support in $A \cup V$. This and Definition 1.4 show that one can assume that $\text{Supp } \alpha_{m,k}(u)$ is contained in $\text{Supp } \pi_{m,k}(u) \cup V$ (possibly after decreasing L). $\square$

3.4. Codominant families. While the above description of V -conformal operators gives a lot of information, the description of codominant families is yet more rigid:

Theorem 3.16. *Consider quasiprojective varieties Z and S , and a codominant conformal family $\mathbf{C}_s$ on S acting in $\mathfrak{MM}(Z)$. Then for any $l \in \mathbb{N}$ there are $M_l, N_l \in \mathbb{N}$ and a mapping $\alpha_l: S \rightarrow Z^{M_l}/\mathfrak{S}_{M_l}$ such that the mapping $\alpha_{m,k}$ from Theorem 3.13 can be taken to be*

$$\sum a_i z_i \mapsto N_l \sum a_i z_i + \alpha_k(s).$$

with α_k depending only on $s \in S$, but not on z_i .

Proof. Suppose that $\mathbf{C}_\bullet$ is defined on $U \subset S \times Z$. Define $U^{\{m\}} \subset S \times Z^m/\mathfrak{S}_m$ by the condition that the fiber of $U^{\{m\}}$ over any $s \in S$ is the m -th Chow power of the fiber of U over s :

$$U^{\{m\}} \cap (\{s\} \times Z^m/\mathfrak{S}_m) = \{s\} \times ((U \cap (\{s\} \times Z))^m / \mathfrak{S}_m).$$

Obviously, Amplification 3.14 can be generalized to families of conformal operators. This generalization provides a blow-up $\tilde{U} \xrightarrow{\pi} U^{\{m\}}$ and a mapping β from $\tilde{U}$ to $Z^M/\mathfrak{S}_M$ for an appropriate $M > 0$. For a generic $u \in \tilde{U}$ the supports of $\beta(u)$ and $\pi(u)$ do not intersect, for any u and any open $V \subset Z$ the support of $\beta(u) \setminus \pi(u)$ is in V as far as the S -projection of u is in an appropriate non-empty open dense subset $S_V \subset S$.

We want to show that $\beta(u)$ depends on S -projection of $u \in S \times Z^m/\mathfrak{S}_m$ only. To show this in the case $\dim Z \geq 2$, it is enough to prove

Proposition 3.17. *Suppose that $\dim Z \geq 2$. For any open dense $U \subset Z$ one can find an open dense $V \subset Z$ such that if a regular mapping b from any blow-up $\widehat{U} \xrightarrow{p} U^m/\mathfrak{S}_m$ to $Z^M/\mathfrak{S}_M$ satisfies*

$$\text{Supp}(b(u) \setminus p(u)) \subset V \quad \text{for any } u \in \widehat{U}$$

and

$$\text{Supp}(b(u)) \cap \text{Supp}(p(u)) = \emptyset \quad \text{for a generic } u \in \widehat{U},$$

then b is a constant mapping to $V^M/\mathfrak{S}_M$.

Proof. One may suppose that Z is irreducible and affine, $Z \subset \mathbb{A}^L$, $\dim Z = d$, $U = Z$. Note that for any fixed $Z_0 \in Z^{m-1}/\mathfrak{S}_{m-1}$ and any curve $\Gamma \subset Z$ the mapping $z \mapsto z + Z_0$, $z \in \Gamma$, sends Γ to $Z^m/\mathfrak{S}_m$. This mappings lifts to any blow-up of $Z^m/\mathfrak{S}_m$. Denote the corresponding curve in such a blowup $\Gamma \vee Z_0$.

Choose any $d-1$ coordinates $\eta_{i_1}, \dots, \eta_{i_{d-1}}$ on $\mathbb{A}^L$, consider the corresponding projection $Z \xrightarrow{H} \mathbb{A}^{d-1}$. Suppose that H is dominant. For an open subset $W \subset \mathbb{A}^{d-1}$ the preimages $\Gamma_w = H^{-1}(w)$, $w \in W$, are curves with the same number N of points “at infinity”, i.e., points in $\bar{\Gamma}_w \setminus \Gamma_w$, $\bar{\Gamma}_w$ being the (maximal) completion of Γ_w . Consider the mapping $\Gamma_w \rightarrow Z^M/\mathfrak{S}_M$ induced by the mapping $b|_{\Gamma_w \vee Z_0}$. Note that $\tilde{\Gamma}$ has no more than NM points at infinity. In particular, any regular mapping from $\tilde{\Gamma}$ to $\mathbb{A}^1 \setminus \sigma$ is constant if σ contains at least NM different points.

Suppose than no coordinate function η_j is constant on Z . Fix an arbitrary subset σ of $NM + m$ points on $\mathbb{A}^1$, let $V = Z \setminus \bigcup_j \eta_j^{-1}\sigma$. Then any mapping from $\tilde{\Gamma}_w$ to $V \cup Z'$ is constant if any η_j -projection of Z' contains at most m points. Consider the function $\varepsilon_k = \eta_{i_k} \circ b'$ restricted to $\tilde{\Gamma}_w$, $1 \leq k \leq d-1$. The first condition of the proposition shows that ε_k sends $\tilde{\Gamma}_w$ to $(\mathbb{A}^1 \setminus \sigma) \cup \varepsilon_k(\tilde{\Gamma}_w) \cup \varepsilon_k(Z_0)$. Since $\eta_{i_k}|_{\tilde{\Gamma}_w}$ is constant, $\varepsilon_k(\tilde{\Gamma}_w) \cup \varepsilon_k(Z_0)$ contains at most m points, thus ε_k is constant on $\tilde{\Gamma}_w$.

This shows that the composition of b with the projection of $Z^M/\mathfrak{S}_M$ to $(\mathbb{A}^{d-1})^M/\mathfrak{S}_M$ is constant on any $\Gamma_w \vee Z_0$ for any Z_0 and $w \in W$. Note that one can do the same with all $(d-1)$ -tuples of independent coordinates on Z (one may need to increase N to do it).

For a generic Z_0 and z the preimage $p^{-1}(z + Z_0)$ contains one point only. The above observation applied to all independent $(d-1)$ -tuples of coordinates shows that for any coordinate η_j on $\mathbb{A}^L$ the composition $\hat{\eta}_j \circ b \circ p^{-1}(z + Z_0)$ (here $\hat{\eta}_j$ is the extension of η_j -projection to a mapping of $Z^M/\mathfrak{S}_M$ to $(\mathbb{A}^1)^M/\mathfrak{S}_M$) depends only on $\eta_j(z)$ and Z_0 for generic z and Z_0 . Denote by ζ_j (or ζ_{j,Z_0}) the corresponding mapping

$$\mathbb{A}^1 \supset W_j \rightarrow (\mathbb{A}^1)^M/\mathfrak{S}_M: \eta_j(z) \mapsto \eta_j(b(z + Z_0))$$

defined on an appropriate open $W_j = W_{j,Z_0}$.

Let $\eta_j, \eta_{j'}$ be two independent coordinates on $Z \subset \mathbb{A}^L$ (this is where we use the condition that $\dim Z > 1$). We may assume that $\eta_j + \eta_{j'}$ coincides with another

coordinate $\eta_{j''}$ on $\mathbb{A}^L$ when restricted on Z . If we consider $\zeta_j, \zeta_{j'}, \zeta_{j''}$ as multivalued functions on open subsets of $\mathbb{A}^1$, the above remark shows that $\zeta_j(a) + \zeta_{j'}(b)$ is a (multivalued) function of $a + b$. As in the case of usual functions, this shows that all the branches of ζ_j and $\zeta_{j'}$ are parallel translations of each other:

Lemma 3.18. *Consider curves $\Gamma, \Gamma' \subset \mathbb{A}^2$. Denote coordinates on $\mathbb{A}^2 \times \mathbb{A}^2$ by x_1, x_2, y_1, y_2 . Consider the projection $\pi: \mathbb{A}^2 \times \mathbb{A}^2 \rightarrow \mathbb{A}^2$ given by $(x_1 + y_1, x_2 + y_2)$. If $\pi|_{\Gamma_1 \times \Gamma_2}$ is not dominant, there is an additive mapping $\rho: \mathbb{A}^1 \rightarrow \mathbb{A}^1$ such that Γ and Γ' are unions of graphs of $\rho + \text{const}$.*

Proof. All the tangent lines to Γ and Γ' have the same slope t . Doing a coordinate change $x'_2 = x_2 - tx_1$, one may assume that this slope is 0. Hence the equations of Γ can be written as $F(x_1^p, x_2) = 0$, $p = \text{char } \mathbb{K}$. If $p \neq 0$, and $F(s, t)$ depends on s , do a coordinate change $x'_1 = x_1^p$ and repeat the argument enough times. $\square$

In general there may be several components of the graphs Γ and Γ' of ζ_i and ζ_j , so we may need to apply the lemma to some components of $\Gamma \times \Gamma'$ only. Thus ζ_i and ζ_j may be unions of several parts, each of these parts being described by the lemma. Thus ζ_i, ζ_j may be unions of graphs of $\rho_t + \text{const}$, with the same collection of additive functions ρ_t for both ζ_i and ζ_j .

Let $k = 2$ if $\text{char } \mathbb{K} \neq 2$, otherwise $k = 3$. One may suppose that η_j^k coincides on Z with another coordinate $\eta_{j'''}$ on $\mathbb{A}^L$ when restricted on Z . Applying the lemma to $\eta_{j'''}$ and $\eta_{j'}$ shows that the additive functions ρ_t which correspond to the pair $(\eta_j, \eta_{j'})$ commute with taking k -th power. Thus for a fixed t either $\rho_t = 0$, or $\rho_t(X) = X^l$, here $l = p^a$, $a \geq 0$, if $\text{char } \mathbb{K} = p \neq 0$, and $l = 1$ if $\text{char } \mathbb{K} = 0$.

If $\text{char } \mathbb{K} \neq 0$, and $\mathbb{K}$ contains a transcendental λ , then, as above, one can conclude that ρ_t commute with multiplication by λ . Thus $a = 0$ in the above case, or otherwise $\mathbb{K}$ is the algebraic closure of $\mathbb{F}_p$, and ρ is a power of Frobenius morphism Φ .

If $\rho = 0$, then the corresponding part of ζ is a constant mapping $\mathbb{A}^1 \rightarrow (\mathbb{A}^1)^{M_1} / \mathfrak{S}_{M_1}$. If $\rho = \text{id}$, then the corresponding part of ζ is constant, thus $bp^{-1}(z + Z_0)$ contains a multiple of z , what contradicts the second condition of the proposition, since Z_0 is arbitrary. One can see that in general $bp^{-1}(z + Z_0)$ (for a fixed Z_0) is some constant term plus $\sum_{a=0}^A c_a \Phi^a(z)$.

The constant term in this expression may depend on Z_0 , breaking Z_0 into a sum $z' + Z_1$ and repeating above arguments one can see that b is

$$\sum_{a=0}^A c_a \Phi^a p + C, \quad c_a \geq 0, \quad C \in Z^{M'} / \mathfrak{S}_{M'}, \quad \sum_a c_a + M' = M.$$

Use $\dim Z \geq 2$ again. Taking a projection on $\mathbb{A}^2$, it is enough to consider the case $Z \subset \mathbb{A}^2$. One may suppose that $Z \cap (\mathbb{A}^1 \times \{0\}) \neq \emptyset$. Let $V = Z \setminus (\mathbb{A}^1 \times \{0\})$. Obviously, if the above b satisfies the conditions of proposition with this V , then $c_a = 0$ for any a . This finishes the proof of the proposition. $\square$

Consider the case $\dim Z = 1$. Decreasing U , one may assume that Z is projective and normal. Fix Z_0 again, restrict β to elements of the form $(s, z + Z_0)$, $z \in Z$. This defines a mapping β_{Z_0} from an open subset of $S \times Z$ to $Z^M/\mathfrak{S}_M$, replacing S by an open subset, one may extend it to a mapping of $S \times Z$. Note that to a mapping $Z \xrightarrow{\gamma} Z^M/\mathfrak{S}_M$ one may associate a curve $\Gamma \subset Z \times Z$ given by the condition that $\gamma(z_0)$ is the intersection of $\{z_0\} \times Z$ with Γ (it is possible that one may need to assign multiplicities to components of Γ). This gives a family Γ_s of curves in $Z \times Z$ parameterized by S . All these curves have the same degree of projection to the second component, call it L . To show that β depends on s only, it is enough to show that $L = 0$.

Suppose that $L > 0$. Taking a finite mapping of Z to $\mathbb{P}^1$, and multiplying M by degree, one can see that it is enough to consider $Z = \mathbb{P}^1$. Fix a generic $s_0 \in S$, consider $\Gamma = \Gamma_{s_0} \subset \mathbb{P}^1 \times \mathbb{P}^1$. We know that Γ does not contain the diagonal as an irreducible component. Consider two projections $\pi_{1,2}$ of Γ to $\mathbb{P}^1$ of degrees M and L as meromorphic functions on Γ . Then $\pi_1 - \pi_2$ is a correctly defined meromorphic function on Γ which is not zero on any irreducible component of Γ , thus has no more than $L + M$ zeros. Suppose $Z \setminus U$ contains d point, let V_0 be $\mathbb{P}^1$ without any collection of N points, $N > L + M + Md$. Obviously, $\Gamma \cap U \times \mathbb{P}^1$ cannot be contained in $(\mathbb{P}^1 \times V_0) \cup \Delta$, Δ being a diagonal in $\mathbb{P}^1 \times \mathbb{P}^1$.

This shows that a family for which $L > 0$ cannot be V_0 -conformal on any open subset of S , which proves the theorem. $\square$

Definition 3.19. Say that the family $\mathbf{C}_s$ is *clean* if Theorem 3.16 remains true with one mapping $\alpha: S \rightarrow Z^M/\mathfrak{S}_M$ instead of a sequence of mappings $\alpha_l: S \rightarrow Z^{M_l}/\mathfrak{S}_{M_l}$.

Remark 3.20. One should expect that an appropriate modification of definition of a conformal field will make all codominant conformal fields to be clean. With the definitions used here, any infinite sum $\mathbf{C}^{(1)} + \mathbf{C}^{(2)} + \dots$ of conformal families on S acting on $\mathfrak{MM}(Z)$ is a conformal family again, as far as all the summands but a finite number vanish on $X_{\mathcal{L}}$ for any given line bundle $\mathcal{L}$ on M . It is the acceptability of such infinite sums which makes it easy to construct examples of non-clean codominant conformal families.

Example 3.21. Consider the family E_z , $z \in Z$. Then one can consider $\alpha(z) = z$, thus this is a clean family.

A composition $\mathbf{C}_s \mathbf{C}'_t$ of two clean families is clean again. A subfamily of a clean family is clean, similarly, all the residues of a clean family are clean.

Remark 3.22. Note that, inversely, any conformal family may be written as a sum of clean families satisfying conditions of Remark 3.20.

Remark 3.23. Say that a conformal operator $\mathfrak{MM}(U) \xrightarrow{\mathbf{c}} \mathfrak{MM}(Z)$ is *geometric* if it is a homomorphism of algebras, say that it is *semigeometric* if there is a homomorphism of algebras $\mathfrak{MM}(U) \xrightarrow{\mathbf{c}'} \mathfrak{MM}(Z)$ such that $\mathbf{c}$ is a corresponding homomorphism of

modules, i.e., $\mathbf{c}(ab) = \mathbf{c}'(a) \mathbf{c}(b)$. Since $\mathfrak{M}\mathcal{M}(Z)$ is considered as a field of functions on $\mathcal{M}(Z)$, one should consider a geometric operator as a rational mapping $\mathcal{M}(Z) \xrightarrow{f} \mathcal{M}(Z)$, and a semigeometric operator as a rational fiber-wise mapping of line bundles over $\mathcal{M}(Z)$.

To describe a geometric mapping $\mathbf{c}$ it is enough to describe the family $\mathbf{c}E_z$, $z \in Z$, of elements of $\mathfrak{M}\mathcal{M}(Z)$. In particular, to any such mapping one may associate a mapping $Z \rightarrow Z^m/\mathfrak{S}_m$ for an appropriate $m > 0$, or, what is the same, a mapping β from a finite extension $\tilde{Z}$ of Z to Z . As before, one can show that a geometric mapping may be included into a codominant family only if β is constant on $\tilde{Z} \setminus \bar{Z}$, and either $\bar{Z} \subset \tilde{Z}$ is empty, or is an irreducible component of $\tilde{Z}$ which projects isomorphically on Z , and $\beta|_{\bar{Z}}$ coincides with this projection. Note that in this case the element α of $Z^M/\mathfrak{S}_M$ from Theorem 3.16 may be taken to be $\beta(\tilde{Z} \setminus \bar{Z})$.

Note that a semigeometric mapping $\mathbf{c}$ uniquely determines the corresponding geometric mapping $\mathbf{c}'$, since $\mathbf{c}'(a) = \mathbf{c}(a)/\mathbf{c}(1)$. One can write $\mathbf{c}(1) = \Phi(E_{z_1}, \dots, E_{z_m})$ with an appropriate rational function Φ (possibly including some derivatives of E_{z_k} as well), thus $\mathbf{c} = \Phi(\mathbf{E}_{z_1}, \dots, \mathbf{E}_{z_m}) \mathbf{c}'$. The mapping $\mathbf{c}$ may be included into a codominant family iff $\mathbf{c}'$ may. The corresponding element α of $Z^M/\mathfrak{S}_M$ is the union of the corresponding element for $\mathbf{c}'$ and of $\{z_1, \dots, z_m\}$. One may see that smooth families of semigeometric mappings are clean, and Theorem 3.16 gives estimates for the positions of z_i .

Note that all the examples of conformal fields considered in this paper are semigeometric.

3.5. Conformal algebras. Consider conformal fields $\mathbf{C}(s)$ and $\mathbf{C}'(t)$, $s \in S$, $t \in T$, acting on $\mathfrak{M}\mathcal{M}(Z)$. Suppose that $\mathbf{C}$ is defined on $U \subset S \times Z$, $\mathbf{C}'$ is a clean codominant family with the associated mapping $\alpha: T \rightarrow Z^m/\mathfrak{S}_m$. Define the *exceptional subset* $\sigma_{\mathbf{C}\mathbf{C}'} \subset S \times T$ by

$$(s, t) \in \sigma_{\mathbf{C}\mathbf{C}'} \text{ if } \alpha(t) \cap R_s \neq \emptyset,$$

here $R_{s_0} \subset Z$ is the subset of Z where $\mathbf{C}(s_0)$ is not defined, i.e., $(S \times Z \setminus U) \cap (\{s_0\} \times Z)$.

The following obvious theorem shows that a product of several clean codominant conformal fields has singularity uniquely determined by singularities of pairwise products. In particular, triple product has no fusions which do not correspond to fusions of pairwise products. This sets a foundation stone for the theory of conformal algebras, since it allows one to investigate relations on fusions implied by associativity of multiplication.

Theorem 3.24. *Consider smooth clean codominant families $\mathbf{C}_1(s_1)$, $\mathbf{C}_2(s_2)$, $\dots$, $\mathbf{C}_k(s_k)$, $s_i \in S_i$, $i = 1, \dots, k$, acting on $\mathfrak{M}\mathcal{M}(Z)$. Then the product family*

$$\mathbf{C}_1(s_1) \mathbf{C}_2(s_2) \dots \mathbf{C}_k(s_k)$$

is smooth at $(s_1, s_2, \dots, s_k)$ if $(s_i, s_j) \notin \sigma_{\mathbf{C}_i \mathbf{C}_j}$ for any $1 \leq i < j \leq k$.

Remark 3.25. Note that by complicating notations by having a sequence of subschemes $\sigma_{\mathbf{C}_i \mathbf{C}_j}^{(l)}$ instead of one subscheme $\sigma_{\mathbf{C}_i \mathbf{C}_j}$ one can remove the requirement that the families are clean. Similarly, one may remove smoothness requirement by introducing subschemes $\sigma_{\mathbf{C}_i} \subset S_i$ where $\mathbf{C}_i$ is not smooth, and stating that

$$\mathbf{C}_1(s_1) \mathbf{C}_2(s_2) \dots \mathbf{C}_k(s_k)$$

is smooth at $(s_1, s_2, \dots, s_k)$ if $s_i \notin \sigma_{\mathbf{C}_i}$ for any $1 \leq i \leq k$, and $(s_i, s_j) \notin \sigma_{\mathbf{C}_i \mathbf{C}_j}$ for any $1 \leq i < j \leq k$.

Thus the only important requirement in the theorem 3.24 is the codominantness. Recall that the codominantness ensures that the family may be composed on the left with an arbitrary conformal operator.

Remark 3.26. Yet another reference to classification of meromorphic families in $\mathfrak{MM}(Z)$ shows that for any $k, m \geq 0$ there is a number $L = L(k, m)$ such that for a “big enough” family $F_t, t \in T$, the family

$$\Phi(s_1, s_2, \dots, s_k) \mathbf{C}_1(s_1) \mathbf{C}_2(s_2) \dots \mathbf{C}_k(s_k) F_t$$

is smooth in $(s_1, s_2, \dots, s_k)$ (in the sense that it is smooth on a subset of $S_1 \times \dots \times S_k \times T$ which projects onto $S_1 \times \dots \times S_k$) as far as Φ has a zero of order L on the subsets given by $s_i \in \sigma_{\mathbf{C}_i}$ for any $1 \leq i \leq k$, and by $(s_i, s_j) \in \sigma_{\mathbf{C}_i \mathbf{C}_j}$ for any $1 \leq i < j \leq k$.

Remark 3.27. Theorem 3.24 shows that in the settings of this paper the conformal associativity relations are corollaries of geometry of punctual Hilbert schemes. Note the similarity with [8], where the associativity relations are obtained as corollaries of geometry of moduli spaces.

The same arguments show

Proposition 3.28. *Consider clean codominant families $\mathbf{C}(s)$ and $\mathbf{C}'(t)$, $s \in S$, $t \in T$, let $\mathbf{D}$ be a fusion of $\mathbf{C}$ and $\mathbf{C}'$ on an irreducible subvariety $P \subset S \times T$. If P projects onto S and onto T , then*

- (1) $\mathbf{D}$ is clean codominant;
- (2) if $\mathbf{D}'(r)$, $r \in R$, is another clean codominant family, then

$$\sigma_{\mathbf{D} \mathbf{D}'} \subset ((\sigma_{\mathbf{C} \mathbf{D}'} \times T) \cup (S \times \sigma_{\mathbf{C}' \mathbf{D}'})) \cap (P \times R) \subset S \times T \times R.$$

3.6. Fractions of formal series. Consider a meromorphic function F on Z which has no pole at $z_0 \in Z$. Then one can consider F as a regular function on a formal completion $\hat{Z}$ of Z at z_0 , i.e., as a formal function near z_0 . Choosing a coordinate system $(\eta_1, \dots, \eta_d)$ near a smooth point z_0 makes F into a formal series in $\eta_1, \dots, \eta_d$.

If one lifts the restriction that F has no pole at z_0 , then some extension of the algebra $\mathbb{K}[[\eta_1, \dots, \eta_d]]$ of formal series in $\eta_1, \dots, \eta_d$ should be introduced. A natural choice is the field of fractions of $\mathbb{K}[[\eta_1, \dots, \eta_d]]$, but it is a pain to work with if $d > 1$, since $\mathbb{K}[[\eta_1, \dots, \eta_d]]$ has too many minimal prime ideals. One way to construct a

field which contains formal functions *and* meromorphic functions is to consider an iterated construction of residue from Section 1.9.

Consider a *flag* $Z = Z_d \supset \cdots \supset Z_0 = \{z_0\}$ of irreducible subvarieties of Z (or ∞ -jets of subvarieties of Z near z_0 , i.e., subvarieties of $\widehat{Z}$). A *normalized flag* is a flag with a lifting $\iota_k: \widetilde{Z}_k \hookrightarrow \widetilde{Z}_{k+1}$ of the inclusion $Z_k \hookrightarrow Z_{k+1}$ into the normalizations $\widetilde{Z}_k, \widetilde{Z}_{k+1}$ of Z_k and Z_{k+1} , for every $0 \leq k < d$. Fix an *equation* η_{k+1} of Z_k in Z_{k+1} , i.e., a meromorphic function on Z_{k+1} which has a zero of the first order at $\iota_k \widetilde{Z}_k \subset \widetilde{Z}_{k+1}$. We will also fix meromorphic vector fields v_{k+1} on $\widetilde{Z}_{k+1}$ such that $\langle v_{k+1}, d\eta_{k+1} \rangle = 0$, and the restriction of v_{k+1} to $\iota_k \widetilde{Z}_k$ is defined and not tangent to $\iota_k \widetilde{Z}_k$. Such data allows one taking residues of meromorphic functions on Z_{k+1} along Z_k (compare Section 1.9). Note that if all Z_k are normal, $(\eta_1, \dots, \eta_d)$ may be extended to become a *regular sequence* of functions on an open subset of Z (see [2]), in smooth case one may assume instead that $\eta_\bullet$ gives a coordinate system near z_0 .

Any meromorphic function F on Z (or an element of the field of fractions of $\mathbb{K}[[\eta_1, \dots, \eta_d]]$) is uniquely determined by its residues on Z_{d-1} of all orders, which are in turn meromorphic functions on Z_{d-1} (or elements of the corresponding field of fractions). Now one can take residues of these meromorphic functions on Z_{d-2} , and so on.

As a result, one associates to F a *formal Laurent series*¹⁹

$$(3.1) \quad \sum_{k_1, \dots, k_d} \phi_{k_1, \dots, k_d} \eta_1^{k_1} \cdots \eta_d^{k_d},$$

and the coefficients $\phi_{k_1, \dots, k_d}$ vanish if

$$k_1 < -K_1, k_2 < -K_2(k_1), \dots, k_d < -K_d(k_1, \dots, k_{d-1}).$$

Note that the series as the above one may be added, multiplied, *and* divided. Denote the field of such formal series by $\mathbb{K}((\eta_d))((\eta_{d-1})) \cdots ((\eta_1))$. We consider elements of this field as functions on the *formal completion* $\widehat{Z}'$ of Y at the *normalized flag* $Z = Z_d \supset \cdots \supset Z_0 = \{z_0\}$.

If a different sequence $(\eta'_1, \dots, \eta'_d)$ results in the same flag, then one can easily establish a canonical isomorphism of $\mathbb{K}((\eta_d))((\eta_{d-1})) \cdots ((\eta_1))$ and $\mathbb{K}((\eta'_d))((\eta'_{d-1})) \cdots ((\eta'_1))$. However, completions at different normalized flags are not related, as the following example shows:

Consider a transposition $\sigma \in \mathfrak{S}_d$, suppose that η_i and v_i are extended to Z , and $(\eta_{\sigma_1}, \dots, \eta_{\sigma_d})$ is a regular sequence too. Then it defines a different normalized flag, and a different completion $\widehat{Z}''$ of Z . A question arises whether one is able to compare functions on $\widehat{Z}'$ and $\widehat{Z}''$ for equality.

Suppose that a function F in $\mathbb{K}((\eta_d))((\eta_{d-1})) \cdots ((\eta_1))$ becomes an element F_1 in $\mathbb{K}[[\eta_1, \dots, \eta_d]]$ after multiplication by an element $F_2 \in \mathbb{K}[[\eta_1, \dots, \eta_d]]$, i.e., $F =$

¹⁹Note that though η_i are defined on subvarieties Z_i only, one can extend them to (at least) the formal neighborhood of Z_i in Z using vector fields v_i .

F_1/F_2 . Obviously, F can be represented as an element of $\mathbb{K}((\eta_{\sigma_d}))((\eta_{\sigma_{d-1}})) \dots ((\eta_{\sigma_1}))$ as well.

Example 3.29. Consider $F = \sum_{k \geq 0} \eta_1^k \eta_2^{-k} \in \mathbb{K}((\eta_2))((\eta_1))$. Then $F = \frac{\eta_2}{\eta_2 - \eta_1}$, thus can be written as $F = -\sum_{k \geq 1} \eta_1^{-k} \eta_2^k \in \mathbb{K}((\eta_1))((\eta_2))$.

Similarly, one can define a relation of *being an analytic continuation* of each other for completions of Z at arbitrary pair of normalized flags, but as with usual analytic continuation, it can be made 1-to-1 on very small subsets only.

Remark 3.30. Consider how the Laurent coefficient at a normalized flag are related to the “usual” Laurent coefficients of a function of two complex variables. Given an analytic function $\xi(z_1, z_2)$ defined on an open subset U of $\mathbb{C}^2$ one may define Laurent coefficients as integrals

$$\xi_{kl} = \frac{1}{(2\pi i)^2} \iint_{\substack{|z_1|=\varepsilon_1 \\ |z_2|=\varepsilon_2}} \frac{\xi(z_1, z_2)}{z_1^{k_1+1} z_2^{k_2+1}} dz_1 dz_2,$$

here $\varepsilon_1, \varepsilon_2$ are appropriate numbers such that the torus $\mathbb{T}_{\varepsilon_1 \varepsilon_2}^2 = \{|z_1| = \varepsilon_1, |z_2| = \varepsilon_2\}$ is in U . The coefficients do not change if one deforms $\varepsilon_1, \varepsilon_2$ in such a way that $\mathbb{T}_{\varepsilon_1 \varepsilon_2}^2$ remains in U during the deformation.

Consider the flag defined by the sequence (z_1, z_2) . Laurent coefficients near this flag correspond to considering the residues at $z_1 = 0$, and residues of these residues at $z_2 = 0$. In other words, one takes the limit $\varepsilon_1 \rightarrow 0$, and then the limit $\varepsilon_2 \rightarrow 0$. On the other hand, if F is meromorphic, then the possibility of description of roots of denominator by Puiseux series shows that there exist $\varepsilon > 0$ and $K \geq 0$ such that $F(z_1, z_2)$ is defined on

$$0 < |z_1| < |z_2|^K, 0 < |z_2| < \varepsilon.$$

Obviously, the above double limit coincides with the “usual” Laurent coefficients for any $0 < \varepsilon_1 < \varepsilon_2^K, 0 < \varepsilon_2 < \varepsilon$.

Similarly, the flag given by the sequence (z_2, z_1) corresponds to taking $0 < \varepsilon_2 < \varepsilon_1^K, 0 < \varepsilon_1 < \varepsilon$. When one deforms the contour of integration from one region to another, it may hit the divisor of poles of F , each time adding residues at the corresponding irreducible component of this divisor.

3.7. Laurent coefficients of conformal fields. Similarly to what is done in the previous section, given a meromorphic function F on $S \times X$, and a normalized flag $\{S_i\}$ on S equipped with equations η_i and vector fields v_i , one can consider F as an element of $\mathcal{M}(X)((\eta_d))((\eta_{d-1})) \dots ((\eta_1))$. Moreover, if F_s is a meromorphic function on $\mathcal{M}(Z)$ parameterized by $s \in S$, one can consider F as an element of $\mathfrak{MM}(Z)((\eta_d))((\eta_{d-1})) \dots ((\eta_1))$.

Apply the defined above completion to a conformal field $\mathbf{C}$ on S acting in $\mathfrak{MM}(Z)$. Suppose that $\mathbf{C}$ dominates $U \subset Z$. Given a function $F \in \mathfrak{MM}(U)$, $\mathbf{C}(s)F$ is a meromorphic function on $S \times \mathcal{M}(Z)$, thus one can consider it as a function on

$\widehat{S} \times \mathcal{M}(Z)$, $\widehat{S}$ being the formal completion of S at the normalized flag $\{S_i\}$. Consider Laurent coefficients $\phi_{k_1 \dots k_D} \in \mathfrak{M}\mathcal{M}(Z)$ of $\mathbf{C}(s)F$, $D = \dim S$. Each of these coefficient is obtained by a sequence of taking a fine residue (defined in Section 1.10) at subvarieties of the normalized flag $\{S_i\}$ (next subvariety is a hypersurface in the previous one), thus there are operators $\mathbf{c}_{k_1 \dots k_D}$ such that

$$\phi_{k_1 \dots k_D} = \mathbf{c}_{k_1 \dots k_D} F$$

and $\mathbf{c}_{k_1 \dots k_D}$ gives a linear mapping $\mathfrak{M}\mathcal{M}(U) \rightarrow \mathfrak{M}\mathcal{M}(Z)$ (which is not necessarily a conformal operator, since it may fail to send smooth families to smooth families). Call these operators *Laurent coefficients* of $\mathbf{C}$ at the normalized flag $\{S_i\}$. (Strictly speaking, one should associate Laurent coefficients to the sequence of equations η_i and vector fields v_i , but as it was noted above, different systems of equations lead to isomorphic fields of functions).

For any family F_t , $t \in T$, of functions F the functions $\mathbf{c}_{k_1 \dots k_D} F_t$ vanish for sufficiently big negative $(k_1, \dots, k_D)$ (big in the same sense as in (3.1)). As we have seen in Remark 1.31, the operators $\mathbf{c}_{k_1 \dots k_D}$ do not necessarily vanish for big negative $(k_1, \dots, k_D)$.

Proposition 3.31. *Consider a clean codominant conformal field $\mathbf{C}(s)$ on S acting on $\mathfrak{M}\mathcal{M}(Z)$ and a flag $S_\bullet$ on S . Suppose that $\mathbf{C}$ dominates $U \subset Z$. Then there is a finite subset $Z_0 \subset Z$ such that for any multiindex $(k_1, \dots, k_D)$ operator $\mathbf{c}_{k_1 \dots k_D}$ sends $\mathfrak{M}\mathcal{M}(U_1) \rightarrow \mathfrak{M}\mathcal{M}(U_2)$ if $U_1 \subset U$, $U_2 \supset U_1 \cup Z_0$.*

Proof. This is an immediate corollary of Theorem 3.16 and the following obvious

Lemma 3.32. *Consider a family F_s in $\mathfrak{M}\mathcal{M}(Z)$ parameterized by S . Suppose that this family corresponds to a mapping $\beta: S \rightarrow \mathfrak{Q}_{k,m}$. Let π be the projection $\mathfrak{Q}_{k,m} \rightarrow Z^{[m]}$. Then for any flag $S_0 \subset \dots \subset S_d$ in S the Laurent coefficients $F_{k_1 \dots k_d}$ of F at $S_\bullet$ are in $\mathfrak{M}\mathcal{M}(U)$ if $U \supset \pi(\beta(S_0))$.*

□

3.8. Laurent coefficients of a composition. Consider two conformal fields, $\mathbf{C}(s)$ on S , and $\mathbf{C}'(t)$ on T , both acting on $\mathfrak{M}\mathcal{M}(Z)$. Suppose that $\mathbf{C}'$ is codominant, thus the composition $\mathbf{C}(s)\mathbf{C}'(t)$ is defined. Let $\mathcal{S} = \{S_i\}$, $i = 0, \dots, D$, be a normalized flag on S , $\mathcal{T} = \{T_j\}$, $j = 0, \dots, D'$, be a normalized flag on T , define a normalized flag $\mathcal{S} * \mathcal{T}$ on $S \times T$ to be

$$S_D \times T_{D'} \supset \dots \supset S_D \times T_1 \supset S_D \times T_0 \supset \dots \supset S_0 \times T_0.$$

Note that similarly defined $\mathcal{T} * \mathcal{S}$ can be considered as a flag on $S \times T \simeq T \times S$ too, and that additional data η_i, v_i for $\mathcal{S}$ and $\mathcal{T}$ gives similar data for $\mathcal{S} * \mathcal{T}$ and $\mathcal{T} * \mathcal{S}$.

Theorem 3.33. *Consider Laurent coefficients $\mathbf{c}_\bullet, \mathbf{c}'_\bullet$ of $\mathbf{C}$ and $\mathbf{C}'$ at $\mathcal{S}$ and $\mathcal{T}$. Suppose that $\mathbf{C}'$ is codominant, let Z_0 is the corresponding finite subset of Z , as in Proposition 3.31. Suppose that $\mathbf{C}$ dominates $U \supset Z_0$, $\mathbf{C}'$ dominates U' , so the*

composition $\mathbf{c}_K \circ \mathbf{c}'_{K'}$ is well-defined on $\mathfrak{MM}(U \cap U')$. Then $\mathbf{c}_K \circ \mathbf{c}'_{K'}$ restricted to $\mathfrak{MM}(U \cap U')$ coincides with the (K, K') -th Laurent coefficient of $\mathbf{C} \circ \mathbf{C}'$ at $\mathcal{S} * \mathcal{T}$.

Proof. Fix a function $F \in \mathfrak{MM}(U \cap U')$, then $\mathbf{C}'(t)F$ and $\mathbf{C}(s)(\mathbf{C}'(t)F)$ are well-defined meromorphic families. Apply the classification of meromorphic families of Theorem 3.9 to $\mathbf{C}'(t)F$. This leads to a blow-up T' of T .

Lemma 3.34. *Consider a normalized flag $T_\bullet$ in T , and a blow-up $T' \rightarrow T$. Then $T_\bullet$ may be naturally lifted to a normalized flag $T'_\bullet$ in T' , each T'_k being a blow-up of T_k . Given equations $\eta_\bullet$ of $T_\bullet$ and vector fields $v_\bullet$ (as in Section 3.6), there is a natural lifting of these data to T' .*

Proof. Let $d = \dim T$. It is enough to consider T_{d-1} , and then proceed by induction. The normalization $\tilde{T}'$ of T' is a blow-up of the normalization $\tilde{T}$ of T in a closed subscheme $Z_0 \subset \tilde{T}$ of codimension greater than 2. In particular, Z_0 does not contain $\iota_{d-1}\tilde{T}_{d-1}$, thus there is a canonically defined blowup $\tilde{T}'_{d-1}$ of $\tilde{T}_{d-1}$ and a mapping $\tilde{T}'_{d-1} \rightarrow \tilde{T}' = \tilde{T}'_d$. Obviously, $\tilde{T}'_{d-1}$ is a normalization of its image in T' .

The vector field v_d and the meromorphic function η_d on $\tilde{T} = \tilde{T}_d$ may be lifted to $\tilde{T}'$ in a canonical way. What we need to prove is that the lifting of η_d has a zero at $\tilde{T}'_{d-1}$, and the lifting of v_d is defined at $\tilde{T}'_{d-1}$, and is not tangent to $\tilde{T}'_{d-1}$. Both parts follow from $\tilde{T}'_{d-1}$ being isomorphic to $\tilde{T}_{d-1}$ at generic point. $\square$

Call such a lifting a *strict preimage* of the flag $T_\bullet$. Note that one can also define a strict preimage for *incomplete* normalized flags $T_k \subset T_{k+1} \subset \dots \subset T_d = T$ as far as each subvariety is a hypersurface in the next subvariety.

Obviously, for a given meromorphic family F_t in $\mathfrak{MM}(Z)$, $t \in T$, and a normalized flag $T_\bullet$ on T the Laurent coefficients of F_t at $T_\bullet$ do not change after any blowup of T (with the corresponding lifting of η_i and v_i). Moreover, multiplication of $\mathbf{C}(s)$ and $\mathbf{C}'(t)$ by scalar meromorphic functions of s and t does not change the validity of the theorem. In particular, one may assume that both $\mathbf{C}$ and $\mathbf{C}'$ are smooth, and in conditions of Theorem 3.9 the family $F_t = \mathbf{C}'(t)F$ leads to no blow-up of T .

This eliminates the field $\mathbf{C}'(t)$, instead one may consider Laurent coefficients of $\mathbf{C}(s)F_t$ for an arbitrary smooth family F_t in $\mathfrak{MM}(U)$.

Note that for an incomplete normalized flag $R_k \subset R_{k+1} \subset \dots \subset R_d = R$ with each subvariety being a hypersurface in the next subvariety (with associated data η_i and v_i), and for a meromorphic function on R one can define *partial Laurent coefficients* $F_{l_k \dots l_d}$ of F which are meromorphic functions on R_k .

Proposition 3.35. *Suppose that a conformal field $\mathbf{C}(s)$ on S acting in $\mathfrak{MM}(Z)$ dominates $U \subset S$, and F_t , $t \in T$, is a smooth family in $\mathfrak{MM}(U)$. Fix a normalized flag $\mathcal{T}$ in T . Then $(l_1, \dots, l_{D'})$ -partial Laurent coefficient of $\mathbf{C}(s)F_t$ on the partial flag $S \times \mathcal{T} \stackrel{\text{def}}{=} \{S \times T_{k-D}\}_{k=D, \dots, D+D'}$ coincides with $\mathbf{C}(s)F_{l_1 \dots l_{D'}}$, $F_{l_1 \dots l_{D'}}$ being $(l_1, \dots, l_{D'})$ -Laurent coefficient of F at the flag $\mathcal{T}$.*

Proof. Again, $\mathbf{C}(s)F_t$ is a family on $R = S \times T$, thus Theorem 3.9 leads to a blow-up R' of R . There is a mapping $\beta: R' \rightarrow \mathfrak{Q}_{k,m}$ which induces the family $\mathbf{C}(s)F_t$.

Suppose that R' is the blow-up of R in $\rho \subset R = S \times T$. For s in an open subset $V \subset S$ the intersection $\rho \cap (\{s\} \times T)$ depends smoothly on s (i.e., $\rho \cap (V \times T)$ is flat over V). Decreasing V , one may assume that $\rho \cap (V \times T_k)$ is also flat over V for any $k \leq \dim T$. This shows that for $s \in V$ the preimage $T_\bullet^{(s)}$ in R' of the flag $\{s\} \times T_\bullet$ in $\{s\} \times T \subset R$ is well-defined, and depends smoothly on s . In particular, for any $s \in V$ one can canonically define a lifting of T_0 to R' , i.e., there is a mapping $\tau_0: V \rightarrow R'$. Let V' be the image of this mapping.

On the other hand, consider an incomplete flag $\mathcal{R} = \{\mathcal{R}_k\} = S \times \mathcal{T}$ on $S \times T$. It has a strict preimage $\mathcal{R}'$ in R' . Obviously, on the preimage of $V \times T$ the subvariety $\mathcal{R}'_D$ coincides with the image of τ_0 .

Denote the smallest variety of the flag $\mathcal{T}$ by $\{t_0\}$. Since $\mathbf{C}(s)$ dominates U , $\mathbf{C}(s)F_{t_0}$ is smooth on an open subset of S . Decrease V so that $\mathbf{C}(s)F_{t_0}$ is smooth if $s \in V$. Consider $\beta_1 = \beta \circ \tau_0$, it sends V into $\mathfrak{Q}_{k,m}$. Thus for any $s_0 \in V$ there is an open subset $W^{(s_0)} \subset T^{(s_0)}$ such that $W^{(s_0)} \ni \tau_0(s_0)$ and β sends $W^{(s_0)}$ into $\mathfrak{Q}_{k,m}$, hence the family $\mathbf{C}(s_0)F_t$ becomes smooth when restricted/lifted to an open subset $W^{(s_0)}$ of a blowup of T .

Lemma 3.36. *Consider a conformal operator $\mathbf{C}: \mathfrak{MM}(U) \rightarrow \mathfrak{MM}(Z)$ and a smooth family F_t of elements of $\mathfrak{MM}(U)$ parameterized by T . Fix a flag $T_\bullet$ in T . Then the Laurent (=Taylor) coefficients of $\mathbf{C}F_t$ in t at $T_\bullet$ are images by $\mathbf{C}$ of Laurent coefficients of F_t in t .*

Proof. This obvious for $(0, \dots, 0)$ -th Laurent coefficient, since $\mathbf{C}$ sends smooth families to smooth families. Note that $\mathbb{K}$ -linearity of $\mathbf{C}$ and the property of Laurent coefficients we know already imply that

- (1) taking a blowup of T does not change validity of the lemma;
- (2) if φ is a regular scalar function on T , then $\mathbf{C}\varphi(t)F_t = \varphi(t)\mathbf{C}F_t$;
- (3) thus if F_t has a zero of k -th order at a hypersurface $T_{d-1} \subset T$, $\mathbf{C}F_t$ has a zero of k -th order at the same hypersurface;
- (4) If F_t does not depend on t , $\mathbf{C}F_t$ does not depend on t .
- (5) If F_t depends on t as $\varphi(t)F^{(0)}$, $\mathbf{C}F_t$ is $\varphi(t)\mathbf{C}F^{(0)}$.

Consider the case $\dim T = 1$ first. The lemma follows from the above properties alone. Indeed, the first property shows that it is enough to consider normal curves. The third property insures that the k -th Laurent coefficient of $\mathbf{C}F_t$ depends only on l -th coefficients of $\mathbf{C}_t$ with $l \leq k$, and the last one shows that l -th coefficient of $\mathbf{C}_t$ influences l -th coefficient of $\mathbf{C}F_t$ only. (Note that in case $\dim T > 1$ there may be no projection $T \rightarrow T_{d-1}$, which breaks the last argument.)

Now proceed by induction in $d = \dim T$. One may assume that the statement is true when the family is restricted to T_{d-1} . It is enough to show that residues of F_t and $\mathbf{C}F_t$ on T_{d-1} are related by $\mathbf{C}$. On the other hand, since the lemma is already

proven if $d = 1$, one may apply it to curves in T_d which are transversal to T_{d-1} , which proves the statement about the residues. $\square$

This finishes the proof of the proposition, since a blow-up does not change Laurent coefficients. $\square$

By definition, Laurent coefficients at $\mathcal{S} * \mathcal{T}$ are Laurent coefficients at $\mathcal{S} \times T_0$ of partial Laurent coefficients at $S \times \mathcal{T}$. Since partial Laurent coefficients of $\mathbf{C}(s)F_t$ at $S \times \mathcal{T}$ can be written as $\mathbf{C}(s)F_{l_1 \dots l_{D'}}$, and Laurent coefficients of $\mathbf{C}(s)G$ are (by definition) Laurent coefficients of $\mathbf{C}$ applied to $F_{l_1 \dots l_{D'}}$, this finishes the proof of the theorem. $\square$

Remark 3.37. Note that since a blow-up of a manifold is not necessarily a manifold, use of non-smooth flags is necessary (as far as one follows the same arguments). I do not know whether one can omit consideration of non-normal flags.

Remark 3.38. Suppose that conformal fields $\mathbf{C}(s)$ and $\mathbf{C}'(t)$ commute:

$$\mathbf{C}(s)\mathbf{C}'(t) = \mathbf{C}'(t)\mathbf{C}(s).$$

However, a fundamental asymmetry between $\mathcal{S} * \mathcal{T}$ and $\mathcal{T} * \mathcal{S}$ in Theorem 3.33 shows that Laurent coefficients of $\mathbf{C}$ and of $\mathbf{C}'$ *do not necessarily commute!* As we have seen it in Section 3.6, the non-commutativity corresponds to fusions of $\mathbf{C}$ and $\mathbf{C}'$ on subvarieties passing through $s_0 \times t_0$, here $\{s_0\}$ and $\{t_0\}$ are smallest elements of flags on S and T correspondingly.

This observation is the core of the *vertex commutation relations*, considered in the following section.

Remark 3.39. Note that Example 4.3 will provide some pseudo-explanation why it is better to pick up the flag $\mathcal{S} * \mathcal{T}$ than $\mathcal{T} * \mathcal{S}$. Indeed, if Theorem 3.33 would contain $\mathcal{T} * \mathcal{S}$, then (4.2) would have another sign, thus (4.3) would give a wrong sign for commutators of Laurent coefficients of ψ and ψ^+ .

4. CHIRAL ALGEBRAS AND COMMUTATION RELATIONS

Recall that in Section 3.8 it was shown that Laurent coefficients of commuting conformal fields do not necessarily commute, and the commutator may be calculated basing on fusions of these conformal fields. Moreover, in Section 3.5 it was shown that products of several codominant conformal fields have fusions at subvarieties which are determined by pairwise fusions only.

In this section we investigate how the properties of conformal fields studied in Section 3 lead to the notion of a chiral algebra, study vertex commutation relations in the case of a chiral algebra. We also show that Laurent coefficients of the operator ψ, ψ^+ (constructed in Section 2.4) satisfy the same (anti)commutation relations as fermion destruction/creation operators, and that Laurent coefficients of operators $\mathcal{G}(r, s)$ (constructed in Section 2.3) (and their smooth parts) generate Lie algebras $\mathfrak{gl}_\infty$ and its central extension $\mathfrak{g}(\infty)$.

Note that the calculation in the homology space of a vector space with several removed hyperspaces coincide literally with the calculation done in for conformal fields studied in physics-related literature, and in the approach of formal power series. However, they have a different meaning, since we perform them with actual correctly defined families of operators, instead of some symbols which depend on a parameter in a formal sense only.

Consider conformal fields $\mathbf{C}_1, \dots, \mathbf{C}_k$ on the same variety S acting in the same space $\mathfrak{M}\mathcal{M}(Z)$. Say that these conformal fields are *conformally based on* $(S, S^{(1)}, S^{(2)})$, $S^{(1)} \subset S$, $S^{(2)} \subset S \times S$, if all $\mathbf{C}_i$ are smooth in $S \setminus S^{(1)}$, and pairwise products of $\mathbf{C}_i$ and $\mathbf{C}_j$ have singularities at components of $S^{(2)}$ only. If one can take $S^{(1)} = \emptyset$, $S^{(2)} = \emptyset$, call the fields *conformally trivial*.

We start with the simplest case when $S^{(2)}$ is the diagonal $\Delta = \Delta_S$ of S .

4.1. Chiral algebras. Let S be a curve. Consider clean codominant conformal fields $\mathbf{C}(s)$, $\mathbf{C}'(s)$, $s \in S$, acting on $\mathfrak{M}\mathcal{M}(Z)$ which are conformally based on $(S, \emptyset, \Delta)$. By definition one can write²⁰

$$(4.1) \quad \mathbf{C}(r) \mathbf{C}'(s) = \sum_{l \geq 1}^L \frac{\mathbf{D}^{(l)}(r)}{(s-r)^l} + \mathbf{D}(r, s),$$

here $\mathbf{D}(\bullet, \bullet)$ has no pole at $x \approx y$, and the conformal families $\mathbf{D}^{(l)}$ are smooth (otherwise $\mathbf{C}$ and $\mathbf{D}$ would have fusions on subvarieties of $S \times S$ of codimension 2, so they would not be based on $(S, \bullet, \Delta)$). Note that we do not exclude the case $L = \infty$, but in this case $\mathbf{D}^{(l)}F$ with $l \gg 1$ vanishes for any fixed F (or F changing in a finite-dimensional family).

As Proposition 3.28 shows, the fields $\mathbf{D}^{(l)}$ are also codominant. As Remark 3.26 shows, for a generic fixed F (or F changing in a generic finite-dimensional family) and $k \geq 2$ there is an appropriate M such that

$$\left(\prod_{i < j} (s_i - s_j) \right)^M \mathbf{A}_1(s_1) \mathbf{A}_2(s_2) \dots \mathbf{A}_k(s_k) F$$

is smooth in s_i , here $\mathbf{A}_i$ is $\mathbf{C}$ or $\mathbf{C}'$. In particular, the only non-zero residues of $\mathbf{A}_1(s_1) \mathbf{A}_2(s_2) \dots \mathbf{A}_k(s_k)$ are on subvarieties given by several equations $s_{i_m} = s_{j_m}$.

Therefore, fusions of $\mathbf{D}^{(l)}(r)$ with $\mathbf{C}(s)$, $\mathbf{C}'(s)$, or with other $\mathbf{D}^{(m)}(s)$ are all smooth, and defined on the diagonal $r = s$. Same is true for fusions of these fusions, and so on. One obtains a huge family of conformal fields conformally based on $(S, \emptyset, \Delta)$. Following [6] and [14], introduce chiral algebras.

²⁰To avoid defining what $s_i - s_j$ means, one may assume that in this expression $\mathbb{K} = \mathbb{C}$, and it is written in local analytic coordinate system on S , or that the expression is written in a *formal* coordinate system on a completion of S in a particular point.

However, since $\mathbf{D}^{(l)}$ depends on l -jet of such coordinate systems only, one can easily define it the algebraic case too.

Definition 4.1. A *chiral algebra* on S is a vector space $\mathcal{C}$ of conformal families on S acting in $\mathfrak{MM}(Z)$ such that

- (1) If $\mathbf{C}(s) \in \mathcal{C}$ and $\Phi(s)$ is a holomorphic function on S , then $\Phi(s)\mathbf{C}(s) \in \mathcal{C}$;
- (2) any fusion of any two families $\mathbf{C}_{1,2} \in \mathcal{C}$ is defined on $\Delta_S \simeq S$ and is an element of $\mathcal{C}$.²¹

It is possible to codify this definition by defining an *abstract chiral algebra* in terms of operations of taking fusion only, without any reference to the action in $\mathfrak{MM}(Z)$. After such a formalization one can consider the object in the above definition as a *representation* of an abstract chiral algebra in the lattice of vector spaces $\mathfrak{MM}(U)$, $U \subset Z$, compare [1, 6].

Given a chiral algebra spanned by families $\mathbf{C}_i$, $i \in I$, one can interpret (4.1) as *chiral commutation relations*

$$\mathbf{C}_i(r) \mathbf{C}_j(s) = \sum_{\substack{1 \leq l \leq L \\ m}} \Gamma_{ij}^{lm}(r) \frac{\mathbf{C}_m(r)}{(s-r)^l} + [\text{a smooth family near } r=s].$$

4.2. Conformal commutation relations. Show how standard calculations with chiral algebras look in our context.

Consider a curve S and two conformal fields $\mathbf{C}(s)$, $\mathbf{C}'(s)$ based on $(S, \emptyset, \Delta)$. Suppose that

$$\mathbf{C}(s) \mathbf{C}'(s') = q \mathbf{C}'(s') \mathbf{C}(s), \quad q \in \mathbb{K},$$

and that $\mathbf{C}$, $\mathbf{C}'$ have chiral commutation relations

$$\mathbf{C}(r) \mathbf{C}'(s) = \sum_{m \geq 1}^L \Gamma^{(m)}(r) \frac{\mathbf{C}^{(m)}(r)}{(s-r)^m} + [\text{a smooth family near } r=s].$$

Consider a smooth point $s_0 \in S$ and Laurent coefficients $\mathbf{C}_k$, $\mathbf{C}'_{k'}$ of these fields near s_0 . Denote by $\mathbf{C}_k^{(m)}$ the Laurent coefficients of the families $\mathbf{C}^{(m)}(s)$, by $\Gamma_k^{(m)}$ the Laurent coefficients of the functions $\Gamma^{(m)}$.

Proposition 4.2. *These Laurent coefficients have commutation relations*

$$(4.2) \quad \mathbf{C}_k \mathbf{C}'_{k'} - q \mathbf{C}'_{k'} \mathbf{C}_k = - \sum_{\substack{1 \leq m \leq L \\ t \in \mathbb{Z}}} \Gamma_t^{(m)} \binom{-k' - 1}{m - 1} \mathbf{C}_{(k+k'+m-t)}^{(m)},$$

Proof. By Theorem 3.33, $\mathbf{C}_k \mathbf{C}'_{k'}$ is (k, k') -Laurent coefficient of $\mathbf{C}(r) \mathbf{C}'(s)$ w.r.t. the flag $\mathcal{F}_1 = \{(s_0, s_0)\} \subset S \times \{s_0\} \subset S \times S \ni (r, s)$, $\mathbf{C}'_{k'} \mathbf{C}_k$ is (k', k) -Laurent coefficient of $\mathbf{C}'(s) \mathbf{C}(r)$ w.r.t. the flag $\mathcal{F}_2 = \{(s_0, s_0)\} \subset \{s_0\} \times S \subset S \times S \ni (r, s)$.

Thus the difference in the left-hand side of (4.2) is the difference of Laurent coefficients of $\mathbf{C}(r) \mathbf{C}(s)$ w.r.t. the flags $\mathcal{F}_1$ and $\mathcal{F}_2$. As it was seen in Section 3.6, if

²¹The standard definition of the chiral algebra has an additional structure: a linear mapping from vector fields on X to smooth families in $\mathcal{C}$.

Thus to any vector field on X one can associate a smooth family from the vector space.

$\mathbb{K} = \mathbb{C}$, one can write the Laurent coefficients as integrals along the torus $|s| = |r|^K$ in the case of $\mathcal{F}_1$, and $|r| = |s|^K$ in the case of $\mathcal{F}_2$, in both cases with $K \gg 1$, $|s| = \varepsilon \ll 1$.

Note that if $\gamma > \alpha > \beta$ and $\delta < \alpha$

$$\int_{|r|=\alpha} \int_{|s|=\beta} - \int_{|r|=\alpha} \int_{|s|=\gamma} = \int_{|r|=\alpha} \left(\int_{|s|=\beta} - \int_{|s|=\gamma} \right) = - \int_{|r|=\alpha} \int_{|s-r|=\delta}$$

in homology of $U = \mathbb{C}^2 \setminus (\{r=0\} \cup \{s=0\} \cup \{r=s\})$. Thus for any holomorphic function $\Phi(r, s)$ on U the difference of Laurent coefficients of Φ w.r.t. $\mathcal{F}_1$ and $\mathcal{F}_2$ is

$$-\frac{1}{(2\pi i)^2} \int_{|r|=\alpha} \int_{|s-r|=\delta} \Phi(r, s) \frac{dr}{r^{k+1}} \frac{ds}{s^{k'+1}}.$$

Since δ may be arbitrarily small, one should be able to express the integral $\int_{|y-x|=\delta}$ via the singular part of Φ on $r = s$ only. Indeed, the part $\frac{dr}{r^{k+1}} \frac{ds}{s^{k'+1}}$ is smooth near $r = s$, $r \neq 0$, so only the poles of $\Phi(r, s)$ along $r = s$ matter. Suppose that $\Phi(r, s) = \sum_{1 \leq l \leq L} \frac{\Phi^{[l]}(r)}{(s-r)^l}$, with Laurent coefficients of $\Phi^{[l]}(r)$ being $\Phi_k^{[l]}$.

Writing the holomorphic on $\{(r, s) \mid r \approx s, r \neq 0\}$ differential form $\frac{dr}{r^{k+1}} \frac{ds}{s^{k'+1}}$ as a Taylor series in $(s-r)$ for a fixed r , one gets

$$\frac{dr}{r^{k+1}} \frac{ds}{s^{k'+1}} = \sum_{m \geq 0} \binom{-k'-1}{m} \frac{dr}{r^{k+k'+2+m}} (s-r)^m d(s-r)$$

thus the difference of Laurent coefficients of Φ is

$$-\frac{1}{2\pi i} \int_{|r|=\alpha} \sum_{m \geq 0} \binom{-k'-1}{m} \Phi^{[m+1]}(r) \frac{dr}{r^{k+k'+2+m}} = - \sum_{m \geq 0} \binom{-k'-1}{m} \Phi_{k+k'+1+m}^{[m+1]}$$

Applying this to $\Phi(r, s) = \mathbf{C}(r) \mathbf{C}'(s) F$, one obtains

$$\mathbf{C}_k \mathbf{C}'_{k'} - q \mathbf{C}'_{k'} \mathbf{C}_k = - \sum_{\substack{0 \leq m \leq L-1 \\ t \in \mathbb{Z}}} \Gamma_t^{(m+1)} \binom{-k'-1}{m} \mathbf{C}_{(k+k'+m-t+1)}^{(m+1)}.$$

In the case of generic $\mathbb{K}$ one gets the same answer, since homology of $S \times S \setminus (\{s_0\} \times S \cup S \times \{s_0\} \cup \Delta)$ does not depend on the field. $\square$

If the fields $\mathbf{C}^{(m)}(s)$ are independent, one can incorporate coefficients $\Gamma^{(m)}(s)$ into the fields, and assume $\Gamma^{(m)}(s) = 1$. Then

$$\mathbf{C}_k \mathbf{C}'_{k'} - q \mathbf{C}'_{k'} \mathbf{C}_k = - \sum_{1 \leq m \leq L} \binom{-k'-1}{m-1} \mathbf{C}_{(k+k'+m)}^{(m)}.$$

Example 4.3. Consider fields $\psi(s), \psi^+(s)$. Denote the Laurent coefficients ψ_k, ψ_k^+ . One concludes that the anticommutators satisfy

$$(4.3) \quad [\psi_k, \psi_{k'}]_+ = 0, \quad [\psi_k^+, \psi_{k'}^+]_+ = 0, \quad [\psi_k, \psi_{k'}^+]_+ = -\delta_{k+k'+1},$$

since the only fusion of these fields is $\mathbf{1}$, which does not depend on s , thus has all the Laurent coefficients but the 0-th one being 0.

One can easily check the sign in this identity, since by Corollary 2.5

$$\psi_{-1}(1) = 0, \quad \psi_0(1) = T, \quad \psi_0^+(1) = T^{-1}, \quad \psi_{-1}(T^{-1}) = -T^{-1}\psi_0(1) = -1.$$

4.3. The family $\mathcal{G}$. Consider a conformal family $\mathcal{G}(r, s)$ on $\mathbb{A}^2$ defined in Section 2.3:

$$(4.4) \quad (\mathcal{G}(r, s)F)(\xi) = \frac{1}{s-r} \frac{\xi(r)}{\xi(s)} F\left(\frac{z-r}{z-s} \xi(z)\right).$$

As calculations in Section 2.4 show, $\mathcal{G}(r, s) = \psi(r)\psi^+(s)$.

As it was seen in Section 2.3, the family $\mathcal{G}(r, s)$ is based on $(\mathbb{A}^2, D_0, D_1)$ with $D_0 = \Delta_{\mathbb{A}^1} \subset \mathbb{A}^2$, $D_1 = \{(r_1, s_1, r_2, s_2) \mid s_1 \neq r_2, s_2 \neq r_1\}$. The residue of $\mathcal{G}(r, s)$ on $r = s$ is $\mathbf{1}$. In fact the definition of $\mathcal{G}(r, s)$ shows that in fact $\mathcal{G}(r, s)$ can be considered as a family based on $((\mathbb{P}^1)^2, \Delta_{\mathbb{P}^1}, \bar{D}_1)$, here $\bar{D}_1 = \{(r_1, s_1, r_2, s_2) \mid s_1 \neq r_2, s_2 \neq r_1\} \subset (\mathbb{P}^1)^4$.

Consider Laurent coefficients of $\mathcal{G}(r, s)$ for $r, s \approx 0$. Consider two different cases, one corresponds to the flag $\mathcal{F}^< = \{(0, 0)\} \subset \mathbb{A}^1 \times \{0\} \subset \mathbb{A}^2$ (i.e., taking a contour with $|r| < |s|$), another to the flag $\mathcal{F}^> = \{(0, 0)\} \subset \{0\} \times \mathbb{A}^1 \subset \mathbb{A}^2$ (or to $|r| > |s|$). Denote the corresponding Laurent coefficients $\mathcal{G}_{nm}^<$ and $\mathcal{G}_{nm}^>$ (assume that n gives the power of r , m the power of s). In the same way as we did it with ψ and ψ^+ , one gets commutation relations for $\mathcal{G}_{nm}^{<, >}$ (a more complicated case of $\mathcal{G}_{nm}$ is done in details below):

$$\begin{aligned} [\mathcal{G}_{nm}^<, \mathcal{G}_{kl}^<] &= \mathcal{G}_{nl}^< \delta_{m+k+1} - \mathcal{G}_{km}^< \delta_{n+l+1}, \\ [\mathcal{G}_{nm}^>, \mathcal{G}_{kl}^>] &= \mathcal{G}_{nl}^> \delta_{m+k+1} - \mathcal{G}_{km}^> \delta_{n+l+1}. \end{aligned}$$

On the other hand, since $\mathcal{G}(r, s)$ has a simple pole at $r \approx s$ with residue $-\mathbf{1}$, one can write the integrals for $\mathcal{G}_{nm}^< - \mathcal{G}_{nm}^>$ as

$$\int_{|r|=\alpha} \int_{|s|=\beta} - \int_{|r|=\gamma} \int_{|s|=\beta} = \int_{|r-s|=\alpha'} \int_{|s|=\beta}, \quad \alpha, \alpha' < \beta < \gamma,$$

thus $\mathcal{G}_{nm}^< - \mathcal{G}_{nm}^> = \delta_{n+m+1} \mathbf{1}$. To understand the relationship between $\mathcal{G}_{nm}^<$ and $\mathcal{G}_{nm}^>$ better, consider the smooth part $\mathcal{G}^{\text{smooth}}(r, s)$ of $\mathcal{G}(r, s)$:

$$\mathcal{G}^{\text{smooth}}(r, s) = \mathcal{G}(r, s) - \frac{1}{s-r} \mathbf{1}.$$

Denote by $\mathcal{G}_{nm}$ the Laurent coefficients of the smooth function $\mathcal{G}^{\text{smooth}}$ (they do not depend on the choice of the flag). Writing Laurent series for $\frac{1}{r-s}$ near flags $\mathcal{F}_1$ and $\mathcal{F}_2$ (i.e., in regions $|r| < |s|$ and $|r| > |s|$), one gets

$$\mathcal{G}_{nm} = \mathcal{G}_{nm}^< - \delta_{n+m+1} \chi_{n \geq 0} = \mathcal{G}_{nm}^> + \delta_{n+m+1} \chi_{n < 0}.$$

Here χ_S is the characteristic function of a set S . Using these relationship, one can easily get the commutation relations for $\mathcal{G}_{nm}$:

$$[\mathcal{G}_{nm}, \mathcal{G}_{kl}] = \mathcal{G}_{nl}\delta_{m+k+1} - \mathcal{G}_{km}\delta_{n+l+1} + 2\delta_{n+l+1}\delta_{m+k+1} (\chi_{n \geq 0} - \chi_{k \geq 0}).$$

Indeed, the divisor of poles for $\mathcal{G}(a, b)\mathcal{G}(c, d) = -\psi(a)\psi(c)\psi^+(b)\psi^+(d)$ is

$$\frac{1}{b-a}\psi(c)\psi^+(d) - \frac{1}{b-c}\psi(a)\psi^+(d) - \frac{1}{d-a}\psi(c)\psi^+(b) + \frac{1}{d-c}\psi(a)\psi^+(b),$$

since $\mathcal{G}^{\text{smooth}}(a, b)\mathcal{G}^{\text{smooth}}(c, d)$ coincides with

$$\mathcal{G}(a, b)\mathcal{G}(c, d) - \frac{1}{b-a}\mathcal{G}^{\text{smooth}}(c, d) - \frac{1}{d-c}\mathcal{G}^{\text{smooth}}(a, b) - \frac{1}{(b-a)(d-c)}$$

the poles coincide with poles of

$$\frac{1}{(b-a)(d-c)} - \frac{2}{(b-c)(d-a)} - \frac{1}{b-c}\mathcal{G}^{\text{smooth}}(a, d) - \frac{1}{d-a}\mathcal{G}^{\text{smooth}}(c, b).$$

To calculate the commutation relations for Laurent coefficients one should take difference of integrals along $|a| \approx |b| < |c| \approx |d|$ and $|a| \approx |b| > |c| \approx |d|$. Take contours C_1 with $|a| < |b| < |c| < |d|$ and C_2 with $|c| < |d| < |a| < |b|$.

When pulling C_1 to C_2 preserving $|a| < |b|$ and $|c| < |d|$, we cross poles at $b = c$ and at $a = d$, thus the first summand out of four gives no contribution at all. During intersection of $b = c$ the fourth one gives no contribution, and during intersection of $a = d$ the third gives no contribution.

The pole at $b = c$ has residue $\frac{2}{d-a} + \mathcal{G}^{\text{smooth}}(a, d)$ w.r.t. $c - b$, thus its contribution to $[\mathcal{G}_{nm}, \mathcal{G}_{kl}]$ is

$$\frac{1}{(2\pi i)^2} \int_{|a| < |d|} \frac{da}{a^{n+1}} \frac{dd}{d^{l+1}} \left(\frac{2}{d-a} + \mathcal{G}^{\text{smooth}}(a, d) \right) \delta_{k+m+1},$$

i.e., it is δ_{k+m+1} times the Laurent coefficient of $\frac{2}{d-a} + \mathcal{G}^{\text{smooth}}(a, d)$. Note that during this intersection $|a| < |d|$, thus the contribution is

$$+2\chi_{n \geq 0}\delta_{k+m+1}\delta_{n+l+1} + \delta_{k+m+1}\mathcal{G}_{nl}.$$

Similarly, the pole at $a = d$ gives contribution of δ_{n+l+1} times the Laurent coefficient of $-\frac{2}{b-c} - \mathcal{G}^{\text{smooth}}(c, b)$, and during this intersection $|b| > |c|$. Thus this pole gives

$$-2\chi_{k \geq 0}\delta_{k+m+1}\delta_{n+l+1} - \delta_{n+l+1}\mathcal{G}_{km}.$$

This proves commutation relations for $\mathcal{G}_{ab}$.

4.4. Generators X_{ij} . Let $X_{ij} = \mathcal{G}_{i-j-1}$, same with $X_{ij}^<$ and $X_{ij}^>$. It is obvious now that $X_{ij}^<$ commute as elementary matrices of infinite size (same with $X_{ij}^>$), thus generate $\mathfrak{gl}_\infty$, and X_{ij} commute as generators of central extension $\mathfrak{g}(\infty)$ of $\mathfrak{gl}_\infty$.

Fix $F \in \mathfrak{MM}(Z)$. Note that $X_{ij}^<F$ is a Laurent coefficient of the family $\mathcal{G}(r, s)F$. This function is smooth if $r, s \notin \text{Supp } F$ and $r \neq s$. Since $\text{Supp } F$ may contain 0, one can conclude that $X_{ij}^<F$ is a Laurent coefficient of a function which is smooth

in $0 < |r| < |s| \ll 1$. Similarly, $X_{ij}F$ is a Laurent coefficient of a function which is smooth in $0 < |r|, |s| \ll 1$.

Consider a collection a_{ij} , $i, j \in \mathbb{Z}$, such that the sum $\sum a_{ij}\phi_{ij}$ contains only a finite number of non-zero summands if ϕ_{ij} are Laurent coefficients of a meromorphic function ϕ which is smooth in $0 < |r| < |s| \ll 1$. Then $a_{ij} = 0$ if $j > J(-i) - i$ or $i > I(-i - j)$, here J and I are appropriate non-decreasing functions. If the same is true if ϕ is smooth in $0 < |r| < |s| \ll 1$, then $a_{ij} = 0$ if $j > J(-i)$ or $i > I(-j)$. This is compatible with the facts that $\mathfrak{g}(\infty)$ contains arbitrary sums $\sum a_{ij}X_{ij}$, with $a_{ij} = 0$ if $j < J(-i)$ or $i > I(j)$, and that cocycle for $\mathfrak{g}(\infty)$ is a coboundary on sums $\sum a_{ij}X_{ij}$, with $a_{ij} = 0$ if $j < -J(-i) + i$ or $i > I(j - i)$.

One can construct geometric realizations of some particular infinite sums $\sum a_{ij}X_{ij}$. Consider

$$Y_{kl} = \sum_{i \in \mathbb{Z}} \binom{i}{k} X_{i-i-k-l} = \sum \binom{i}{k} \mathcal{G}_{i-i+k+l-1}^{\text{smooth}}, \quad k \geq 0.$$

Taking into account that $\mathcal{G}_{ab}$ are Laurent coefficients for $\mathcal{G}(r, s)$, one can see that

$$Y_{kl} = \frac{1}{(2\pi i)^2} \sum_i \binom{i}{k} \iint \mathcal{G}(r, s) r^{-i-1} s^{i-k-l} dr ds = \frac{1}{2\pi i} \int \mathcal{G}^{\text{smooth}(k,0)}(s, s) \frac{ds}{s^l},$$

so Y_{kl} is the $(l-1)$ -th Laurent coefficient of $-k$ -th residue of $\psi(r)\psi^+(s)$ at $r = s$ (the vector field v being $\partial/\partial r$).

Let us calculate Laurent coefficients of 0-th residue $B(s)$ of $\psi(r)\psi^+(s)$ at $r = s$. By (4.4)

$$(B(s)F)\xi = -\frac{\xi'(s)}{\xi(s)}F(\xi(z)) - \frac{dF\left(\left(1 + \frac{\tau}{z-s}\right)\xi(z)\right)}{d\tau} \Big|_{\tau=0}.$$

In coordinates

$$\xi(z) = Te^{\sum_k t_k z^k}$$

one can write B as

$$B(s)F(T, t) = -\left(\sum_{l \geq 1} lt_l s^{l-1}\right)F(T, t) + \sum_{l \geq 1} \frac{1}{s^{l+1}} \partial_{t_l} F(T, t) + \frac{1}{s} \frac{\partial_T}{T} F(T, t).$$

We conclude that $Y_{0l} = -lt_l$ if $l > 0$, $Y_{0l} = \partial_{t_{-l}}$ if $l < 0$, and $Y_{00} = \frac{\partial_T}{T}$.

Taking the “holomorphic in $|s| \leq 1$ part” of the function $B(s)$, one gets the standard boson creation operator in the Fock space $\mathbb{K}[t_1, t_2, \dots]$, and the “holomorphic in $|s| \geq 1$ part” gives the boson destruction operator. Moreover, one can see that $B(s)$ lies in the completion of algebra generated by $\psi(s)$ and $\psi^+(s)$. On the other hand, the formulae for vertex operators show that visa versa, $\psi(s)$ and $\psi^+(s)$ are in the completion of algebra generated by Laurent coefficients of $B(s)$.

Note that Laurent coefficients of $\psi(s)$, $\psi^+(s)$ commute as fermion destruction/creation operators (see (4.3)), and Laurent coefficients of $B(s)$ commute as t_k , $\frac{\partial}{\partial t_k}$, i.e., as

boson creation/destruction operators. This the base of so called *boson-fermion correspondence*, which is studied in the following section.

5. BOSON-FERMION CORRESPONDENCE

In this section we establish a *geometric boson-fermion correspondence* (in dimension 1+1), show that this correspondence provides heuristic for considering $\mathcal{M}(\mathcal{M}(\mathbb{P}^1))$ as a configuration space for a system of fermions close to Dirac's vacuum state, show that the Fock space coincides with the localization of $\mathcal{M}(\mathcal{M}(\mathbb{P}^1))$ near $0 \in \mathbb{P}^1$, and investigate the relation of the geometric boson-fermion correspondence with the usual, algebraic boson-fermion correspondence.

The mechanical description of a physical system consists of two parts: kinematics describes the set of possible *states* of the system, and dynamics examines how the states change when time goes. As far as we are concerned here, dynamics is not a question, so we restrict our attention to kinematics only.

The quantum mechanics describes *states* of a system as elements of a Hilbert space called (*quantum*) *configuration space*. In the case of d space dimensions, this space may be the Hilbert space of square-integrable functions on $\mathbb{R}^d$. A particle “concentrated” near a point $Y \in \mathbb{R}^d$ is described by a function with a support in a neighborhood of Y . Two proportional vectors represent the same state of the system.

In general the relation to the classical mechanics is that the quantum configuration space is the vector space of functions on *classical configuration space* ($\mathbb{R}^d$ above), and functions with support in a small neighborhood of the given point represent quantum states which are “almost classical”.

In what follows we ignore the Hilbert structure completely (though it can be added back later at some moment). This would allow one to consider more general functions than square-integrable one, say, one can consider functions of the form

$$(5.1) \quad \sum_{k=1}^K a_k \delta(Y - Y_k),$$

vectors Y_i being some points in $\mathbb{R}^d$. If $K = 1$, the above state describes a particle concentrated *exactly* at Y_1 , thus a classical state.

5.1. Bosonic space. What is described above is the approach of *primary quantization*, when the number of particles is fixed, thus we can model m particles in d -dimensional space by one particle in md -dimensional space. To allow the number of particle to be variable, one needs some modifications.

In what follows we assume $d = 1$.

If the quantum system is a union of two parts, each one described by a (Hilbert) space $H_{1,2}$, then the system in whole is described by the tensor product $H_1 \otimes H_2$, vector $v_1 \otimes v_2$ describing the state of the system when the first part is in the state v_1 ,

the second in the state v_2 . Thus the system of n particles having the space of states V each is described by

$$\underbrace{V \otimes \cdots \otimes V}_{n \text{ times}}.$$

If the particles are indistinguishable one needs to require that $v_1 \otimes v_2$ and $v_2 \otimes v_1$ describe the *same* state of the system, thus in this case the space of states of the system is some quotient of the above space. In this quotient the images of $v_1 \otimes v_2$ and $v_2 \otimes v_1$ should be proportional.

A little bit of representation theory of the symmetric group $\mathfrak{S}_n$ says that there are only two such quotients: the symmetric power $S^n V$ of V and the skewsymmetric one $\Lambda^n V$. Which one is relevant is determined by the physical properties of the particle in question. Particles for which S^n should be taken are called *bosons*, the other ones *fermions*.

Summing up, we conclude that the configuration space of n bosons on a line $\mathbb{R}$ consists of symmetric functions on $\mathbb{R}^n$. If one consider states of one boson of the form (5.1), one should consider the space B_n which consists of symmetric generalized functions²² on $\mathbb{R}^n$ which are sums of δ -functions.

To consider variable number of particles, one should consider the direct sum of spaces of the above form for all possible values of n , $0 \leq n < \infty$:

$$B = \sum_{k \geq 0} B_k.$$

5.1.1. *The dual space.* The above space is too convoluted to work with, thus in fact we are going to describe *some* dual space to this one. Note that there is a canonically defined non-degenerate pairing between the above space of generalized functions on $\mathbb{R}^n$ and the space of symmetric polynomials on $\mathbb{R}^n$. This pairing is the calculation of value

$$\langle P(Y), \delta(Y - Y_0) \rangle = P(Y_0), \quad Y \in \mathbb{R}^n.$$

5.1.2. *Quotient space.* One can identify the space of symmetric polynomial with the space of polynomials in new variables $(\sigma_1, \dots, \sigma_n)$, which are elementary symmetric functions in the old variables z_i . Associate with the collection $(\sigma_1, \dots, \sigma_n)$ the polynomial

$$\pi(z) = z^n - \sigma_1 z^{n-1} + \sigma_2 z^{n-2} - \cdots \pm \sigma_n = (z - z_1)(z - z_2) \dots (z - z_n).$$

Denote the manifold of polynomials of degree n with the leading coefficient 1 by $\mathcal{P}_n$. One concludes that the dual space to the bosonic configuration space can be described as the space of polynomials on the affine variety $\mathcal{P}_n$. Similarly, the bosonic space itself can be identified with the space spanned by δ -functions on $\mathcal{P}_n$.

²²I.e., generalized functions which are invariant w.r.t. permutations of coordinates.

To consider variable number of particles, one should consider the variety

$$\mathcal{P} = \bigcup_n \mathcal{P}_n.$$

5.2. Boson creation operator. Fix a point $z_0 \in \mathbb{R}$. Consider the transformation of adding a boson at the point z_0 . Obviously, if the initial state of the system was described by $\delta(Y - Y_0)$, $Y, Y_0 \in \mathbb{R}^n$, the new state is $\delta(\bar{Y} - \bar{Y}_0)$, $\bar{Y}, \bar{Y}_0 \in \mathbb{R}^{n+1}$, $\bar{Y}_0 = (z_0, Y_0)$. In other words, it is described by the mapping

$$\mathbb{R}^n \rightarrow \mathbb{R}^{n+1}: (z_1, \dots, z_n) \mapsto (z_0, z_1, \dots, z_n)$$

(followed by symmetrization, of course).

Similarly, in the language of quotient space, this operator is

$$\mathcal{P}_n \rightarrow \mathcal{P}_{n+1}: \pi(z) \mapsto (z - z_0)\pi(z).$$

Multiplication law for polynomials shows that this mapping is compatible with the affine structures on the spaces $\mathcal{P}_\bullet$.

In the dual language, the dual operator sends symmetric polynomial on $\mathbb{R}^{n+1}$ to symmetric polynomials on $\mathbb{R}^n$:

$$P(z, z_1, \dots, z_n) \mapsto Q(z_1, \dots, z_n), \quad Q(z_1, \dots, z_n) = P(z_0, z_1, \dots, z_n).$$

In the dual language for the quotient, when one considers polynomials $f(\pi)$ on $\mathcal{P}$, this mapping is written as

$$(5.2) \quad \alpha_{z_0}: f \mapsto \alpha_{z_0}f, \quad (\alpha_{z_0}f)(\pi) = f((z - z_0)\pi).$$

Remark 5.1. Note that this formula for the boson creation operator coincides with the formula for the operator $\mathbf{M}_\xi$ from Section 1.3 in the case $\xi(z) = z - z_0$ and the rational function F being the polynomial π . Thus it is the restriction of operator $a^{-1}(s)$ considered in Section 0.

5.3. Naive boson-fermion correspondence. Consider the configuration space for the system of n fermions on the line $\mathbb{R}$. As described above, it is the space of skew-symmetric functions on $\mathbb{R}^n$. To use the dual language, note that there is a canonical pairing of the space of skew-symmetric generalized functions on $\mathbb{R}^n$ (with compact support) and the space of skew-symmetric polynomials.

Thus we consider the space of skew-symmetric polynomials on $\mathbb{R}^n$ as the dual space to the configuration space. On the other hand, any skew-symmetric polynomial q in z_i can be written as

$$q(z_i) = \text{Disc}(z_i)p(z_i), \quad \text{Disc}(z_i) = \prod_{i < j} (z_i - z_j),$$

with p being a symmetric polynomial.

Call the correspondence $p \iff q$ the *(geometric) boson-fermion correspondence*. This correspondence makes it possible to describe states of fermions in terms of states of bosons, via an explicit isomorphism of configuration spaces.

5.4. Fermion creation operator. Consider a transformation of configuration space which corresponds to adding a fermion at the point $z_0 \in \mathbb{R}$. As in the case of bosons, it is described by the mapping

$$\mathbb{R}^n \rightarrow \mathbb{R}^{n+1}: (z_1, \dots, z_n) \mapsto (z_0, z_1, \dots, z_n),$$

the only difference being that it should be followed by skewsymmetrization (instead of symmetrization). In the dual language, the formula is the same as for bosons:

$$P(z, z_1, \dots, z_n) \mapsto Q(z_1, \dots, z_n), \quad Q(z_1, \dots, z_n) = P(z_0, z_1, \dots, z_n),$$

but it is applied to skew-symmetric functions. Let us rewrite this operator in terms of bosonic configuration space using the boson-fermion correspondence. Let $P_{\text{bos}}, Q_{\text{bos}}$ be the images of P and Q under boson-fermion correspondence:

$$P(z, z_i) = \text{Disc}(z, z_i) \cdot P_{\text{bos}}(z, z_i), \quad Q(z_i) = \text{Disc}(z_i) \cdot Q_{\text{bos}}(z_i).$$

One can see that in terms of bosons this operator is

$$\begin{aligned} P_{\text{bos}}(z, z_1, \dots, z_n) &\mapsto Q_{\text{bos}}(z_1, \dots, z_n), \\ Q_{\text{bos}}(z_1, \dots, z_n) &= \frac{\text{Disc}(z_0, z_i)}{\text{Disc}(z_i)} P_{\text{bos}}(z_0, z_1, \dots, z_n) \\ &= \left(\prod_{k=1}^n (z_0 - z_k) \right) P_{\text{bos}}(z_0, z_1, \dots, z_n). \end{aligned}$$

On the other hand, in terms of dual language for the quotient, the factor $(\prod_{k=1}^n (z_0 - z_k))$ is just $\pi(z_0)$, thus the fermion creation operator is

$$(5.3) \quad \psi_{z_0}^+ : f \mapsto \psi_{z_0}^+ f, \quad (\psi_{z_0}^+ f)(\pi) = \pi(z_0) f((z - z_0)\pi).$$

Remark 5.2. Note that the last formula coincides with Equation (2.1) of Section 2.4 for ψ^+ in the case of the rational function ξ being the polynomial π .

5.5. Fock space I. The above description of the configuration space in the fermion case is extremely naive, since it deals with collections of a finite number of fermions. However, the physically meaningful picture deals with collections close to the Dirac vacuum state.

This state corresponds to having infinitely many fermions, all the fermions with “negative energy” being present, all with “positive energy” being absent. (This is not relevant to the discussion in this paper, but we state it nevertheless: in terms of functions on $\mathbb{R}$, we say that the state $f(z)$, $z \in \mathbb{R}$, has negative energy if its Fourier transform $\widehat{f}(\xi)$ vanishes for positive ξ , similarly for positive energy. However, we use “complex logarithmic” coordinate system $z = e^{i\tilde{z}}$, thus will use Laurent transform instead of Fourier transform.)

What follows may have no mathematical meaning, but it provides a heuristic for relation of conformal field theory to the secondary quantization in 1-dimensional case. Note that this heuristic is yet incomplete, compare with Remark 5.3.

Fix a finite collection S of points on the line $\mathbb{R}$. The classical states which correspond to bosons sitting at the points of S and at some other points are described by the subset

$$p_S \cdot \mathcal{P} \subset \mathcal{P}, \quad p_S(z) = \prod_{s \in S} (z - s).$$

Clearly, the same is true for fermion states after applying boson-fermion correspondence.

If we consider a classical state corresponding to S as a new *ground state*,²³ then the parameter change $\pi = p_S \cdot \pi_S$ describes a new coordinate π_S on the set of admissible states such that the ground state corresponds to $\pi_S = 1$.

Suppose that the collection S is very big, but one want to consider classical states which are *close* to S , say, add only a few additional bosons or fermions. One finishes with the subset of classical configuration space:

$$p_S \cdot \mathcal{P}_{\leq K} \subset \mathcal{P}, \quad \mathcal{P}_{\leq K} = \bigcup_{k \leq K} \mathcal{P}_k.$$

If we allow taking a few particles from the collection S as well as adding a few particles, the set of classical states becomes

$$(p_S \cdot \mathcal{Q}_K) \cap \mathcal{P} = p_S \cdot \mathcal{Q}_K^{(S)} \subset \mathcal{P}.$$

Here $\mathcal{Q}_K$ is the set of rational functions of the form $\frac{p_1}{p_2}$, $p_{1,2} \in \mathcal{P}_{\leq k}$, and $\mathcal{Q}_K^{(S)} = \mathcal{Q}_K \cap (p_S^{-1} \cdot \mathcal{P})$.

We conclude that if we consider classical states *close to* the ground state S (in the above sense), then the corresponding subset of configurations space can be identified with the set $\mathcal{Q}_K^{(S)}$ for some small value of K . When we increase the set S , the set $\mathcal{Q}_K^{(S)}$ increases and becomes closer and closer to $\mathcal{Q}_K$. In other words, the set $\mathcal{Q}_K$ describes *relative* classical states of a configuration of particles with respect to a *very big* collection S , or, in other words, a very populated ground state. The corresponding quantum states are described by functions on $\mathcal{Q}_K$.

Let us describe boson and fermion creation operators in terms of representation by functions on $\mathcal{Q}_K$. Obviously, the boson creation operator α_{z_0} is described by the same formula (5.2), thus the expression for this operator does not depend on S . To write down the fermion creation operator $\psi_{z_0}^+$, one need to fix the family S . Writing down $\psi_{z_0}^+$ in terms of $\pi = p_S \cdot \pi_S$, $\pi_S \in \mathcal{Q}_K^{(S)}$, one gets

$$\psi_{z_0}^{+S} : f \mapsto \psi_{z_0}^{+S} f, \quad (\psi_{z_0}^{+S} f)(\pi_S) = p_S(z_0) \pi_S(z_0) f((z - z_0) \pi_S), \quad \pi_S \in \mathcal{Q}_K^{(S)},$$

or $\psi_{z_0}^{+S} = p_S(z_0) \psi_{z_0}^+$. Since the operators $\psi_{z_0}^+$ and $\psi_{z_0}^{+S}$ differ by a constant $p_S(z_0)$, and vectors which differ by a constant describe the same quantum state, one can safely use $\psi_{z_0}^+$ instead of $\psi_{z_0}^{+S}$.

²³Say, one can suppose that the Hamiltonian for some physical system has this collection as a minimum.

It is obvious that the (extended as above) operator α_{z_0} sends polynomial functions on $\mathcal{Q}_K$ to polynomial functions on $\mathcal{Q}_{K+1}$. This is not true for $\psi_{z_0}^+$, since $\pi_S \mapsto \pi_S(z_0)$ is not a *polynomial* function of $\pi_S \in \mathcal{Q}_K^{(S)}$. However, $\pi_S \mapsto \pi_S(z_0)$ is a *rational* function of $\pi_S \in \mathcal{Q}_K^{(S)}$, thus the extended operator $\psi_{z_0}^+$ sends rational functions on $\mathcal{Q}_K$ to rational functions on $\mathcal{Q}_{K+1}$.

Obviously, the operators α_{z_0} , $\psi_{z_0}^+$ are compatible with natural inclusions $\mathcal{Q}_K \hookrightarrow \mathcal{Q}_{K+L}$, $L \geq 0$, thus one can extend them to rational functions on

$$\mathcal{Q} = \lim_{\rightarrow} \mathcal{Q}_K.$$

Note that the description of $\mathcal{Q}$ as of inductive limit leads to the same definition of the vector space of rational functions on $\mathcal{Q}$ as we had in the Section 1.1, taking into account $\mathcal{Q} = \mathcal{M}(\mathbb{P}^1)$.

Heuristic . *Extend operators α_{z_0} and $\psi_{z_0}^+$ to the space of rational functions on $\mathcal{Q}$. These operators describe addition of a boson or a fermion (correspondingly) to a collection of particles relative to a very populated ground state.*

Remark 5.3. Note that the given heuristic does not describe why the ground state which is a limit of collections of particles with more and more of them should correspond to the *Fock ground state*. However, as we will see later, it is possible to add *fermion destruction operators* ψ_{z_0} which have correct commutations relations with the operators $\psi_{z_0}^+$, and which annihilate the quantum state corresponding to the classical state $\pi = 1$, i.e., the classical ground state.

Some additional arguments are required to find a heuristic for this coincidence.

5.6. Fock space II. Now we can show that the operators constructed in Section 5.5 can be indeed considered as fermion creation operator. One needs to construct a subspace of the space of meromorphic functions on $\mathcal{Q}$ (the *space of states*) and to show that there are fermion destruction operators $\psi(s)$ such that

- (1) they satisfy correct (anti)commutation relations with operators $\psi^+(s)$;
- (2) the negative Laurent coefficients of $\psi(s)$ kill the vacuum state;
- (3) the negative Laurent coefficients of $\psi^+(s)$ kill the vacuum state;
- (4) all the states may be obtained as combinations of Laurent coefficients of $\psi(s)$ and $\psi^+(s)$ applied to the vacuum state.

Recall that in the boson case the classical vacuum state corresponds to having no bosons, thus to the polynomial 1 which has no roots, so the quantum vacuum state corresponded to the function on $\cup_{k \geq 0} \mathcal{P}_k$ which vanishes on $\mathcal{P}_{\geq 1}$, and is 1 on the one-point set $\mathcal{P}_0$. As seen in Section 5.5, introduction of Fock space corresponds to multiplication by an arbitrary polynomial π_S , and taking the limit when the degree of this polynomial goes to infinity. These multiplications (for different π_S of fixed degree k) spread the point $\mathcal{P}_0$ over the whole space $\mathcal{P}_k$. Thus it is natural to consider a constant function 1 on $\mathcal{Q} = \mathcal{M}(\mathbb{P}^1)$ as a natural analogue of the above boson ground state.

Since this function 1 is translation-invariant, one can see that $M_\xi 1 = 1$, thus $\psi(s) 1 = E_s$. On the other hand, E_s depends smoothly on s , thus negative Laurent coefficients of $\psi(s)$ kill 1. This is obviously so in $\mathfrak{MM}(\mathbb{P}^1)$.

Similarly, consider the operators $\psi^+(s)$ acting on meromorphic functions on $\mathcal{Q}$. Clearly, $\psi^+(s) 1 = E_s^{-1}$. This expression depends smoothly on s considered as an element of $\mathfrak{MM}(\mathbb{P}^1)$.

The (anti)commutation relations for Laurent coefficients of $\psi(s)$ and $\psi^+(s)$ were already proved in (4.3).

Consider now the completeness part. Note that by Corollary 2.5 the action of Laurent coefficients of $\psi(s)$ and $\psi^+(s)$ preserves the subspace of $\mathfrak{MM}(\mathbb{P}^1)$ consisting of polynomials in T , T^{-1} and t_k . This means that the space of states is contained in the space of these polynomials.

On the other hand, as seen in Section 4.4, the operators $T \frac{\partial}{\partial T}$, $\frac{\partial}{\partial t_k}$ and the operators of multiplication by t_k may be expressed via Laurent coefficients of $\psi(s)$ and $\psi^+(s)$. All these operators preserve the degree in T , but $\psi(s)$ and $\psi^+(s)$ do not, thus *all* the polynomials in T , T^{-1} and t_k may be obtained by application of combinations of Laurent coefficients of $\psi(s)$ and $\psi^+(s)$.

This would finish the description of the Fock space as a subspace of the space of meromorphic functions on $\mathcal{Q}$, but note that above we had shown that E_s and E_s^{-1} depend smoothly on s as elements of $\mathfrak{MM}(\mathbb{P}^1)$, not as elements of the space of states. However, the formulae of Corollary 2.5 show that this is true in the space of states too.

We conclude that the subspace of $\mathfrak{MM}(\mathbb{P}^1)$ which corresponds to the Fock space is a subspace of $\mathfrak{MM}(\widehat{Z})$, here $\widehat{Z}$ is the localization of $\mathbb{P}^1$ near $0 \in \mathbb{P}^1$. Recall that $\mathfrak{MM}(\widehat{Z})$ consists of rational expressions of $E_0^{\frac{dz^k}{dz^k}}$, $k \geq 0$, and the subspace consists of polynomials in $E_0^{\frac{dz^k}{dz^k}}$, $k \geq 0$, and in E_0^{-1} .

5.7. Bosons: geometric vs algebraic. Our approach introduced a geometric connection between a boson with the creations operators given by Laurent coefficients of the conformal field $\alpha(s) = a^{-1}(s)$, and a fermion with the creation operators given by Laurent coefficients of the conformal field $\psi^+(s)$. However, usually people consider a different correspondence, which associate to this fermion a boson with the creation operator given by Laurent coefficients $B_+(s)$ and destruction operator given by Laurent coefficients of $B_-(s)$, here $B_\pm$ are positive and negative parts of the field $B(s)$ from Section 4.4:

$$\begin{aligned} (B_+(s) F) \xi &= -\frac{\xi'(s)}{\xi(s)} F(\xi(z)), \\ (B_-(s) F) \xi &= -\frac{dF\left(\left(1 + \frac{\tau}{z-s}\right) \xi(z)\right)}{d\tau} \Big|_{\tau=0}, \end{aligned}$$

with Laurent coefficients

$$B_+(s) F(T, t) = - \left(\sum_{l \geq 1} l t_l s^{l-1} \right) F(T, t),$$

$$B_-(s) F(T, t) = \sum_{l \geq 1} \frac{1}{s^{l+1}} \partial_{t_l} F(T, t) + \frac{1}{s} \frac{\partial_T}{T} F(T, t).$$

It is easy to see that $B_+(s) = -\mathbf{E}_s^{\frac{d}{ds}} / \mathbf{E}_s$, $B_-(s) = -\frac{d}{d\tau} \mathbf{M}_{1+\frac{\tau}{z-s}}$. It follows that $B_+(r)$ commutes with $B_+(s)$ without fusions, similarly for $B_-(r)$ and $B_-(s)$, $[B_+(r), B_-(s)] = \frac{1}{(r-s)^2}$, and $B_+(r) B_-(s)$ has no fusion, while the fusions of $B_-(s) B_+(r)$ are residues of $\frac{1}{(r-s)^2}$.

Any pair of fields $B'_+(s)$, $B'_-(s)$ with the same commutation and fusion relations may be considered as fields of creation/destruction of a boson. Since $\alpha(s) = \mathbf{M}_{z-s}$, $\alpha(s)$ and $\alpha(r)$ commute for any r and s without fusion, thus they definitely fulfill the share of properties required from a boson creation field taken separately from boson destruction field. On the other hand, $\alpha(s)^{-1} \frac{d}{ds} \alpha(s) = B_-(s)$! Heuristically, this means that $\alpha(s) = C \exp \int B_-(s) ds$. Thus one should expect that $[B_+(r), \alpha(s)] = \frac{1}{r-s} \alpha(s)$, what is easy to check by a direct calculation. Also, $B_+(r) \alpha(s)$ has no fusion.

It is easy to see that if the integral $\beta(s) = \int B_-(s) ds = \log \alpha(s)$ existed, then the pair $\beta(s)$, $\beta^+(s) = -\frac{d}{ds} B_+(r)$ would satisfy the required properties for a boson destruction/creation fields. Note that $\beta^+(s) = \frac{\left(\mathbf{E}_s^{\frac{d}{ds}} \right)^2 - \mathbf{E}_s \mathbf{E}_s^{\frac{d}{ds} \frac{d}{ds}}}{\mathbf{E}_s^2}$.

Thus we decompose the algebraic boson-fermion correspondence into the following 3 steps

$$B(s) \xrightarrow{\int} \beta(s) \xrightarrow{\exp} \alpha(s) \xrightarrow{\text{Disc}} \psi^+(s),$$

here Disc is an “extrapolation” to Dirac vacuum state of the operation of multiplication by the discriminate Disc (see Section 5.3). As we have seen it above, the first step is not literally defined for operators acting in $\mathfrak{MM}(Z)$, but formally $\beta(s)$ is the field of destruction of a boson (which is “an integral” of algebraic boson $B(s)$). The exponent of this field becomes a creation operator $\alpha(s)$ for a geometric boson, and geometric boson-fermion correspondence expresses a fermion creation operator $\psi^+(s)$ in terms of boson creation operator $\alpha(s)$.

Since the first two operations correspond to integration of a connection, the geometric boson may be considered as a connection, and the algebraic boson as a differential form corresponding to a connection. This fact is widely used in physical literature to quantify connections. I do not know a heuristic which would explain why exponent of β -boson *destruction* operator appears a geometric boson *creation* operator.

REFERENCES

1. Richard E. Borcherds, *Vertex algebras*, <http://xxx.lanl.gov/e-print/q-alg/9706008>, [arXiv:q-alg/9706008](https://arxiv.org/abs/q-alg/9706008), June 1997.
2. Robin Hartshorne, *Algebraic geometry*, Springer-Verlag, New York, 1977, Graduate Texts in Mathematics, No. 52.
3. Victor Kac, *Vertex algebras for beginners*, University Lecture Series, vol. 10, American Mathematical Society, Providence, RI, 1997.
4. Anthony A. Iarrobino, *Punctual Hilbert schemes*, Mem. Amer. Math. Soc. **10** (1977), no. 188, viii+112.
5. M. M. Kapranov, *Chow quotients of Grassmannians. I*, I. M. Gel'fand Seminar, Adv. Soviet Math., vol. 16, Amer. Math. Soc., Providence, RI, 1993, pp. 29–110.
6. Alexander Beilinson and Vladimir Drinfeld, *Chiral algebras*, manuscript.
7. David Mumford, *Lectures on curves on an algebraic surface*, Princeton University Press, Princeton, N.J., 1966, With a section by G. M. Bergman. Annals of Mathematics Studies, No. 59.
8. Maxim Kontsevich, *Deformation quantization of Poisson manifolds, I*, ([arXiv:q-alg/9709040](https://arxiv.org/abs/q-alg/9709040)).
9. M. Kapranov and E. Vasserot, *Vertex algebras and the formal loop space*.
10. Hiraku Nakajima, *Lectures on Hilbert schemes of points on surfaces*, <http://www.kusm.kyoto-u.ac.jp/~nakajima/TeX.html>, October 1996.
11. Andreas Kriegl and Peter W. Michor, *The convenient setting of global analysis*, American Mathematical Society, Providence, RI, 1997. MR **98i**:58015
12. Ilya Zakharevich, *On algebraic geometry of the space of meromorphic functions*, in preparation.
13. I. Grojnowski, *Instantons and affine algebras. I. The Hilbert scheme and vertex operators*, Math. Res. Lett. **3** (1996), no. 2, 275–291.
14. Yi-Zhi Huang and James Lepowsky, *On the $\mathcal{D}$ -module and formal-variable approaches to vertex algebras*, Topics in geometry, Progr. Nonlinear Differential Equations Appl., vol. 20, Birkhäuser Boston, Boston, MA, 1996, pp. 175–202.

E-mail address: math@ilyaz.org